ON THE PROBLEM OF EVANESCENT PROCESSES

Z. DITZIAN AND M. NADKARNI

1. Let R denote the set of real numbers. Let G be a countable dense subgroup of R . We construct a nontrivial σ -finite measure m on R such that

- (i) m is nonatomic, i.e. $\mu(\{x\})=0$ for every real number x ,
- (ii) m is singular with respect to the Lebesgue measure L on R ,
- (iii) m is invariant under translation by members of G , i.e., $m(A+g) = m(A)$ for all $g \in G$ and for all Borel subsets $A \in \mathfrak{B}$. (Here and in sequel $\mathfrak{B}$ denotes the class of Borel subsets of R .)

Such measures are intimately connected with the problem of evanescent processes and analytic functions on compact tori raised by Henry Helson and David Lowdenslager in their paper [3]. We establish this connection in §3. In §2 we shall study the group of unitary operators T^g , $g \in G$, defined on $L_2(R, m)$ by $(T^g f)(\lambda) = f(\lambda + g)$, $f \in G$. In the references we list papers connected with the present work. We proceed with the construction of the measure in steps.

Step 1 (Cantor's decimal set D). Expand every number x in the unit interval $I = \{x = 0 \leq x < 1\}$ in the decimal system, i.e. write $x = \sum_{n=1}^{\infty} (\alpha_n / 10^n)$, $\alpha_n = 0, 1, 2, \dots, 9$; $n = 1, 2, \dots$ and let D be the set of all those numbers x in whose expansion α_n takes values 0 or 9. More accurately D is the set of all numbers x in the unit interval I such that x can be expanded by using 0 and 9 alone. Geometrically D is the Cantor set obtained by deleting the middle 8/10ths.

Step 2. Here we state a known result and indicate its proof. Let A_5 denote the set of all those numbers in the unit interval I whose decimal expansion does not involve the number 5.

LEMMA 1. A_5 has Lebesgue measure zero.

PROOF. Let Q_n be the set of numbers $x \in I$ such that 5 is in the n th decimal place of the expansion of x but not in the first $(n-1)$ places. Then each Q_n is measurable and its Lebesgue measure can be shown to be $9^{n-1}/10^n$. Q_n 's form a disjoint sequence of sets and the Lebesgue measure of $\bigcup_{n=1}^{\infty} Q_n$ is $\sum_{n=1}^{\infty} (9^{n-1}/10^n) = 1$. But $A_5 = I - \bigcup_{n=1}^{\infty} Q_n$. So the Lebesgue measure of A_5 is zero, q.e.d.

Step 3. Let $Q = \{x - y: x, y \in D\}$.

LEMMA 2. Q has Lebesgue measure zero.

PROOF. $Q = Q_+ \cup Q_-$ where

Received by the editors July 18, 1966.

$$Q_+ = \{x - y: x, y \in D, x \geq y\},$$

$$Q_- = \{x - y: x, y \in D, x \leq y\} = -Q_+.$$

So it is enough to show that Q_+ has Lebesgue measure zero. We shall show that $Q_+ \subset A_5$ of Lemma 1. Let $x, y \in D$, with $x > y$, have decimal expansions $\cdot\alpha_1\alpha_2 \cdots$ and $\cdot\beta_1\beta_2 \cdots$ respectively. Let x_n and y_n be the numbers obtained from x, y by terminating their decimal expansions at the n th stage. Then

$$x_n - y_n = \cdot\alpha_1\alpha_2 \cdots \alpha_n - \cdot\beta_1\beta_2 \cdots \beta_n.$$

Since α 's and β 's take values 0 or 9 only (since $x, y \in D$) it follows that $x_n - y_n$ does not involve the number 5 in its decimal expansion. Hence $x - y = \lim_{n \rightarrow \infty} (x_n - y_n)$ does not involve the number 5 in its decimal expansion. So the Lebesgue measure of $Q_+ = 0$. Hence $L(Q) = L(Q_+) + L(Q_-) = 0$. *q.e.d.*

REMARK. It is interesting to note that if D were the well-known Cantor ternary set, then the set $Q = \{x - y: x, y \in D\}$ would be the entire interval from -1 up to 1 .

Step 4. Let Q be the set of Lebesgue measure zero of Step 3. Write $F = \{(x + m)/n: x \in Q, m, n \text{ arbitrary integers, } n \neq 0\}$.

Since Q has Lebesgue measure zero, F has Lebesgue measure zero. Hence there exists an irrational number $\lambda \notin F$.

Step 5. Choose an irrational number $\lambda \notin F$, where F is as in Step 4. Let G be the group $m + \lambda n$, where m, n are integers. The group G is dense in R . (A nondense subgroup of R is necessarily isomorphic to the group of integers.) Let D be the Cantor decimal set of Step 1.

LEMMA 3. *Translates of D by members of G are disjoint.*

PROOF. Let $D + m + \lambda n, D + p + \lambda q$ be two translates of D . Suppose that $(D + m + \lambda n) \cap (D + p + \lambda q) \neq \emptyset$. Then there exists $x, y \in D$ such that $x + m + \lambda n = y + p + \lambda q$, i.e. $x - y = p - m + \lambda(q - n)$. If $q = n$, then $p = m$ (since $0 \leq x, y < 1$) so that we do not have distinct translates. If $q \neq n$, then $(x - y + m - p)/(q - n) = \lambda$; but $(x - y + m - p)/(q - n) \in F$ and $\lambda \notin F$, so we again get a contradiction. Hence translates of D by members of G are disjoint.

Step 6 (*The Cantor function f*). Let $x \in I$ have the decimal expansion $x = \cdot\alpha_1\alpha_2\alpha_3, \cdots, \alpha_i = 0, 1, 2, \cdots, 9$. Let $n = n(x)$ be the first index for which $\alpha_n \in \{1, 2, \cdots, 8\}$ and $\alpha_n \notin \{0, 9\}$. If there is no such n , i.e., if $x \in D$, write $n(x) = \infty$. Define the function f by

$$f(x) = \frac{1}{9} \left(\sum_{i=1}^{n-1} \frac{\alpha_i}{2^i} \right) + \frac{1}{2^n}, \quad n = n(x).$$

The function f is continuous and monotonically nondecreasing with points of increase only in the set D of Lebesgue measure zero.

Step 7 (Construction of m). Let μ be the finite measure induced by the monotone function f of Step 6. μ is obviously nonatomic and singular with respect to the Lebesgue measure on I . Extend μ by setting $\mu(A)=0$ for sets A outside I . Let G be the countable dense subgroup of Step 5 and define m by

$$m(A) = \sum_{n, \lambda_n = -\infty}^{\infty} \mu(A + m + \lambda_n) = \sum_{g \in G} \mu(A + g), \quad A \in \mathfrak{B}.$$

Clearly m is invariant under translation by members of G . Further m is nonatomic. Finally we observe that m is supported on $\bigcup_{g \in G} (D + g)$, the union of countable number of disjoint sets $D + g$, $g \in G$, and that $m(D + g) = m(D) = 1$. Hence m is σ -finite. This completes the construction of m .

REMARK. We have constructed the measure m invariant under translation by a dense subgroup with two generators. But this is not a restriction. With little manipulation one can construct a σ -finite nonatomic singular measure invariant under translation by any countable subgroup of the real line.

2. From now on we shall denote by G a fixed countable dense subgroup of R . A measure m on $\mathfrak{B}$ is called nonatomic singular G -invariant if

- (i) m is nonatomic,
- (ii) m is singular with respect to the Lebesgue measure on R ,
- (iii) $m(A + g) = m(A)$ for all $A \in \mathfrak{B}$ and $g \in G$,
- (iv) There exists a Borel set D of finite m measure such that the translates $D + g$, $g \in G$, of D by members of G are pair wise disjoint and $\bigcup_{g \in G} (D + g)$ supports m .

A method of constructing such measures was given in §1.

Now fix a continuous singular G -invariant measure m on R . Let $L_2(R, m)$ be the linear space of functions square integrable with respect to m . Let D be the set (the existence of which is guaranteed by (iv)) such that the sets $D + g = D_g$ are pair wise disjoint and $\bigcup_{g \in G} D_g$ supports m . Then clearly $L_2(R, m) = \sum_{g \in G} \oplus L_2(D_g, m)$ where $L_2(D_g, m)$ is the set of functions in $L_2(R, m)$ that vanish outside D_g . The orthogonal projection of $f \in L_2(R, m)$ on $L_2(D_g, m)$ is given by fI_g , where I_g is the characteristic function of D_g .

Now m is G -invariant so we get a group T^g , $g \in G$, of unitary operators defined on $L_2(R, m)$ by $(T^g f)(\lambda) = f(\lambda + g)$, $f \in L_2(R, m)$, $g \in G$.

Let us define a spectral measure E on $\mathfrak{B}$ by writing $E(\sigma)f = I_\sigma f$, $f \in L_2(R, m)$ where I_σ is the characteristic function of σ . It is easily

varified that E and T^σ are connected by the relation $T^\sigma E(\sigma) T^{-\sigma} = E(\sigma - g)$ for all $g \in G$. But T^σ is not the only commutative group of unitary operators which satisfies this equation with E . The general commutative group U^σ , which with E , satisfies $U^\sigma E(\sigma) U^{-\sigma} = E(\sigma - g)$, $g \in G$, has the following form. U^σ is defined by

$$(U^\sigma f)(\lambda) = A(g, \lambda) f(\lambda + g), \quad g \in G, f \in L_2(R, m),$$

where $A(g, \lambda)$ is an m -measurable function of λ for every fixed g such that

- (i) $|A(g, \lambda)| = 1$,
- (ii) $A(g+h, \lambda) = A(g, \lambda) A(h, \lambda+g)$ for almost all λ with respect to the m -measure.

The set of m -measure zero where (ii) does not hold may vary with the pair (g, h) .

Functions satisfying the functional equation (ii) occur very crucially in the study of spectral measures E on $\mathfrak{B}$ for which there exists a commutative group U^σ , $g \in G$, satisfying the equation $U^\sigma E(\sigma) U^{-\sigma} = E(\sigma + g)$, $g \in G$, $\sigma \in \mathfrak{B}$ (cf. §4).

The group U^σ has a spectral measure associated with it as follows. (See [6, p. 392].)

Let $B = \hat{G}_d$ be the compact dual of G_d , the group G with the discrete topology. Since U^σ is a commutative group of unitary operators, by Godement's extension of Stone's theorem on the representation of unitary operators [1] there exists a Hermitian projection valued spectral measure F on the Borel subsets $\mathfrak{F}$ of B such that $U^\sigma = \int_B \chi_\sigma(\lambda) dF_\lambda$ in the sense that

$$(*) \quad (U^\sigma f, h) = \int_B \chi_\sigma(\lambda) (dF_\lambda f, h), \quad f, h \in L_2(R, m).$$

Here χ_σ denotes the character on B corresponding to $g \in G_d$. For $f, h \in L_2(R, m)$, $(F(\cdot)f, h)$ defines a complex valued finite measure on $\mathfrak{F}$ so that for $\sigma \in \mathfrak{F}$ the value of this measure is $(F(\sigma)f, h)$.

We show that for every f and h , $(F(\cdot)f, h)$ is absolutely continuous with respect to the Haar measure on B .

THEOREM 1. *If $f \in L_2(D_{g_0}, m)$ for some $g_0 \in G$, then the measure $(F(\cdot)f, f)$ is a constant multiple of the Haar measure on B . For any $f, h \in L_2(R, m)$, the measure $(F(\cdot)f, h)$ is absolutely continuous with respect to the Haar measure on B .*

PROOF. Let $f \in L_2(D_{g_0}, m)$, then $U^\sigma f \in L_2(D_{g_0-g}, m)$. Hence, the elements $\{U^\sigma f: g \in G\}$ are mutually orthogonal. Now by $(*)$ $(U^\sigma f, f) = \int_B \chi_\sigma(\lambda) (dF_\lambda f, f) = 0$ if $g \neq 0$. Hence $(F(\cdot)f, f)$ is a constant multiple

of the Haar measure on B . The constant multiple is, of course, non-zero if and only if $f \neq 0$ in $L_2(D_{g_0}, m)$. Now let $f \in L_2(D_{g_0}, m)$, $h \in L_2(D_{g_1}, m)$, then by the polarization formula it is easy to see that $(F(\cdot)f, h)$ is absolutely continuous with respect to the Haar measure on B . Finally choose any $f, h \in L_2(R, m)$. Let $f = \sum_{g \in G} f_g$, $h = \sum_{g \in G} h_g$, $f_g, h_g \in L_2(D_g, m)$. Then clearly

$$(F(\cdot)f, h) = \sum_{g, g' \in G} (F(\cdot)f_g, h_{g'}).$$

Since each $(F(\cdot)f_g, h_{g'})$ is absolutely continuous with respect to the Haar measure on B , it follows that $(F(\cdot)f, h)$ has the same property, q.e.d.

The next theorem shows that Wiener closure theorem has no analogue for a nonatomic singular G -invariant measure.

THEOREM 2. *There is no $f \in L_2(R, m)$ such that $\{U^g f \cdot g \in G\}$ spans $L_2(R, m)$.*

To prove this theorem we need a known result which we state here without proof for the sake of completeness.

LEMMA 4. *Let μ be a finite positive regular measure on the Borel subsets $\mathfrak{F}$ of B . Let $h, f \in L_2(B, \mu)$. Then $\int_B \chi_g(\lambda) h(\lambda) \overline{f(\lambda)} d\mu = 0$ for all $g \in G_d$ if and only if h vanishes almost everywhere with respect to μ on the set where $|f| > 0$.*

This lemma is an easy consequence of the fact that a finite regular Borel measure on a locally compact abelian group is uniquely determined by its Fourier-Stieltjes transform [7, p. 17].

Consider the measures on $\mathfrak{F}$ defined by $\int_{\sigma} |h(\lambda)|^2 d\mu$, $\int_{\sigma} |f(\lambda)|^2 d\mu$. Then the lemma is equivalent to the following fact:

$$\int_B \chi_g(\lambda) h(\lambda) \overline{f(\lambda)} d\mu = 0$$

for all $g \in G_d$ if and only if the measures $\int_{\cdot} |h(\lambda)|^2 d\mu$ and $\int_{\cdot} |f(\lambda)|^2 d\mu$ are mutually singular.

PROOF OF THEOREM 2. Suppose that there exists $f \in L_2(R, m)$ such that $\{U^g f \cdot g \in G\}$ spans $L_2(R, m)$. By (*), $(U^g f, f) = \int \chi_g(\lambda) (dF_{\lambda} f, f) = \int \chi_g(\lambda) d\mu$, where μ is the measure defined by $\mu(\sigma) = (F(\sigma)f, f)$, $\sigma \in \mathfrak{F}$. By Theorem 1, μ is absolutely continuous with respect to the Haar measure on B . The mapping $S: SU^g f = \chi_g$ extends by linearity to an invertible isometry from the space spanned by $\{U^g f: g \in G\}$ to $L_2(B, \mu)$. Now let D_1, D_2 be two disjoint measureable subsets of D

such that $D = D_1 \cup D_2$ and $m(D_1), m(D_2) > 0$. Let h_1 and h_2 denote the characteristic functions of D_1 and D_2 . Write $f_1 = Sh_1$, $f_2 = Sh_2$. It is clear that

- (i) $U^g h_1$ are all mutually orthogonal in $L_2(R, m)$,
- (ii) $U^g h_2$ are all mutually orthogonal in $L_2(R, m)$,
- (iii) $U^g h_1 \perp U^{g'} h_2$ for all $g, g' \in G$.

This is because the translates of D by members of G are disjoint.

Now for all $g \in G$, $(U^g h_1, h_1) = \int \chi_g(\lambda) |f_1(\lambda)|^2 d\mu = \int_B \chi_g(\lambda) (dF_\lambda h_1, h_1) = 0$ if $g \neq 0$. Similarly $(U^g h_2, h_2) = \int \chi_g(\lambda) |f_2(\lambda)|^2 d\mu = \int_B \chi_g(\lambda) (dF_\lambda h_2, h_2) = 0$.

Hence the measures $(F(\cdot)h_1, h_1)$ and $(F(\cdot)h_2, h_2)$ are nonzero constant multiples of the Haar measure on B . Hence the measures $\int(\cdot) |f_1(\lambda)|^2 d\mu$ and $\int(\cdot) |f_2(\lambda)|^2 d\mu$ are nonzero constant multiples of the Haar measure on B . But $U^g h_1 \perp U^{g'} h_2$ for all g, g' by (iii). Hence $(U^g h_1, h_2) = \int_B \chi_g(\lambda) \bar{f}_2 d\mu = 0$ for all g . Hence by Lemma 4 the measures $\int(\cdot) |f_1(\lambda)|^2 d\mu$ and $\int(\cdot) |f_2(\lambda)|^2 d\mu$ are mutually singular. This is a contradiction, q.e.d.

3. In this section we show how the measures of the type discussed in §2 are excluded in the problem of evanescent processes. First we must explain this problem.

Let G and B be as in §2. Let f be a nonzero positive function on B summable with respect to the Haar measure on B . Let $L_2(B, f) = \{\psi: |\psi|^2 f \text{ is summable with respect to the Haar measure on } B\}$. Let H_t be the subspace of $L_2(B, f)$ spanned by $\{\chi_g: g < t\}$, where χ_g denotes the character on B corresponding to the real number $g \in G$. It is clear that $H_t \subseteq H_{t'}$ whenever $t < t'$. It can be shown that either $H_t = H_{t'}$ for all t, t' or $\bigcap_t H_t = \{0\}$ and $H_t \subsetneq H_{t'}$ whenever $t < t'$. This has been shown by Helson and Lowdenslager in their paper [3]. The problem of evanescent processes can be stated as follows: Assume that $H_t \neq H_{t'}$ for $t < t'$, then is it always true that $(\bigcap_{t>0} H_t) \ominus H_0 \neq \{0\}$?

A well-known result of Helson and Lowdenslager [2] answers the question in the affirmative under the assumption that $\log f$ is summable with respect to the Haar measure on B . In what follows we give further evidence in favor of the affirmative answer to the question.

The increasing subspaces H_t give rise to a spectral measure E on the Borel subsets of R . For intervals $(a, b]$, E is given by $E(a, b] =$ orthogonal projection on $H_b \ominus H_a$. In $L_2(B, f)$ there is a commutative group U^g of unitary operators defined by $U^g \psi = \chi_g \psi$, $\psi \in L_2(B, f)$, $g \in G$. Further the following two identities are easily verified

(A) $U^g (H_b \ominus H_a) = H_{b+g} \ominus H_{a+g}$ where a, b ($a < b$) are any two real numbers.

(B) For any $\psi \in L_2(B, f)$, $\|E(a, b]\psi - \psi\|^2 = \|U^g E(a, b]\psi - U^g \psi\|^2$.

(A) and (B) together imply that U^σ and E are connected by the relation $U^\sigma E(\sigma) U^{-\sigma} = E(\sigma + g)$ for all $\sigma \in \mathfrak{B}$ and $g \in G$. Helson and Lowdenslager have shown that if $E\{x\} \neq 0$ for some x , then the spectral measure E is purely discrete and E has no continuous component. Now it can be shown that E cannot have a component absolutely continuous with respect to the Lebesgue measure on R , i.e., there does not exist a nonzero $\psi \in L_2(B, f)$ such that $(E(\cdot)\psi, \psi)$ is absolutely continuous with respect to the Lebesgue measure on R . In what follows we show that E has no component absolutely continuous with respect to a nonatomic singular G -invariant measure on R .

THEOREM 3. *Assume that $E\{x\} = 0$ for all x . There does not exist a Borel set D such that:*

- (i) *the sets $D+g$, $g \in G$ are mutually disjoint,*
- (ii) *$E(D) \neq 0$.*

PROOF. Suppose not. Then there exists a set D such that the sets $D+g$, $g \in G$, are mutually disjoint and $E(D) \neq 0$. Since E has no discrete spectrum, we can find two nonzero vectors Φ, ψ in $E(D)$ such that Φ and ψ are mutually orthogonal. Now $U^\sigma \Phi = U^\sigma E(D)\Phi = E(D+g)U^\sigma \Phi \in E(D+g)$ and similarly $U^\sigma \psi \in E(D+g)$. Since the sets $D+g$, $g \in G$, are mutually disjoint, we see that $U^\sigma \Phi \perp \Phi$, $U^\sigma \psi \perp \psi$ for all $g \neq 0$ and $U^\sigma \Phi \perp U^\sigma \psi$ for all g, g' . So

- (i) $(U^\sigma \Phi, \Phi) = \int_B \chi_g(\lambda) |\Phi(\lambda)|^2 f(\lambda) d\sigma = 0$ for $g \neq 0$.
 - (ii) $(U^\sigma \psi, \psi) = \int_B \chi_g(\lambda) |\psi(\lambda)|^2 f(\lambda) d\sigma = 0$ for $g \neq 0$.
 - (iii) $(U^\sigma \Phi, \psi) = \int_B \chi_g(\lambda) \Phi(\lambda) \overline{\psi(\lambda)} f(\lambda) d\sigma = 0$ for all g .
- (Here $d\sigma$ is the normalized Haar measure on B .)

The first two equations above say that $|\Phi|^2 f d\sigma$ and $|\psi|^2 f d\sigma$ are nonzero constant multiples of the Haar measure on B and the third equation says that $\Phi \bar{\psi} f$ is equal to zero almost everywhere with respect to the Haar measure on B . This is impossible, q.e.d.

4. Let E be a spectral measure on the Borel subsets of R and let G be a countable dense subgroup of R . We call a spectral measure E G -stationary if there exists a commutative group U^σ of unitary operators such that $U^\sigma E(\sigma) U^{-\sigma} = E(\sigma + g)$ for all $\sigma \in \mathfrak{B}$ and $g \in G$. If one tries to obtain the canonical representation of G -stationary spectral measures like the one there is for a pair of commutative groups of unitary operators satisfying Weyl's commutativity relation one at once faces the following question.

Let μ be a finite positive measure on $\mathfrak{B}$. Call μ G -quasi invariant if μ and μ_g are mutually absolutely continuous for all $g \in G$. Here μ_g is defined by $\mu_g(A) = \mu(A+g)$, $A \in \mathfrak{B}$, $g \in G$.

QUESTION 1. μ is G -quasi invariant. Does there exist a σ -definite measure m on $\mathfrak{B}$ such that (i) $m(\sigma+g)=m(\sigma)$ for all $\sigma\in\mathfrak{B}$, $g\in G$, (ii) m and μ are mutually absolutely continuous?

Now suppose that $\mu=\mu^d+\mu^a+\mu^s$ where μ^d is the atomic part of μ , μ^a =part of μ absolutely continuous with respect to the L , the Lebesgue measure, and μ^s =nonatomic singular part of μ . It is easy to see that each component μ^d , μ^a and μ^s is separately G -quasi invariant.

Further μ^a and the Lebesgue measure are mutually absolutely continuous. Thus for μ^a the question raised above has a solution. One can also show easily that the question raised above has a solution for μ^d . Hence in the question raised above one can assume that μ is nonatomic singular measure.

We give a reformulation of our question in terms of the functions $A(g, \lambda)=(d\mu_g/d\mu)(\lambda)$. One verifies very easily that $A(g, \lambda)$ satisfy the relation $A(g+h, \lambda)=A(g, \lambda)A(h, \lambda+g)$ a.e. $[\mu]$.

THEOREM 4. *Question 1 has a solution if and only if there exists a measurable function B such that $A(g, \lambda)=B(\lambda+g)/B(\lambda)$.*

PROOF. Suppose there exists an m as in Question 1. Write $B(\lambda)=(d\mu/dm)(\lambda)$. Then clearly $A(g, \lambda)=(d\mu_g/d\mu)(\lambda)=(d\mu_g/dm)(\lambda)\cdot(dm/d\mu)(\lambda)=(d\mu_g/dm)(\lambda)\cdot 1/B(\lambda)$. Now by the invariance of m under translation by g it is easy to see that $(d\mu_g/dm)(\lambda)=B(\lambda+g)$; thus $A(g, \lambda)=B(\lambda+g)/B(\lambda)$. Conversely suppose that $A(g, \lambda)=B(\lambda+g)/B(\lambda)$ where B is measurable. Define m by $m(\sigma)=\int_{\sigma}[B(\lambda)]^{-1}d\mu$. It is clear that m and μ are mutually absolutely continuous. Next to see the G -invariance of m we note that

$$\begin{aligned} m(\sigma+g) &= \int_{\sigma+g} [B(\lambda)]^{-1}d\mu = \int_{\sigma} [B(\lambda+g)]^{-1}d\mu_g(\lambda) \\ &= \int_{\sigma} [B(\lambda+g)]^{-1} \frac{d\mu_g}{d\mu}(\lambda) d\mu = \int_{\sigma} \frac{[B(\lambda+g)]^{-1}B(\lambda+g)}{B(\lambda)} d\mu \\ &= \int_{\sigma} [B(\lambda)]^{-1}d\mu = m(\sigma), \end{aligned} \quad \text{q.e.d.}$$

We conclude by making the following remarks.

Assume that μ of Question 1 is singular. In order that Question 1 have an affirmative solution it is enough that there is a μ -measurable set D such that $D+g$, $g\in G$ are disjoint and $\bigcup_{g\in G}(D+g)$ supports μ . However, there exist singular G -quasi invariant measures for which no such D exists. We illustrate this by the following example. Let C

be the Cantor ternary set and ψ the Cantor function from C onto $[0, 1]$. ψ is strictly increasing and continuous on C with range $[0, 1]$. Let P be the singular measure on the real line induced by ψ . Let G be the group of real members having finitely many terms in their ternary expansions. Let $g_1, g_2, g_3, \dots$ be a denumeration of G . Write $\mu(A) = \sum_{n=1}^{\infty} (1/2^n) P(A + g_n)$, $A \in \mathfrak{B}$. Clearly μ is G -quasi invariant. Call two members of C equivalent if their difference belongs to G . This equivalence relation partitions C . Choose a member from each equivalence class and call the new set D . Translates $D + g$, $g \in G$ are disjoint and $\bigcup_{g \in G} (D + g)$ supports μ . But D can never be chosen to be μ -measurable, for the difference of two members of $\psi(D)$ has always finite binary expansions, so that $\psi(D)$ is nonmeasurable. Hence $D = \psi^{-1}(\psi(D))$ is non- μ -measurable.

REFERENCES

1. R. Godement, *Sur une généralisation d'un théorème de Stone*, C.R. Acad. Sci. Paris **218** (1944), 901–903.
2. H. Helson and D. Lowdenslager, *Prediction theory and Fourier series in several variables*. I, Acta Math. **99** (1958), 165–202.
3. ———, *Prediction theory and Fourier series in several variables*. II, Acta Math. **106** (1961), 175–213.
4. ———, *Invariant subspaces*, Proc. Internat. Sympos. on Linear Spaces, Jerusalem, 1960, pp. 251–262, Macmillan (Pergamon), New York, 1961.
5. George Mackey, *A theorem of Stone and von Neumann*, Duke Math. J. **16** (1949), 313–326.
6. F. Riesz and B. Sz. Nagy, *Functional analysis*, Ungar, New York, 1955.
7. W. Rudin, *Fourier analysis on groups*, Interscience, New York, 1962.

MICHIGAN STATE UNIVERSITY AND
UNIVERSITY OF MINNESOTA