A NOTE ON REFLEXIVE BANACH SPACES

KUNG-WEI YANG

The purpose of this note is to prove three well-known theorems concerning reflexive Banach spaces by using exact sequences.

Let F denote either the field of real numbers or the field of complex numbers. Let $\mathcal{B}$ be the category whose objects are Banach spaces over the field F and whose morphisms are continuous linear maps $T: X \rightarrow Y$. As usual, $\mathcal{B}(X, Y)$ denotes the set of all continuous linear maps from X to Y . With the norm $|T| = \sup_{|x| \leq 1} |T(x)|$ for each T in $\mathcal{B}(X, Y)$, $\mathcal{B}(X, Y)$ is a Banach space over F . Notice that if $T: X \rightarrow Y$ and $S: Y \rightarrow Z$ then $|ST| \leq |S| |T|$. This implies that

$$\mathcal{B}(T, Z): \mathcal{B}(Y, Z) \rightarrow \mathcal{B}(X, Z)$$

is a morphism in the category $\mathcal{B}$ and $|\mathcal{B}(T, Z)| \leq |T|$. As in [1], we let $\mathcal{B}(X, F) = X^*$ and $\mathcal{B}(T, F) = T^*$.

Let Y be a closed subspace of the Banach space X . Let $i: Y \rightarrow X$ be the inclusion map. Then

$$0 \rightarrow Y \xrightarrow{i} X \xrightarrow{p} Z \rightarrow 0,$$

where $Z = X/Y$ and p is the natural homomorphism, is an exact sequence in $\mathcal{B}$. By the Hahn-Banach Theorem, the sequence

$$0 \leftarrow Y^* \xleftarrow{i^*} X^* \xleftarrow{p^*} Z^* \leftarrow 0$$

is exact. Therefore the sequence

$$0 \longrightarrow Y^{**} \xrightarrow{i^{**}} X^{**} \xrightarrow{p^{**}} Z^{**} \longrightarrow 0$$

is also exact. Clearly, we have the following commutative diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & Y & \xrightarrow{i} & X & \xrightarrow{p} & Z \rightarrow 0 \\ & & n_1 \downarrow & & n_2 \downarrow & & n_3 \downarrow \\ 0 & \rightarrow & Y^{**} & \xrightarrow{i^{**}} & X^{**} & \xrightarrow{p^{**}} & Z^{**} \longrightarrow 0 \end{array}$$

Presented to the Society, February 15, 1967 under the title *Exact sequences and reflexive Banach spaces*; received by the editors September 12, 1966.

where n_1, n_2, n_3 are the natural embeddings and all the rows and columns are exact. Therefore, by "diagram chasing," we have the following commutative diagram (D):

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & Y & \xrightarrow{i} & X & \xrightarrow{p} & Z \longrightarrow 0 \\
 & & n_1 \downarrow & & n_2 \downarrow & & n_3 \downarrow \\
 (D) \quad 0 & \longrightarrow & Y^{**} & \xrightarrow{i^{**}} & X^{**} & \xrightarrow{p^{**}} & Z^{**} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & Y^{**}/Y & \longrightarrow & X^{**}/X & \longrightarrow & Z^{**}/Z \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

where all the rows and columns are exact. In particular, the following sequence (E) is exact.

$$(E) \quad 0 \rightarrow Y^{**}/Y \rightarrow X^{**}/X \rightarrow Z^{**}/Z \rightarrow 0.$$

We observe that the Banach space X is reflexive if and only if the Banach space X^{**}/X in (D) (or (E)) is equal to 0.

THEOREM 1. *If X is a reflexive Banach space and Y is a closed subspace of X , then Y is reflexive.*

PROOF. By the exactness of the sequence (E), we have X is reflexive $\Rightarrow X^{**}/X = 0 \Rightarrow Y^{**}/Y = 0 \Rightarrow Y$ is reflexive.

THEOREM 2. *If X is a Banach space and Y is a closed subspace of X , and if both Y and X/Y are reflexive, then X is reflexive.*

PROOF. Again by the exactness of the sequence (E), we have Y and X/Y are reflexive $\Rightarrow Y^{**}/Y = 0$ and $Z^{**}/Z = 0 \Rightarrow X^{**}/X = 0 \Rightarrow X$ is reflexive.

THEOREM 3. *If X is a reflexive Banach space and Y is a closed subspace of X , then X/Y is reflexive.*

PROOF. By the exactness of the sequence (E), we have X is reflexive $\Rightarrow X^{**}/X = 0 \Rightarrow Z^{**}/Z = 0 \Rightarrow Z = X/Y$ is reflexive.

REFERENCE

1. N. Dunford and J. T. Schwartz, *Linear operators*, Vol. I, Interscience, New York, 1958.

WESTERN MICHIGAN UNIVERSITY

ON TYPE I C^* -ALGEBRAS

SHÔICHIRO SAKAI¹

1. Introduction. Recently, the author [4] proved the equivalence of type I C^* -algebras and GCR C^* -algebras without the assumption of separability. On the other hand, for separable type I C^* -algebras, we have a simpler criterion as follows: a separable C^* -algebra $\mathfrak{A}$ is of type I if and only if every irreducible image contains a nonzero compact operator.

It has been open whether or not this remains true when $\mathfrak{A}$ is not separable (cf. [1], [2], [3]).

In the present paper, we shall show that a C^* -algebra $\mathfrak{A}$ is GCR if and only if every irreducible image contains a nonzero compact operator, so that by the author's previous theorem [4], the above problem is affirmative for arbitrary C^* -algebra.

2. Theorem. In this section, we shall show the following theorem.

THEOREM. *A C^* -algebra $\mathfrak{A}$ is of type I if and only if every irreducible image contains a nonzero compact operator.*

PROOF. Suppose that a C^* -algebra $\mathfrak{A}$ is of type I, then it is GCR and so every irreducible image contains a nonzero compact operator (cf. [1], [2], [3], [4]).

Conversely suppose that every irreducible image of $\mathfrak{A}$ contains a nonzero compact operator. It is enough to assume that $\mathfrak{A}$ has the unit I . We shall assume that $\mathfrak{A}$ is not of type I. Then it is not GCR; then there is a separable nontype I C^* -subalgebra $\mathfrak{B}$ of $\mathfrak{A}$ (cf. [2], [4]). Take a pure state ϕ on $\mathfrak{B}$ such that the image of $\mathfrak{B}$ under the irreducible $*$ -representation $\{U_\phi, \mathfrak{H}_\phi\}$ of $\mathfrak{B}$ constructed via ϕ does not contain any nonzero compact operator, where $\mathfrak{H}_\phi$ is a Hilbert space.

Received by the editors September 12, 1966.

¹ This paper was written with partial support from ONR Contract NR-551(57).