NTH ROOTS OF OPERATORS

M. R. EMBRY

I. Introduction. Let A and E be continuous linear transformations which map a Banach space into itself. If E commutes with A^n , for some integer n greater than 1, we do not necessarily know whether E commutes with A . The main purpose of this paper is to exhibit a proof of the following theorem:

Let A and E be continuous linear transformations which map a complex Banach space into itself. Let n be an integer greater than 1. If $\sigma(A) \cap \sigma(e^{2\pi i k/n} A) = \emptyset$ for $k=1, \dots, n-1$ and E commutes with A^n , then E commutes with A .

A significant corollary of this theorem is as follows:

If A is a continuous linear transformation from a Hilbert space into itself and A^n is normal, then A is normal, whenever the spectrum of A satisfies the condition stated in the theorem.

Throughout the paper the following notation and terminology will be used. $\mathfrak{X}$ is a complex Banach space and $\mathfrak{L}(\mathfrak{X})$ is the space of continuous linear transformations from $\mathfrak{X}$ into $\mathfrak{X}$. The identity element of $\mathfrak{L}(\mathfrak{X})$ will be denoted by I . If $A \in \mathfrak{L}(\mathfrak{X})$, $\sigma(A)$ will denote the spectrum of A and $\rho(A)$ the resolvent set of A . If C is a simple closed rectifiable curve in the complex plane, C will be assumed to be oriented in a counterclockwise direction. By the interior of C we mean the bounded component of the complement of C ; the exterior of C is the unbounded component.

II. Essential background material. The proof of the above-stated theorem relies heavily on the following results, which are found in *The spectrum of linear transformations* by E. R. Lorch [1]:

THEOREM 1. *Let C be a simple closed rectifiable curve lying entirely within $\rho(A)$, where $A \in \mathfrak{L}(\mathfrak{X})$. Then the contour integral*

$$P = \frac{1}{2\pi i} \int_C (zI - A)^{-1} dz$$

exists and represents an element of $\mathfrak{L}(\mathfrak{X})$.

The transformation P is unchanged if the curve C is continuously

Received by the editors July 22, 1966 and, in revised form, December 8, 1966.

deformed into a curve C' , providing only that the deformation is effected without going outside of $\rho(A)$ [1, p. 241].

THEOREM 2. *The operator P defined by the curve C as in Theorem 1 has the following properties:*

- (a) $P=0$ if and only if every point in the interior of C lies in $\rho(A)$.
- (b) $P=I$ if and only if every point in the exterior of C lies in $\rho(A)$ [1, p. 244].

III. Basic lemmas. It is well known that if $A \in \mathfrak{L}(\mathfrak{X})$, $\sigma(A)$ is a nonempty, compact subset of the complex plane. If for some integer n , $n > 1$, $\sigma(A) \cap \sigma(e^{2\pi ik/n}A) = \emptyset$ for $k=1, \dots, n-1$, then $M > 0$, where

$$M = \min_{k=1, \dots, n-1} \text{g.l.b.} \{ |z - w| : z \in \sigma(A), w \in \sigma(e^{2\pi ik/n}A) \}.$$

Since $\sigma(A)$ is compact, there exist $z_1, \dots, z_m$ in $\sigma(A)$ such that $\sigma(A) \subset \bigcup_{j=1}^m B(z_j; M/3)$, where $B(z; M/3)$ is the open ball in the complex plane of radius $M/3$ and center z . Let $V_1, \dots, V_p$ be the components of $\bigcup_{j=1}^m B(z_j; M/3)$ and let

$$s_{p(k-1)+j} = \sigma(e^{2\pi ik/n}A) \cap (e^{2\pi ik/n}V_j)$$

for $k=1, \dots, n$ and $j=1, \dots, p$. Note the following facts:

- (1) for each j , s_j is a compact subset of the complex plane;
- (2) for $q \neq j$, $s_q \cap s_j = \emptyset$;
- (3) for $k=1, \dots, n$, $\sigma(e^{2\pi ik/n}A) = \bigcup_{j=1}^p s_{p(k-1)+j}$.

Let $S = \{s_k : k=1, \dots, np\}$. We will say that s_j is *interior* to s_q if $j \neq q$ and whenever C is a simple closed rectifiable curve such that s_q is in the interior of C , then s_j is also in the interior of C . We will denote this by $s_j < s_q$. Because of the way in which S was constructed, if $s_j < s_q$, there exists a simple closed rectifiable curve C' such that s_j is in the interior of C' and s_q is in the exterior of C' .

C will be called a *proper curve* for s_j if C is a simple closed rectifiable curve, lying entirely in $\bigcap_{k=1}^n \rho(e^{2\pi ik/n}A)$, such that (1) s_j is in the interior of C and (2) whenever s_q is in the interior of C , then $s_q = s_j$ or $s_q < s_j$.

An element s_k of S will be said to be of *order* 0 if there exists no s_j in S such that $s_j < s_k$. Since S is a finite set, there must exist elements of S of order 0. A *branch of length* k in S is a sequence $\{t_j : j=1, \dots, k\}$ of elements of S such that (1) t_1 is of order 0, (2) $t_j < t_{j+1}$ for $j=1, \dots, k-1$, and (3) there exists no s_i in S such that $t_j < s_i < t_{j+1}$. An element s_i of S is of *order* k , $k \geq 1$, if there exists a branch $\{t_j : j=1, \dots, k\}$ of length k in S such that $t_k < s_i$ and there

exists no branch $\{\bar{i}_j: j=1, \dots, k+1\}$ of length $k+1$ in S such that $\bar{i}_{k+1} < s_i$.

LEMMA 1. Let n be a positive integer. There exist nonzero complex numbers $b_1, \dots, b_n$ such that

$$(z^n I - A^n)^{-1} = A^{-(n-1)} \sum_{j=1}^n b_j (zI - e^{2\pi i j/n} A)^{-1},$$

whenever A is an invertible element of $\mathfrak{L}(\mathfrak{X})$ and $z \in \bigcap_{j=1}^n \rho(e^{2\pi i j/n} A)$.

PROOF. Let $\{b_1, \dots, b_n\}$ be the unique solution of the system of equations

$$\begin{aligned} \sum_{k=1}^n b_k e^{2\pi i k(j-1)/n} &= 0 \quad \text{for } j = 1, \dots, n-1, \\ \sum_{k=1}^n b_k e^{2\pi i k(n-1)/n} &= 1. \end{aligned}$$

Let z be any element of $\bigcap_{j=1}^n \rho(e^{2\pi i j/n} A)$. Then

$$\begin{aligned} A^{n-1} &= \sum_{j=1}^n z^{n-j} A^{j-1} \left(\sum_{k=1}^n b_k e^{2\pi i k(j-1)/n} \right) \\ &= \sum_{k=1}^n b_k \left(\sum_{j=1}^n z^{n-j} (e^{2\pi i k/n} A)^{j-1} \right) \\ &= \sum_{k=1}^n b_k (zI - e^{2\pi i k/n} A)^{-1} (z^n I - A^n) \\ &= (z^n I - A^n) \sum_{k=1}^n b_k (zI - e^{2\pi i k/n} A)^{-1}. \end{aligned}$$

Therefore, $(z^n I - A^n)^{-1} = A^{-(n-1)} \sum_{k=1}^n b_k (zI - e^{2\pi i k/n} A)^{-1}$.

To see that each of the b_j is nonzero we need only consider a special case. Let $A = I$. Let C be a simple closed rectifiable curve such that $e^{2\pi i j/n}$ is in the interior of C and $e^{2\pi i k/n}$, $k \neq j$, is in the exterior of C . Then, using the last equation in the preceding paragraph, we see that

$$\begin{aligned} \frac{1}{2\pi i} \int_C (z^n - 1)^{-1} dz &= \sum_{k=1}^n b_k \frac{1}{2\pi i} \int_C (z - e^{2\pi i k/n})^{-1} dz \\ &= b_j. \end{aligned}$$

However,

$$\frac{1}{2\pi i} \int_C (z^n - 1)^{-1} dz = \prod_{k=1; k \neq j}^n (e^{2\pi i j/n} - e^{2\pi i k/n})^{-1} \neq 0.$$

Therefore, $b_j \neq 0$, $j=1, \dots, n$.

The most serious problem which arose in the proof of Theorem 3 was the interrelation between the components of $\sigma(A)$ and the components of $\sigma(e^{2\pi ik/n}A)$, $k=1, \dots, n-1$. If there exists a simple closed rectifiable curve C such that $\sigma(A)$ is interior to C and $\sigma(e^{2\pi ik/n}A)$ is exterior to C for $k=1, \dots, n-1$, then by Theorem 2 we have

$$\int_C (zI - e^{2\pi ik/n}A)^{-1} dz = 0 \quad \text{for } k = 1, \dots, n-1$$

and $(1/2\pi i) \int_C (zI - A)^{-1} dz = I$. Using Lemma 1, we see that

$$\frac{1}{2\pi i} \int_C (z^n I - A^n)^{-1} dz = A^{-(n-1)} b_n.$$

Therefore, if E commutes with A^n , E also commutes with $A^{-(n-1)}$ and with A . However, in the case in which there exists no such curve C , the situation is much more complicated.

LEMMA 2. *Let A and E be elements of $\mathfrak{L}(\mathfrak{X})$ and let n be an integer greater than 1. Suppose that*

$$\sigma(A) \cap \sigma(e^{2\pi ik/n}A) = \emptyset \quad \text{for } k = 1, \dots, n-1$$

and that E commutes with A^n . If s_j is an element of S and C is a proper curve for s_j , then E commutes with

$$A^{-(n-1)} \int_C (zI - e^{2\pi ik/n}A)^{-1} dz, \quad k = 1, \dots, n.$$

PROOF. The proof of this lemma will be by induction on the order of the elements of S . Note that if C is any simple closed rectifiable curve, lying in $\bigcap_{k=1}^n \rho(e^{2\pi ik/n}A)$, then $\int_C (z^n I - A^n)^{-1} dz$ is an element of $\mathfrak{L}(\mathfrak{X})$ and commutes with E [1, pp. 240-241].

Let s_j be an element of S of order 0 and let C be a proper curve for s_j . Assume that $s_j \subset \sigma(e^{2\pi ik/n}A)$. By Theorem 2,

$$(1) \quad \int_C (zI - e^{2\pi iq/n}A)^{-1} dz = 0 \quad \text{for } q \neq k \quad \text{and } q = 1, \dots, n.$$

Using (1) and Lemma 1, we have

$$(2) \quad \int_C (z^n I - A^n)^{-1} dz = b_k A^{-(n-1)} \int_C (zI - e^{2\pi ik/n}A)^{-1} dz.$$

From (1) and (2) it follows that E commutes with

$$A^{-(n-1)} \int_C (zI - e^{2\pi iq/n}A)^{-1} dz \quad \text{for } q = 1, \dots, n$$

whenever C is a proper curve for an element of S of order 0.

Assume now that whenever s_j is of order p , $0 \leq p \leq m$, and C is a proper curve for s_j , E commutes with

$$A^{-(n-1)} \int_C (zI - e^{2\pi i q/n} A)^{-1} dz \quad \text{for } q = 1, \dots, n.$$

Let s_{j_0} be an element of S of order $m+1$ and C be a proper curve for s_{j_0} . Let $t_1, \dots, t_r$ be the elements of S such that $t_j < s_{j_0}$, $j=1, \dots, r$ and there exists no s_k in S such that $t_j < s_k < s_{j_0}$. Let C_j , $j=1, \dots, r$, be a proper curve for t_j . Since each of the t_j , $j=1, \dots, r$, is of order p , $p \leq m$, then E commutes with

$$A^{-(n-1)} \int_{C_j} (zI - e^{2\pi i k/n} A)^{-1} dz \quad \text{for } k = 1, \dots, n \text{ and } j = 1, \dots, r.$$

Assume that $s_{j_0} \subset \sigma(e^{2\pi i k/n} A)$. It follows from Theorems 1 and 2 that for $q \neq k$,

$$A^{-(n-1)} \int_C (zI - e^{2\pi i q/n} A)^{-1} dz = \sum_{p=1}^r A^{-(n-1)} \int_{C_p} (zI - e^{2\pi i q/n} A)^{-1} dz.$$

Therefore, the induction hypothesis implies that E commutes with

$$A^{-(n-1)} \int_C (zI - e^{2\pi i q/n} A)^{-1} dz \quad \text{for } q \neq k \text{ and } q = 1, \dots, n.$$

Moreover, since E commutes with $\int_C (z^n I - A^n)^{-1} dz$ and

$$\int_C (z^n I - A^n)^{-1} dz = \sum_{q=1}^n b_q A^{-(n-1)} \int_C (zI - e^{2\pi i q/n} A)^{-1} dz,$$

E also commutes with $A^{-(n-1)} \int_C (zI - e^{2\pi i k/n} A)^{-1} dz$. Thus the proof by induction is complete.

IV. Theorems.

THEOREM 3. Let A and E be elements of $\mathfrak{L}(\mathfrak{X})$ and let n be an integer greater than 1. If $\sigma(A) \cap \sigma(e^{2\pi i k/n} A) = \emptyset$ for $k=1, \dots, n-1$, and A^n commutes with E , then A commutes with E .

PROOF. Let $t_1, \dots, t_r$ be the elements of S which are interior to no other elements of S . Let C_j , $j=1, \dots, r$, be a proper curve for t_j . Let C be a simple closed rectifiable curve such that t_j is interior to C for $j=1, \dots, r$. By Theorem 2

$$(1) \quad \frac{1}{2\pi i} \int_C (zI - A)^{-1} dz = I$$

since $\sigma(A)$ lies entirely in the interior of C . Moreover, by Theorem 1 and Theorem 2

$$(2) \quad \frac{1}{2\pi i} \int_C (zI - A)^{-1} dz = \sum_{j=1}^r \frac{1}{2\pi i} \int_{C_j} (zI - A)^{-1} dz.$$

Combining (1) and (2), we see that

$$(3) \quad \sum_{j=1}^r \frac{A^{-(n-1)}}{2\pi i} \int_{C_j} (zI - A)^{-1} dz = A^{-(n-1)}.$$

Lemma 2 asserts that E commutes with each of the summands in the left hand side of equation (3). Therefore E commutes with $A^{-(n-1)}$ as well as A^n , which implies that E commutes with A .

One of the simplest situations in which the hypotheses of Theorem 3 are satisfied is when $\mathfrak{X}$ is a Hilbert space and either $\operatorname{Re} A \gg p > 0$ or $\operatorname{Im} A \gg p > 0$, for some real number p . Then A and A^2 commute with exactly the same operators, since $\sigma(A) \cap \sigma(-A) = \emptyset$. If in addition A^2 is normal, $AA^* = A^{*2}A$ since $A^2A^* = A^*A^2$. But this implies that $AA^* = A^*A$ since $\sigma(A^*) \cap (-A^*) = \emptyset$. More generally, we have by an analogous proof:

THEOREM 4. *Let $\mathfrak{X}$ be a Hilbert space. If A is an element of $\mathfrak{L}(\mathfrak{X})$ such that A^n is normal for some n greater than 1 and $\sigma(A) \cap \sigma(e^{2\pi i k/n} A) = \emptyset$ for $k=1, \dots, n-1$, then A is normal.*

REFERENCE

1. E. R. Lorch, *The spectrum of linear transformations*, Trans. Amer. Math. Soc. 52 (1942), 238-248.

UNIVERSITY OF NORTH CAROLINA AT CHARLOTTE