A LOWER BOUND FOR THE FIRST EIGENVALUE OF SECOND ORDER ELLIPTIC OPERATORS¹

H. OGAWA AND M. H. PROTTER

Let L be the second order operator

$$L = \sum_{i,j=1}^n a_{ij}(x) \frac{\partial^2}{\partial x_i \partial x_j} + \sum_{i=1}^n b_i(x) \frac{\partial}{\partial x_i} + c(x),$$

where the matrix (a_{ij}) is symmetric and positive definite in a bounded domain D . The coefficients are assumed to be real and so smooth that the adjoint operator L^* has continuous coefficients on $\bar{D}$. Let f and ϕ be positive functions on $\bar{D}$. By the methods of Hooker [1] and Protter [2], it can be shown that the lowest eigenvalue λ_1 of the problem

$$(1) \quad Lu + \lambda fu = 0 \quad \text{in } D,$$

$$(2) \quad u = 0 \quad \text{on } \partial D,$$

satisfies the inequality

$$(3) \quad \operatorname{Re} \lambda_1 \geq \inf_D (-(L\phi + L^*\phi)/2f\phi).$$

If L is selfadjoint, the inequality reduces to the extension of the inequality of Barta

$$\operatorname{Re} \lambda_1 \geq \inf_D (-L\phi/f\phi),$$

which Protter and Weinberger [3] recently established for nonselfadjoint second order elliptic operators.

The purpose of this paper is to show that the inequality (3) can be generalized to

$$(4) \quad \operatorname{Re} \lambda_1 \geq \inf_D \left[-\frac{1}{2f} \left(\frac{L\phi}{\phi} + \frac{L^*\psi}{\psi} \right) \right]$$

for arbitrary positive functions ϕ and ψ of class $C^2(\bar{D})$. Furthermore, we shall show that (4) also holds under more general boundary conditions, provided additional conditions are imposed on ϕ and ψ at the boundary. Such problems were studied by Hooker [1].

Let g and h be nonnegative functions on ∂D such that $g^2 + h^2 > 0$.

Received by the editors December 29, 1966.

¹ This work was supported by the National Science Foundation and the Air Force Office of Scientific Research.

We shall impose a boundary condition of the form

$$(5) \quad Bu \equiv gu + h \partial u / \partial \nu = 0 \quad \text{on } \partial D,$$

where $\partial u / \partial \nu$ denotes the conormal derivative associated with the operator L ; i.e.,

$$\frac{\partial}{\partial \nu} = \sum_{i,j} a_{ij} n_i \frac{\partial}{\partial x_j},$$

where n_i denotes the component of the outward unit normal to ∂D in the direction x_i . We shall also employ the adjoint boundary operator defined by

$$B^* = g + h(k + \partial / \partial \nu),$$

where

$$k = \sum_{i,j} \frac{\partial}{\partial x_j} (a_{ij}) n_i - \sum_i b_i n_i.$$

THEOREM. Let ϕ and ψ be positive functions of class $C^2(\overline{D})$ satisfying

$$(6) \quad (B\phi/\phi) + (B^*\psi/\psi) \geq 0 \quad \text{on } \partial D.$$

Then the lowest eigenvalue of the problem (1), (5) satisfies the inequality (4).

PROOF. Let v be a positive function of class $C^2(\overline{D})$ and set

$$I(v, \lambda) = -\frac{1}{2} \int_D v [\bar{u}(Lu + \lambda fu) + u(L\bar{u} + \bar{\lambda} f\bar{u})] dx.$$

Several applications of Green's Theorem yield

$$(7) \quad \begin{aligned} & I(v, \lambda) \\ &= \int_D \left\{ v \sum_{i,j} a_{ij} \frac{\partial u}{\partial x_i} \frac{\partial \bar{u}}{\partial x_j} - \left[\frac{1}{2} L^* v + \frac{1}{2} cv + (\operatorname{Re} \lambda) f v \right] |u|^2 \right\} dx \\ & \quad + \int_{\partial D} \left[\frac{1}{2} |u|^2 \left(\frac{\partial v}{\partial \nu} + kv \right) - \frac{1}{2} v \frac{\partial}{\partial \nu} |u|^2 \right] dS. \end{aligned}$$

If P_i , $i = 1, 2, \dots, n$, are functions of class $C^1(\overline{D})$, then

$$\int_D \sum_i \frac{\partial}{\partial x_i} (P_i |u|^2) dx - \int_{\partial D} |u|^2 \sum_i P_i n_i dS = 0.$$

Adding this to (7), we obtain

$$\begin{aligned}
 I(v, \lambda) = & \int_D \left\{ v \sum_{i,j} a_{ij} \frac{\partial u}{\partial x_i} \frac{\partial \bar{u}}{\partial x_j} + \sum_i P_i \left(u \frac{\partial \bar{u}}{\partial x_i} + \bar{u} \frac{\partial u}{\partial x_i} \right) \right. \\
 (8) \quad & \left. - \left[\frac{1}{2} L^* v + \frac{1}{2} c v + (\operatorname{Re} \lambda) f v - \sum_i \frac{\partial P_i}{\partial x_i} |u|^2 \right] \right\} dx \\
 & + \int_{\partial D} \left[\frac{1}{2} |u|^2 \left(\frac{\partial v}{\partial \nu} + k v \right) - \frac{1}{2} v \frac{\partial}{\partial \nu} |u|^2 - |u|^2 \sum_i P_i n_i \right] dS.
 \end{aligned}$$

For ϕ and ψ positive functions of class $C^2(\bar{D})$, an easy computation yields

$$\begin{aligned}
 & \frac{1}{2} L^* \left(\frac{\psi}{\phi} \right) + \frac{1}{2} c \frac{\psi}{\phi} \\
 & = \frac{\psi}{\phi} \left(\frac{1}{2} \frac{L\phi}{\phi} + \frac{1}{2} \frac{L^*\psi}{\psi} \right) - \sum_{i,j} \left[\frac{\partial}{\partial x_i} \left(\frac{a_{ij}\psi}{\phi^2} \frac{\partial \phi}{\partial x_j} \right) + \frac{a_{ij}\psi}{\phi^3} \frac{\partial \phi}{\partial x_i} \frac{\partial \phi}{\partial x_j} \right].
 \end{aligned}$$

Thus, if we set $v = \psi/\phi$ and

$$P_i = - \frac{\psi}{\phi^2} \sum_j a_{ij} \frac{\partial \phi}{\partial x_j}$$

and make use of the boundary condition (5), the equation (8) assumes the form

$$\begin{aligned}
 I\left(\frac{\psi}{\phi}, \lambda\right) = & \int_D \frac{\psi}{\phi} \left\{ \sum_{i,j} a_{ij} \left(\frac{\partial u}{\partial x_i} - \frac{u}{\phi} \frac{\partial \phi}{\partial x_i} \right) \left(\frac{\partial \bar{u}}{\partial x_j} - \frac{\bar{u}}{\phi} \frac{\partial \phi}{\partial x_j} \right) \right. \\
 & \left. - \left[\frac{1}{2} \frac{L\phi}{\phi} + \frac{1}{2} \frac{L^*\psi}{\psi} + (\operatorname{Re} \lambda) f \right] |u|^2 \right\} dx \\
 & + \int_{\Sigma} \frac{1}{2} \frac{\psi}{\phi h} |u|^2 \left(\frac{B\phi}{\phi} + \frac{B^*\psi}{\psi} \right) dS,
 \end{aligned}$$

where Σ is the subset of ∂D on which h is positive. Hence, because of the hypothesis (6), if λ is chosen so that

$$\frac{1}{2}(L\phi/\phi + L^*\psi/\psi) + (\operatorname{Re} \lambda)f < 0,$$

then $I(\psi/\phi, \lambda)$ is positive for all u not identically zero. Consequently, λ is not an eigenvalue and the theorem is proved.

For the case of Dirichlet boundary data, we have $h=0$, $g=1$, and hence condition (6) is satisfied for arbitrary positive functions ϕ and ψ .

Finally, we note that a generalization of (4) involving a finite num-

ber of pairs of positive functions, ϕ_k and ψ_k , can be found by considering the sum $\sum I(\psi_k/\phi_k, \lambda)$.

BIBLIOGRAPHY

1. W. W. Hooker, *Lower bounds for the first eigenvalue of elliptic equations of orders two and four*, Tech. Rep. 10. AF 49(638)–398, Univ. of California, Berkeley, Calif., 1960.
2. M. H. Protter, *Lower bounds for the first eigenvalue of elliptic equations*, Ann. of Math. **71** (1960), 423–444.
3. M. H. Protter and H. F. Weinberger, *On the spectrum of general second order operators*, Bull. Amer. Math. Soc. **72** (1966), 251–255.

UNIVERSITY OF CALIFORNIA, RIVERSIDE AND
UNIVERSITY OF CALIFORNIA, BERKELEY