ON STARLIKE FUNCTIONS¹

J. T. POOLE

I. Let S_k be the class of functions

$$f(z) = z + \sum_{n=k+1}^{\infty} a_n z^n,$$

regular and univalent in $|z| < 1$; let Σ_k be the class of functions

$$\tilde{f}(z) = z + \sum_{n=k-1}^{\infty} a_n z^{-n}, \quad f(z) \neq 0,$$

regular and univalent in $|z| > 1$, except for the simple pole at infinity. Finally, let S_k^* and Σ_k^* be the corresponding subclasses of starlike functions.

In a recent paper [6], T. H. MacGregor proved that if $f \in S_k^*$ then

$$|a_n| \leq \frac{mk}{n-1} \left| \binom{-2/k}{m} \right|$$

for $mk+1 \leq n \leq (m+1)k$, $m=1, 2, \dots$, thus generalizing Golusin's result [2] that if $f(z) = z + \sum_{m=1}^{\infty} a_{mk+1} z^{mk+1}$ belongs to S_k^* then

$$|a_{mk+1}| \leq \left| \binom{-2/k}{m} \right|.$$

It is the purpose of this paper to extend, in Theorem 1, MacGregor's result and then, in §II, to obtain some interesting corollaries to Theorem 1.

The proof of Theorem 1 follows MacGregor's method of proof [6].

THEOREM 1. Suppose $f \in S_k^*$ and for integral $t \geq 1$ let

$$f(z)^{-t} = z^{-t} + \sum_{\nu=-t+k}^{\infty} a_{\nu}^{(-t)} z^{\nu}.$$

Then

$$|a_{\nu}^{(-t)}| \leq \frac{mk}{\nu+t} \binom{2t/k}{m}$$

Received by the editors January 31, 1967.

¹ This research was supported in part by the National Science Foundation under grant NSF-GP-817.

for $-t + mk \leq \nu \leq -t + (m+1)k - 1$, m a positive integer, such that $t - (m-1)k \geq 0$.

PROOF. Let

$$(1) \quad g(z) = \frac{1}{-t} \frac{z[f(z)^{-t}]'}{f(z)^{-t}} = \frac{zf'(z)}{f(z)}$$

and define

$$(2) \quad w(z) = \frac{g(z) - 1}{g(z) + 1} = \sum_{\nu=k}^{\infty} \omega_{\nu} z^{\nu}.$$

From (1) and (2) we have

$$(3) \quad \{z[f(z)^{-t}]' - tf(z)^{-t}\}w(z) = z[f(z)^{-t}]' + tf(z)^{-t}.$$

Comparing coefficients in (3), we have

$$(4) \quad -2t\omega_{\nu} = \nu a_{-t+\nu}^{(-t)}, \quad \nu = k, \dots, 2k-1.$$

Since $\operatorname{Re}(g(z)) > 0$ in $|z| < 1$, $|w(z)| < 1$ in $|z| < 1$ and thus $\sum_{\nu=k}^{\infty} |\omega_{\nu}|^2 \leq 1$. Therefore, using (4), we have

$$(5) \quad \sum_{\nu=k}^{2k-1} \nu^2 |a_{-t+\nu}^{(-t)}|^2 = \sum_{\nu=k}^{2k-1} 4t^2 |\omega_{\nu}|^2 \leq 4t^2 \sum_{\nu=k}^{\infty} |\omega_{\nu}|^2 \leq 4t^2.$$

Writing (3) in the form

$$\begin{aligned} & \sum_{\nu=-t+k}^p (\nu+t) a_{\nu}^{(-t)} z^{\nu} + \sum_{\nu=p+1}^{\infty} c_{\nu} z^{\nu} \\ &= w(z) \left[-2tz^{-t} + \sum_{\nu=-t+k}^{p-k} (\nu-t) a_{\nu}^{(-t)} z^{\nu} \right], \end{aligned}$$

multiplying each side of this by its conjugate, integrating around $|z| = r < 1$ and letting $r \rightarrow 1$ yields the inequality

$$\sum_{\nu=-t+k}^p (\nu+t)^2 |a_{\nu}^{(-t)}|^2 \leq 4t^2 + \sum_{\nu=-t+k}^{p-k} (\nu-t)^2 |a_{\nu}^{(-t)}|^2,$$

from which we obtain

$$(6) \quad \sum_{\nu=p-k+1}^p (\nu+t)^2 |a_{\nu}^{(-t)}|^2 \leq 4t^2 + \sum_{\nu=-t+k}^{p-k} -4\nu t |a_{\nu}^{(-t)}|^2.$$

In (6) let $p = -t - 1 + (q+1)k$ and it then becomes

$$\begin{aligned}
 (7) \quad & \sum_{\nu=-t+qk}^{-t+(q+1)k-1} (\nu+t)^2 |a_\nu^{(-t)}|^2 \leq 4t^2 + \sum_{\nu=-t+k}^{-t+qk-1} -4\nu t |a_\nu^{(-t)}|^2 \\
 & = 4t \left[t + \sum_{\nu=-t+k}^{-t+qk-1} -\nu |a_\nu^{(-t)}|^2 \right] \\
 & = 4t \left[t + \sum_{m=1}^{q-1} \sum_{\nu=-t+mk}^{-t+(m+1)k-1} -\nu |a_\nu^{(-t)}|^2 \right].
 \end{aligned}$$

Now

$$\begin{aligned}
 \sum_{\nu=-t+k}^{-t+2k-1} -\nu |a_\nu^{(-t)}|^2 &= \frac{t-k}{k^2} \sum_{\nu=-t+k}^{-t+2k-1} \frac{k^2(-\nu)}{t-k} |a_\nu^{(-t)}|^2 \\
 &\leq \frac{t-k}{k^2} \sum_{\nu=-t+k}^{-t+2k-1} (\nu+t)^2 |a_\nu^{(-t)}|^2 \\
 &\leq (t-k) \frac{4t^2}{k^2} = (t-k) \binom{2t/k}{1},
 \end{aligned}$$

where the last inequality follows from (5). And we assume

$$\sum_{\nu=-t+mk}^{-t+(m+1)k-1} -\nu |a_\nu^{(-t)}|^2 \leq (t-mk) \binom{2t/k}{m}, \quad m = 2, \dots, q-1.$$

Then from (7)

$$\begin{aligned}
 \sum_{\nu=-t+qk}^{-t+(q+1)k-1} (\nu+t)^2 |a_\nu^{(-t)}|^2 &\leq 4t \left[t + \sum_{m=1}^{q-1} \sum_{\nu=-t+mk}^{-t+(m+1)k-1} -\nu |a_\nu^{(-t)}|^2 \right] \\
 &\leq 4t \left[t + \sum_{m=1}^{q-1} (t-mk) \binom{2t/k}{m} \right] \\
 &= (qk)^2 \binom{2t/k}{m}, \quad q \geq 2,
 \end{aligned}$$

so long as $t-(q-1)k \geq 0$. From this it follows that

$$\begin{aligned}
 \sum_{\nu=-t+qk}^{-t+(q+1)k-1} -\nu |a_\nu^{(-t)}|^2 &= \frac{t-qk}{(qk)^2} \sum_{\nu=-t+qk}^{-t+(q+1)k-1} \frac{(qk)^2(-\nu)}{t-qk} |a_\nu^{(-t)}|^2 \\
 &\leq \frac{t-qk}{(qk)^2} \sum_{\nu=-t+qk}^{-t+(q+1)k-1} (\nu+t)^2 |a_\nu^{(-t)}|^2 \\
 &\leq (t-qk) \binom{2t/k}{q}.
 \end{aligned}$$

Thus, by induction, we prove that

$$(8) \quad \sum_{\nu=-t+mk}^{-t+(m+1)k-1} (\nu+t)^2 \left| a_\nu^{(-t)} \right|^2 \leq (mk)^2 \binom{2t/k}{m}^2$$

and

$$(9) \quad \sum_{\nu=-t+mk}^{-t+(m+1)k-1} -\nu \left| a_\nu^{(-t)} \right|^2 \leq (t-mk) \binom{2t/k}{m}^2$$

for m such that $t-(m-1)k \geq 0$.

The theorem follows from (8).

If $\nu = -t + mk$ the inequality of Theorem 1 is sharp: consider $f(z) = z/(1+z^k)^{2/k}$. Also if $-t+k < \nu \leq -t+2k-1$ the inequality is sharp: consider $f(z) = z/(1+z^{\nu+t})^{2/(\nu+t)}$.

II. For $f \in S_k$ let $G = \{z \mid |z| < 1\}$. Suppose D is the image of G under the transformation $w = 1/\zeta$ and let E be the complement of D in the w -plane. E is a bounded continuum of capacity one with the coordinate system origin $0 \in E$. We define the moments $s_n = \int w^n d\mu$, $n = 1, 2, \dots$, where μ is the natural mass distribution on E . The coefficients of f can be expressed as polynomials in s_n [5]. Let $\tilde{f}(z) = 1/f(1/z)$ ($\tilde{f}$ maps $|z| > 1$ conformally onto D) and let $\phi(w) = w + \sum_{n=k+1}^{\infty} b_n w^n$ and $\tilde{\phi}(w) = w + \sum_{n=k-1}^{\infty} \tilde{b}_n w^n$ be the inverse functions to f and $\tilde{f}$ respectively.

It has been proved ([3], [4], [7]) that

1. $b_n = (1/n) a_{-1}^{(-n)}$, $n = 1, 2, \dots$.
2. $\tilde{b}_n = -(1/n) a_1^{(-n)}$, $n = 1, 2, \dots$.
3. $s_n = a_0^{(-n)}$, $n = 1, 2, \dots$.

4. If $s_n(\bar{0})$ are the moments obtained when the origin is situated at the center of gravity $\bar{0}$ ($=s_1$ in terms of 0) of E , i.e., the moments with respect to the center of gravity $\bar{0}$ of E , then

$$s_n = s_n(\bar{0}) + \binom{n}{1} s_{n-1}(\bar{0}) s_1 + \binom{n}{2} s_{n-2}(\bar{0}) s_1^2 + \dots + s_1^n.$$

In particular, if $s_1 = 0$ then $s_n(\bar{0}) = s_n$.

THEOREM 2. If $\phi(w) = w + \sum_{n=k+1}^{\infty} b_n w^n \in S_k^{*-1}$, then

$$|b_n| \leq \frac{mk}{n(n-1)} \binom{2n/k}{m}$$

for $mk+1 \leq n \leq (m+1)k$, $m = 1, 2, \dots$.

THEOREM 3. If $\tilde{\phi}(w) = w + \sum_{n=k-1}^{\infty} \tilde{b}_n w^n \in \Sigma_k^{*-1}$ then

$$|\bar{b}_n| \leq \frac{mk}{n(n+1)} \binom{2n/k}{m}$$

for $mk-1 \leq n \leq (m+1)k-2$, $m=1, 2, \dots$.

THEOREM 4. If $f \in S_k^*$ then

$$|s_n| \leq \frac{mk}{n} \binom{2n/k}{m}$$

for $mk \leq n \leq (m+1)k-1$, $m=1, 2, \dots$.

These theorems follow as corollaries to Theorem 1 and 1, 2 and 3 above, respectively.

If $f \in S_2^*$, i.e., f is starlike and $a_2=0$, then $s_1=0$. In this case $s_n(\bar{0}) = s_n$ and we have the following estimates for $s_n(\bar{0})$ as a corollary to Theorem 4.

COROLLARY. If $f \in S_2^*$ then

$$(10) \quad |s_{2m}(\bar{0})| \leq \binom{2m}{m}, \quad |s_{2m+1}(\bar{0})| \leq \frac{2m}{2m+1} \binom{2m+1}{m}$$

for $m=1, 2, \dots$.

The estimates (10) hold for a larger class of functions.

DEFINITION. Let $\bar{S}^*$ denote the class of univalent functions (functions in S_1) such that $f \in \bar{S}^*$ if and only if the corresponding continuum E contains its center of gravity $\bar{0}$ and is starshaped with respect to $\bar{0}$.

THEOREM 5. If $f \in \bar{S}^*$ then the estimates (10) hold.

PROOF. Suppose $f \in \bar{S}^*$ and has corresponding continuum E . Let $E_0 = \{w' | w' = w - \bar{0}, w \in E\}$ and note that E_0 has its center of gravity at the coordinate system origin and is starshaped with respect to the origin. There exists a function $f_0 \in S_1$ such that its corresponding continuum is E_0 . $f_0 \in S_2$ since the center of gravity of E_0 is $w=0$ which implies $a_2=0$. Since $f_0(z^{-1})^{-1}$ maps $|z| > 1$ onto the complement of E_0 and E_0 is starshaped with respect to $w=0$, $f_0(z^{-1})^{-1} \in \Sigma_2^*$. $f_0(z^{-1})^{-1} \in \Sigma_2^* \Rightarrow f_0 \in S_2^*$. Thus (10) holds for f_0 . But $s_n(\bar{0})$ for E is identical with $s_n(0) = s_n$ for E_0 , since E_0 is a translation of E . This completes the proof.

The estimates (10) are sharp for $s_{2m}(\bar{0})$, $m=1, 2, \dots$, and $s_3(\bar{0})$ (see the comments on the sharpness of Theorem 1).

Since it is known [3] that $|s_2(\bar{0})| \leq 2$ for all $f \in S_1$, it seems reasonable to conjecture that

$$|s_{2m}(\bar{0})| \leq \binom{2m}{m}, \quad m = 1, 2, \dots$$

for all $f \in S_1$. It is also known [1] that $|s_3(\bar{0})| = |-3a_4 + 6a_2a_3 - 3a_2^3| \leq 2$ for all $f \in S_1$.

REFERENCES

1. P. R. Garabedian and M. Schiffer, *A coefficient inequality for schlicht functions*, Ann. of Math. **61** (1955), 116–136.
2. G. Golusin, *On some estimates for functions which map the circle conformally and univalently*, Recueil Math. Moscow **36** (1929), 152–172. (Russian)
3. J. Gorski, *Some sharp estimations of coefficients of univalent functions*, J. Analyse Math. **14** (1965), 199–207.
4. J. Gorski and J. T. Poole, *Some sharp estimations of coefficients of univalent functions*, J. Math. Mech. **16** (1966), 577–582.
5. F. Leja, *Sur les coefficients des fonctions analytiques dans le cercle et les points extrémaux des ensembles*, Ann. Soc. Polon. Math. **23** (1950), 69–78.
6. T. H. MacGregor, *Coefficient estimates for starlike mappings*, Michigan Math. J. **10** (1963), 277–281.
7. J. T. Poole, *Coefficient extremal problems for schlicht functions*, Trans. Amer. Math. Soc. **121** (1966), 455–475.

FLORIDA STATE UNIVERSITY