COHOMOLOGY OF $\Omega(G/U)$

LARRY SMITH¹

Let G be a compact connected Lie group and $U \subset G$ a closed connected subgroup. Our objective is to determine the cohomology of the space of loops on the homogenous space G/U . Our main tool is a spectral sequence of Eilenberg and Moore ([3], [4], [7], [8]) from which we obtain the following:

[THEOREM P1. Let p be a prime and $T \subset G$ a maximal torus of G . Assume that $H^*(G, Z_p)$ is an exterior algebra on odd dimensional generators. Then there exists a filtration $\{F^{-n}H^*(\Omega(G/T), Z_p); n \geq 0\}$ such that

$$E^0H^*(\Omega(G/T), Z_p) = \text{Tor}_{H^*(G/T, Z_p)}(Z_p, Z_p)$$

as Hopf algebras.

THEOREM Q1. Let Q be the rational numbers and let G, T be as in Theorem P1, then there is an isomorphism of algebras

$$H^*(\Omega(G/T), Q) = \text{Tor}_{H^*(G/T, Q)}(Q, Q).$$

THEOREM P2. Let G be simply connected and U a closed connected subgroup and $i: U \subset G$ the inclusion map. Consider $H^*(U, Z_p)$ as an $H^*(G, Z_p)$ module via the map $i^*: H^*(G, Z_p) \rightarrow H^*(U, Z_p)$. Then if $H^*(G, Z_p)$ is an exterior algebra on odd dimensional generators there is a filtration $\{F^{-n}H^*(\Omega(G/U), Z_p); n \geq 0\}$ such that

$$E^0H^*(\Omega(G/U), Z_p) = \text{Tor}_{H^*(G, Z_p)}(Z_p, H^*(U, Z_p))$$

as Hopf algebras.

THEOREM Q2. Let Q be the rational numbers, G, U as in Theorem P2. Then there is an isomorphism of algebras

$$H^*(\Omega(G/U), Q) = \text{Tor}_{H^*(G, Q)}(Q, H^*(U, Q)).$$

1. Algebra. Throughout this section k will denote a fixed field. All modules will be graded k -modules of finite type, all algebras will be connected associative and commutative; $\otimes$ means $\otimes_k$. Our notation will be that of [5].

DEFINITION. If A is an algebra, $a_1, \dots, a_n \in A$, then we say that $a_1, \dots, a_n$ is an ESP-sequence if a_i is not a zero divisor in the algebra

Received by the editors November 4, 1966.

¹ Partially supported by NSF-GP-4037 and NSF-GP-3946.

$A/(a_1, \dots, a_{i-1})$ for $i=1, \dots, n-1$. (Here $(a_1, \dots, a_{i-1})$ denotes the ideal generated by $a_1, \dots, a_{i-1}$.)

DEFINITION. An algebra Λ is called a saturated algebra if it is the quotient of a polynomial algebra $P[x_1, \dots, x_n]$ by an ESP-sequence $r_1, \dots, r_n$.

It is easy to see that if Λ is a saturated algebra then it can be presented as a quotient of a polynomial algebra $P[y_1, \dots, y_m]$ by an ESP-sequence $s_1, \dots, s_m$ where no s_i is indecomposable. We will always assume that our presentations are minimal in this sense.

Our objective in this section is to compute $\text{Tor}_\Lambda(k, k)$ when Λ is a saturated algebra. To this end let

$$\Lambda = P[x_1, \dots, x_n]/(r_1, \dots, r_n)$$

be a minimal presentation of Λ . Let

$$\mathcal{E} = \Lambda \otimes E[u_1, \dots, u_n]$$

where $\deg u_i = (-1, \deg x_i)$. We give $\mathcal{E}$ the structure of a complex by defining a differential ∂ by the formulas

$$\partial(\lambda \otimes 1) = 0, \quad \partial(\lambda \otimes u_i) = \lambda x_i \otimes 1, \quad i = 1, \dots, n, \quad \lambda \in \Lambda,$$

and requiring ∂ to be a derivation of algebras. An easy calculation [10] shows

$$H^*(\mathcal{E}) = E[v_1, \dots, v_n], \quad \deg v_i = (-1, \deg r_i).$$

Now form the algebra

$$\mathcal{R} = \mathcal{E} \otimes \Gamma[z_1, \dots, z_n], \quad \deg z_i = (-2, \deg r_i)$$

and define a derivation d on $\mathcal{R}$ as follows: for each i choose a representative cocycle $\bar{v}_i \in \mathcal{E}$ representing $v_i \in H(\mathcal{E})$. Now define d by

$$d|_{\mathcal{E}} = \partial, \quad d(z_i) = \bar{v}_i, \quad i = 1, \dots, n,$$

and requiring that d be a derivation of algebras with divided powers. (Note that if the characteristic of k is zero then $\Gamma[z_1, \dots, z_n] = P[z_1, \dots, z_n]$.) As in [10] we can now show that $\mathcal{R} \rightarrow k$ is a Λ -free resolution of k and hence

PROPOSITION 1.1. $\text{Tor}_\Lambda(k, k) = E[u_1, \dots, u_n] \otimes \Gamma[z_1, \dots, z_n]$ as a Hopf algebra over k . $\square$

2. Torus subgroups. Let G be a compact connected Lie group and T a maximal torus of G . The inclusion $T \subset G$ induces a fibre bundle

$$\rho: G/T \rightarrow B_T \xrightarrow{\rho} B_G$$

where B_T and B_G are classifying spaces for T and G respectively. Let $k = Z_p$ or Q and assume that $H^*(G, k)$ is an exterior algebra on odd dimensional generators (this is no restriction if $k = Q$). We then have the fundamental result of Borel [2]:

THEOREM ([2], [1]). $H^*(B_G, k) = P[y_1, \dots, y_n]$ $n = \text{rank } G$, $\rho^*(y_1), \dots, \rho^*(y_n)$ is an ESP-sequence in $H^*(B_T, k)$ and $H^*(G/T, k) = H^*(B_T, k) // \rho^*$. Since $H^*(B_T, k) = P[x_1, \dots, x_n]$ $\deg x_i = 2$ it follows that $H^*(G/T, k)$ is a saturated algebra.

PROOF OF THEOREM P1. Let $\{E_r, d_r\}$ denote the Eilenberg-Moore spectral sequence [3], [4], [8] of the pathspace fibration

$$\Omega(G/T) \rightarrow P(G/T) \rightarrow G/T$$

we then have

$$E_r \Rightarrow H^*(\Omega(G/T), Z_p), \quad E_2 = \text{Tor}_{H^*(G/T, Z_p)}(Z_p, Z_p).$$

We will show that $E_2 = E_\infty$ which will complete the proof. From §1 we see that

$$E_2 = E[u_1, \dots, u_m] \otimes \Gamma[z_1, \dots, z_m]$$

where $m \leq n$, $\deg u_i = (-1, 2)$, $\deg z_i = (-2, 2t_i)$, $t_i \geq 2$. Now $\{E_r, d_r\}$ is a spectral sequence of Hopf algebras. Let s be the smallest integer ≥ 2 such that $d_s \neq 0$. Then $E_2 = E_s$. Let $a \in E_2$ of minimal degree such that $d_s(a) \neq 0$. Then we may assume that a is indecomposable, and therefore $\deg a = (-2p^q, 2tp^q)$, $t > 1$. A simple computation shows that $d_s(a)$ must be primitive. Since the total degree of a is even it follows that $d_s(a)$ is a primitive of odd total degree and hence $\deg d_s(a) = (-1, 2)$. Since $\deg d_s = (s, 1-s)$ we see that $s = 2p^q - 1$ and hence we have $\deg d_s(a) = (-1, 2 + 2(t-1)p^q)$. But $t > 1$ and hence $2 + 2(t-1)p^q > 2$ which contradicts our previous calculation of $\deg d_s(a)$. Therefore our assumption that $d_s(a) \neq 0$ must be false and hence $E_2 = E_\infty$. $\square$

PROOF OF THEOREM Q1. We recall [2] that $H^*(G, Q)$ is an exterior algebra on odd dimensional generators. By the argument above we see the Eilenberg-Moore spectral sequence of the fibration

$$\Omega G/T \xrightarrow{j} P(G/T) \xrightarrow{i} G/T$$

with Q coefficients has $E_2 = E_\infty$.

But $E_2 = E[u_1, \dots, u_m] \otimes P[z_1, \dots, z_m]$ is a free commutative algebra and hence the extension as algebras is trivial. $\square$

3. A fibration. Let U be a closed connected subgroup of G and $i: U \subset G$ the inclusion map. By a standard construction [9, p. 99] we can replace i by an equivalent fibre mapping whose fibre is $\Omega(G/U)$. This gives a multiplicative fibre map

$$\Omega(G/U) \rightarrow U \rightarrow G.$$

If G is simply connected then this fibring has for each field k an Eilenberg-Moore spectral sequence $\{\tilde{E}_r, \tilde{d}_r\}$ with

$$\tilde{E}_r \Rightarrow H^*(\Omega(G/U), k), \quad \tilde{E}_2 = \text{Tor}_{H^*(G,k)}(k, H^*(U, k)).$$

Henceforth we shall assume that $k = \mathbb{Z}_p$ or $\mathbb{Q}$ and that $H^*(G, k)$ is an exterior algebra on odd dimensional generators. From [2] it follows that $H^*(G, k)$ is primitively generated and hence cocommutative. Let $H^*(G, k) \backslash i^*$ denote the kernel in the category of bicommutative Hopf algebras of the map $i^*: H^*(G, k) \rightarrow \text{im } i^*$ (see [6], [7], [8]). Then as in [7], [8] we can show

PROPOSITION.

$$\text{Tor}_{H^*(G,k)}(k, H^*(U, k)) = H^*(U, k) \backslash i^* \otimes \text{Tor}_{H^*(G,k)} \backslash i^* (k, k)$$

as Hopf algebras. $\square$

4. The simply connected case. Before proving Theorem P2 we state a preliminary technical result whose proof we postpone until later.

PROPOSITION.

$$\ker\{j^*: H^*(U, \mathbb{Z}_p) \rightarrow H^*(\Omega(G/U), \mathbb{Z}_p)\} = \overline{\text{im } i^*} \cdot H^*(U, \mathbb{Z}_p).$$

PROOF OF THEOREM P2. By the Proposition of §3 we have

$$E_2 = H^*(U, \mathbb{Z}_p) \backslash i^* \otimes \text{Tor}_{H^*(G, \mathbb{Z}_p)} \backslash i^* (\mathbb{Z}_p, \mathbb{Z}_p).$$

From Borel's structure theorem for Hopf algebras over $\mathbb{Z}_p$ [6, 7.11] we deduce that $H^*(G, \mathbb{Z}_p) \backslash i^* = E[u_1, \dots, u_m]$, degree u_i odd, and hence we deduce that as Hopf algebras

$$E_2 = H^*(U, \mathbb{Z}_p) \backslash i^* \otimes \Gamma[y_1, \dots, y_m]$$

where $\deg y_i = (-1, \deg u_i)$. Suppose that $E_2 \neq E_\infty$. A repetition of the argument in Theorem P1 shows that there is an $x \in E_2$ such that $d_r(x) \neq 0 \in E_2^{0,*}$. The map j^* is given by the composition

$$H^*(U, \mathbb{Z}_p) \rightarrow H^*(U, \mathbb{Z}_p) \backslash i^* = E_2^{0,*} \xrightarrow{\epsilon} E_\infty^{0,*} \subset H^*(\Omega(G/U), \mathbb{Z}_p)$$

and hence from the Proposition above it follows that the edge map ϵ is a monomorphism. But this contradicts the fact that $d_r(x) \neq 0$. Therefore it follows that $E_2 = E_\infty$. $\square$

PROOF OF PROPOSITION. Consider the fibration

$$\Omega(G) \rightarrow \Omega(G/U) \xrightarrow{j} U$$

obtained by replacing j by a fibre map. Let $\{\hat{E}^r, \hat{d}^r\}$ denote the homology Serre spectral sequence of this fibration. Consider the diagram

$$\begin{array}{ccc} \Omega(G) & \rightarrow & \Omega(G) \\ \downarrow & & \downarrow \\ \Omega(G/U) & \rightarrow & P(G) \\ \downarrow & & \downarrow \\ U & \rightarrow & G \end{array}$$

and recall that we have by [6, 4.4]

- (1) $H_*(G, Z_p) = E[x_1, \dots, x_n]$
- (2) $H_*(U, Z_p) = E[y_1, \dots, y_t] \otimes M$ where

$$\begin{aligned} i_*(y_i) &= x_i, & 1 \leq i \leq t, \\ i_*/M &= 0, \end{aligned}$$

and

(3) $H_*(\Omega(G), Z_p) = P[z_1, \dots, z_n]$ with $\tau x_i = z_i$ where τ is the transgression in the path space fibring over G . A standard spectral sequence argument now shows that $\hat{E}^\infty = M \otimes P[z_{t+1}, \dots, z_n]$ and since the map j_* is the composition $H_*(U, Z_p) \rightarrow \hat{E}_{0,*}^2 \rightarrow \hat{E}_{0,*}^\infty \subset H_*(\Omega(G/U), Z_p)$ the result follows by duality. $\square$

The proof of Theorem Q2 is an easy exercise and is left to the reader.

In closing we note that if G is simply connected and T is any torus in G then the inclusion map $i: T \subset G$ is null homotopic and hence standard homotopy theory shows that $\Omega(G/T)$ has the homotopy type of $T \times \Omega(G)$, and hence the study of torus subgroups in the simply connected case is trivial.

REFERENCES

1. P. F. Baum, *Cohomology of homogenous spaces*, Topology **7** (1968), 78–106.
2. A. Borel, *Sur la cohomologie des espaces fibre princepeaux*, Ann. of Math. **57** (1953), 115–207.
3. S. Eilenberg and J. C. Moore, *Homology and fibrations*. I, Comment. Math. Helv. **40** (1966), 199–236.

4. ———, *Homology and fibrations*. II, III, . . . , Comment. Math. Helv. (to appear).
5. S. MacLane, *Homology*, Academic Press, New York, 1963.
6. J. Milnor and J. C. Moore, *The structure of Hopf algebras*, Ann. of Math. **81** (1965), 211–264.
7. L. Smith, *Multiplicative fibre maps*, Bull. Amer. Math. Soc. **73** (1967), 334–338.
8. ———, *Cohomology of stable two stage Postnikov systems*, Illinois J. Math. **11** (1967), 310–329.
9. E. H. Spanier, *Algebraic topology*, McGraw-Hill, New York, 1966.
10. J. Tate, *Homology of Noetherian rings and of local rings*, Illinois J. Math. **1** (1957).

YALE UNIVERSITY AND
PRINCETON UNIVERSITY