ON A THEOREM IN COMPLETE $\mathfrak{A}$ -ADIC RINGS

BERNARD KOLMAN

Introduction. In this paper we characterize a local ring R with maximal ideal $\mathfrak{M}$ and ideal $\mathfrak{A}$ such that R is equicharacteristic, $R/\mathfrak{A}$ is regular, and R is Hausdorff and complete with respect to the $\mathfrak{A}$ -adic topology. The key result is an extension of a theorem due to Harrison [5], which is analogous to the first part of Cohen's Theorem 9 [1]. Methods similar to those used in Geddes [3], [4] and in Curtiss [2] are used.

DEFINITION. A local ring R with maximal ideal $\mathfrak{M}$ and an ideal $\mathfrak{A}$ is called a special local ring if it satisfies the following properties:

- (1) R is equicharacteristic.
- (2) $R/\mathfrak{A}$ is regular.
- (3) R is a complete space with respect to the $\mathfrak{A}$ -adic topology.

Harrison [5] has proved the following

THEOREM A. *Let R be a special local ring. If $\mathfrak{A}^2 = 0$, then there exists a subring B of R such that $B + \mathfrak{A} = R$ and $B \cap \mathfrak{A} = (0)$.*

We wish to extend this theorem by showing that the conclusion holds when R is Hausdorff and $\mathfrak{A}^2 \neq 0$. We first prove the following:

LEMMA 1. *Let R be a Noetherian ring, containing an identity 1, $\mathfrak{A}$ an ideal of R such that $R/\mathfrak{A}$ is local, and $\mathfrak{A}^2 = 0$. Then R is local.*

PROOF. Let $\mathfrak{M}'$ be the unique maximal ideal of $R/\mathfrak{A}$, and $\sigma: R \rightarrow R/\mathfrak{A}$ the natural homomorphism. Let $\mathfrak{M} = \sigma^{-1}(\mathfrak{M}')$. If $x \notin \mathfrak{M}$, then $\sigma(x)$ is a unit in $R/\mathfrak{A}$. Thus, there exists $y \in R$ such that $\sigma(x)\sigma(y) = \sigma(1)$, which implies that $xy - 1 \in \mathfrak{A}$. Let $xy - 1 = \alpha$, $\alpha \in \mathfrak{A}$. Then $x(y - \alpha y) = xy - \alpha xy = 1 + \alpha - \alpha(1 + \alpha) = 1 + \alpha - \alpha - \alpha^2 = 1$. Hence, x is a unit in R , and it then follows that R is local.

THEOREM 1. *Let R be a special local ring which is Hausdorff in the $\mathfrak{A}$ -adic topology. If $\mathfrak{A}^2 \neq (0)$, then there exists a subring B of R such that $B + \mathfrak{A} = R$, $B \cap \mathfrak{A} = (0)$.*

PROOF. Let $R' = R/\mathfrak{A}^2$. Then R' is a local ring with maximal ideal $\mathfrak{M}' = \mathfrak{M}/\mathfrak{A}^2$. Moreover, R' is a special local ring with respect to the ideal $\mathfrak{A}' = \mathfrak{A}/\mathfrak{A}^2$. To verify this, we note that (2) in the definition of special local ring follows from $R'/\mathfrak{A}' = (R/\mathfrak{A}^2)/(\mathfrak{A}/\mathfrak{A}^2) \simeq R/\mathfrak{A}$, which is regular. (3) follows from the corresponding property for R . (1) follows from the fact that $R'/\mathfrak{M}' = (R/\mathfrak{A}^2)/(\mathfrak{M}/\mathfrak{A}^2) \simeq R/\mathfrak{M}$.

Received by the editors February 23, 1967.

Applying Theorem A to R' , we have a subring R'_1 of R' such that $R'_1 + \mathfrak{A}' = R'$ and $R'_1 \cap \mathfrak{A}' = \mathfrak{A}'^2$. Letting R_1 be the inverse image of R'_1 under the natural homomorphism $R \rightarrow R' = R/\mathfrak{A}^2$, we have $R_1 + \mathfrak{A} = R$, and $R_1 \cap \mathfrak{A} = \mathfrak{A}^2$. We now assert that $1 \in R_1$. For we have $1 = r_1 + a$, where $r_1 \in R_1$, $a \in \mathfrak{A}$. Then $r_1^2 - r_1 = ar_1 \in R_1 \cap \mathfrak{A} = \mathfrak{A}^2$. Since $a^2 \in R_1$, we have $1 - r_1^2 \in R_1$, and thus $1 \in R_1$.

Now $(R_1/\mathfrak{A}^4)/(\mathfrak{A}^2/\mathfrak{A}^4) \simeq R_1/\mathfrak{A}^2 \simeq R_1/(R_1 \cap \mathfrak{A}) \simeq (R_1 + \mathfrak{A})/\mathfrak{A} \simeq R/\mathfrak{A}$, and hence $(R_1/\mathfrak{A}^4)/(\mathfrak{A}^2/\mathfrak{A}^4)$ is a local ring and $(\mathfrak{A}^2/\mathfrak{A}^4)^2 = \bar{0}$. Also, $1 \in R_1/\mathfrak{A}^4$. By Lemma 1, we conclude that $R_1/\mathfrak{A}^4$ is a local ring. Let $R'_1 = R_1/(\mathfrak{A}^2)^2$, $\mathfrak{A}'' = \mathfrak{A}^2/(\mathfrak{A}^2)^2$ an ideal in R'_1 , $\mathfrak{M}''$ the maximal ideal of R'_1 . We now verify that R'_1 is a special local ring. The residue field of R'_1 is $R'_1/\mathfrak{M}''$, and it then follows that $\text{char } R'_1 = \text{char } R'_1/\mathfrak{M}''$. Moreover, $R'_1/\mathfrak{A}'' = (R_1/\mathfrak{A}^4)/(\mathfrak{A}^2/\mathfrak{A}^4) \simeq R_1/\mathfrak{A}^2 = R_1/(R_1 \cap \mathfrak{A}) \simeq (R_1 + \mathfrak{A})/\mathfrak{A} = R/\mathfrak{A}$, which implies that $R'_1/\mathfrak{A}''$ is regular. Also, if R is complete in its $\mathfrak{A}$ -topology then it is complete in its $\mathfrak{A}^2$ -topology. For if $\{a\}$ is a regular sequence in R with respect to the $\mathfrak{A}^2$ -topology, then given any integer $s \geq 0$ there exists an integer N_1 such that if $n, m \geq N_1$ then $a_n - a_m \in \mathfrak{A}^{2s}$. Moreover, $\{a_j\}$ is regular with respect to the $\mathfrak{A}$ -topology, and thus $\{a_j\}$ has a limit a . Thus, given any $r \geq 0$, there exists N_2 such that if $m > N_2$ then $a_m - a \in \mathfrak{A}^r$. Then if $n > N = \max(N_1, N_2)$ we have $a_n - a \in \mathfrak{A}^{2s} + \mathfrak{A}^r$. Since $\mathfrak{A}^{2s}$ is open in the $\mathfrak{A}$ -topology, it is closed. Thus, $\mathfrak{A}^{2s} = \bigcap_{r=1}^{\infty} (\mathfrak{A}^{2s} + \mathfrak{A}^r)$, which implies that $a_n - a \in \mathfrak{A}^{2s}$ for $n > N$, and thus a is the limit of $\{a_j\}$ in the $\mathfrak{A}^2$ -topology. Thus, R is complete in its $\mathfrak{A}^2$ topology. Since $R_1 \cap \mathfrak{A}^s = \mathfrak{A}^s$, $s \geq 2$, then R_1 is complete with respect to the $\mathfrak{A}^2$ -adic topology. Then $R_1/\mathfrak{A}^4$ is complete in the $\mathfrak{A}^2/\mathfrak{A}^4$ topology. Then by Theorem A, and going back in the natural homomorphism $R_1 \rightarrow R_1/\mathfrak{A}^4$, there exists a subring R_2 of R_1 such that $R_2 + \mathfrak{A}^2 = R_1$ and $R_2 \cap \mathfrak{A}^2 = \mathfrak{A}^4$. Thus, by induction, there exists a sequence of subrings $\{R_n\}$, $R_{n+1} \subset R_n$ such that $R_{n+1} + \mathfrak{A}^{2^n} = R_n$ and $R_{n+1} \cap \mathfrak{A}^{2^n} = \mathfrak{A}^{2^{n+1}}$. Let $B = \bigcap_{i=1}^{\infty} R_i$. We then have $R_i \cap \mathfrak{A} = \mathfrak{A}^{2^i}$, for $\mathfrak{A}^{2^i} = R_i \cap \mathfrak{A}^{2^{i-1}} = R_i \cap R_{i-1} \cap \mathfrak{A}^{2^{i-2}} = R_i \cap R_{i-1} \cap R_{i-2} \cap \cdots \cap R_1 \cap \mathfrak{A} = R_i \cap \mathfrak{A}$. Then

$$B \cap \mathfrak{A} = \left(\bigcap_{i=1}^{\infty} R_i \right) \cap \mathfrak{A} = \bigcap_{i=1}^{\infty} (R_i \cap \mathfrak{A}) = \bigcap_{i=1}^{\infty} \mathfrak{A}^{2^i} = (0).$$

Next, we show that $B + \mathfrak{A} = R$. We first have that $R = R_1 + \mathfrak{A}$, and by induction get $R = R_n + \mathfrak{A}$, for any n . Thus, if $a \in R$, then for any n there exists $b_n \in R_n$ such that $a \equiv b_n (\mathfrak{A})$. Then $b_n - b_{n+1} \in R_n \cap \mathfrak{A}$. Since $R_n \cap \mathfrak{A} = \mathfrak{A}^{2^n}$, $\{b_n\}$ is a regular sequence in R . Since R is complete, $\{b_n\}$ has a limit b in R . Also, $R_n \supset \mathfrak{A}^{2^n}$, and hence R_n is closed in the $\mathfrak{A}$ -adic topology. Thus, $b \in R_n$ for all n , which implies that

$b \in B$. Moreover, since $a \equiv b_n(\mathfrak{A})$, $b_n \rightarrow b$ and we have $a \equiv b(\mathfrak{A})$. Thus, $B + \mathfrak{A} = R$. This completes the proof of Theorem 1.

LEMMA 2. *Let R be a Noetherian ring which is complete with respect to the $\mathfrak{A}$ -adic topology, where $\mathfrak{A} = (u_1, u_2, \dots, u_n)$. Let $S = R\{x_1, x_2, \dots, x_m\}$ be the ring of formal power series in m indeterminates over R . Then S is a Noetherian ring which is complete with respect to the $\mathfrak{A}^*$ -adic topology, where $\mathfrak{A}^* = (u_1, u_2, \dots, u_n, x_1, x_2, \dots, x_m)$. Moreover, if the above basis for $\mathfrak{A}$ is minimal then so is the indicated basis for $\mathfrak{A}^*$.*

PROOF. To show that if R is Noetherian then S is Noetherian we proceed as in the Hilbert basis theorem. We now show that S is complete in the $\mathfrak{A}^*$ -adic topology. Let $\{f^i\}$ be a regular sequence in S . Then $f^i = \sum_{k=0}^{\infty} f_k^i$, where f_k^i is a form in $x_1, x_2, \dots, x_m$ of degree k . Since $\{f^i\}$ is regular in S , given any $s > 0$ then $f^i - f^{i+1} \in \mathfrak{A}^{*s}$ for i sufficiently large. Now

$$\mathfrak{A}^{*s} = \left\{ \sum_{k=0}^{\infty} t_k \mid t_k \in \mathfrak{A}^{s-k} \cdot R[x_1, x_2, \dots, x_n] \ \forall k < s \right\}.$$

Fixing k , and for $s > k$, we then have that the coefficients of $f_k^i - f_k^{i+1} \in \mathfrak{A}^{s-k}$. Thus, the coefficients of each monomial in f_k^i form a regular sequence in R , and since R is complete they have a limit in R . Thus there exists a form f_k of degree k such that $\lim_{i \rightarrow \infty} f_k^i = f_k$. Let $f = \sum_k f_k$. Then $f^i - f = \sum_k (f_k^i - f_k)$, and for any $s \geq 0$, $f_k^i - f_k \in \mathfrak{A}^{s-k} \cdot R[x_1, x_2, \dots, x_n]$ for all $k < s$ and i sufficiently large. Hence, $f^i - f \in \mathfrak{A}^{*s}$ for i sufficiently large, which implies that S is complete.

Now suppose that the above basis of $\mathfrak{A}$ is minimal and that the basis of $\mathfrak{A}^*$ is not minimal. Thus, suppose $x_1 = \sum_{i=1}^n f_i u_i + \sum_{j=2}^m g_j x_j$, $f_i, g_j \in S$. We expand the right-hand side into a power series in $x_1, x_2, \dots, x_m$ and observe that $\sum_{j=2}^m g_j x_j$ contributes no term in x_1 alone. Thus, if $\sum_{i=1}^n f_i u_i$ contributes no such term then $1 = 0$. If $\sum_{i=1}^n f_i u_i$ does contribute such a term then its coefficient $\in \mathfrak{A}$, which implies that $1 \in \mathfrak{A}$ and thus $\mathfrak{A} = R$, a contradiction. If we assume that $u_1 = \sum_{i=2}^n f'_i u_i + \sum_{j=1}^m g_j x_j$, $f'_i, g_j \in S$, then $u_1 = \sum_{i=2}^n f''_i u_i$, where f'_i is the constant term of f_i , which contradicts the minimality of the basis $(u_1, u_2, \dots, u_n)$. This completes the proof.

The following theorem characterizes a special local ring which is Hausdorff with respect to the $\mathfrak{A}$ -adic topology, and is analogous to the second part of Cohen's Theorem 9 [1]. The proof follows along the lines of Cohen's proof.

THEOREM 2. *Let R be a special local ring with maximal ideal $\mathfrak{M}$, Hausdorff in the $\mathfrak{A}$ -adic topology, where $\mathfrak{A}$ is an ideal having a maximal basis of n elements. Then R is a homomorphic image of a formal power series ring $(R/\mathfrak{A})\{x_1, x_2, \dots, x_n\}$.*

PROOF. By Theorem 1, there exists a subring B of R such that $B + \mathfrak{A} = R$, $B \cap \mathfrak{A} = (0)$. Let $\mathfrak{A} = (u_1, u_2, \dots, u_n)$. We first show that R is the closure of $B[u_1, u_2, \dots, u_n]$. Let $c \in R$. Then we find a sequence $\{c_k\}$, $c_k \in B[u_1, u_2, \dots, u_n]$, such that $\lim c_k = c$, i.e., we show that $c - c_k \in \mathfrak{A}^{k+1}$ by induction.

Since $c \in R$, there exists $b \in B$ such that $c \equiv b(\mathfrak{A})$. Let $c_0 = b$. Assume that c_k has been defined. Then $c - c_k \in \mathfrak{A}^{k+1}$, which implies that $c - c^k = \sum_{i=1}^n r_i v_i$, where $r_i \in R$, and v_i is a power product of $u_1, u_2, \dots, u_n$ of degree $k+1$. Since $r_i \in R$, there exists $b_i \in R$ such that $r_i \equiv b_i(\mathfrak{A})$. Then $c - c_k \equiv \sum_{i=1}^n b_i v_i(\mathfrak{A}^{k+2})$. Letting $c_{k+1} = c_k + \sum_{i=1}^n b_i v_i$, we have constructed the desired sequence $\{c_k\}$. Now in the natural mapping of R onto $R/\mathfrak{A}$, B is mapped onto $R/\mathfrak{A}$. There exists a unique homomorphism T of $(R/\mathfrak{A})[x_1, x_2, \dots, x_n]$ onto $B[u_1, u_2, \dots, u_n]$ such that $x_i T = u_i$ and T restricted to $R/\mathfrak{A}$ is the inverse of the natural mapping of B onto $R/\mathfrak{A}$. T is continuous, where the topology of $(R/\mathfrak{A})[x_1, x_2, \dots, x_n]$ is the relative topology determined by $(R/\mathfrak{A})\{x_1, x_2, \dots, x_n\}$. Since R is complete and is the closure of $B[u_1, u_2, \dots, u_n]$ and $(R/\mathfrak{A})\{x_1, x_2, \dots, x_n\}$ is the closure of $(R/\mathfrak{A})[x_1, x_2, \dots, x_n]$, then T can be uniquely extended to a homomorphism of $(R/\mathfrak{A})\{x_1, x_2, \dots, x_n\}$ onto R . This completes the proof of Theorem 2.

BIBLIOGRAPHY

1. I. S. Cohen, *On the structure and ideal theory of complete local rings*, Trans. Amer. Math. Soc. **59** (1946), 54–106.
2. C. W. Curtiss, *The structure of non semi-simple algebras*, Duke Math. J. **21** (1954), 79–85.
3. A. Geddes, *A short proof of the existence of coefficient fields for complete equi-characteristic local rings*, J. London Math. Soc. **29** (1954), 334–341.
4. ———, *On the embedding theorems for complete local rings*, Proc. London Math. Soc. (3) **6** (1956), 343–354.
5. D. K. Harrison, *Commutative algebras and cohomology*, Trans. Amer. Math. Soc. **104** (1962), 191–204.

DREXEL INSTITUTE OF TECHNOLOGY