ON THE HYPERFINITE II_1 -FACTOR

SHÔICHIRO SAKAI¹

1. Introduction. In this paper, we shall consider the following problem which has been asked by several mathematicians: Can we embed any arbitrary finite factor on a separable Hilbert space into the hyperfinite II_1 -factor? The answer is negative.

2. Theorem. First of all, we shall show,

LEMMA 1. *Let M be a finite W^* -algebra on a Hilbert space $\mathfrak{H}$ such that M' has the property P in the sense of Schwartz [2], $B(\mathfrak{H})$ the W^* -algebra of all bounded operators on $\mathfrak{H}$, and let N be a W^* -subalgebra of M , then there exists a linear mapping P on $B(\mathfrak{H})$ to N satisfying the following conditions:*

- (1) $P(x^*) = P(x)^*$ for $x \in B(\mathfrak{H})$,
- (2) $P(h) \geq 0$ for $h(\geq 0) \in B(\mathfrak{H})$, and $P(I) = I$,
- (3) $P(axb) = aP(x)b$ for $a, b \in N$ and $x \in B(\mathfrak{H})$.

PROOF. By the result of Schwartz [2], there is a linear mapping P_1 on $B(\mathfrak{H})$ to M satisfying the same properties. On the other hand, by the result of Umegaki [3], there is a linear mapping P_2 on M to N satisfying the same properties. Now put $P(x) = P_2(P_1(x))$ for $x \in B(\mathfrak{H})$, then clearly P satisfies the required properties. This completes the proof.

Now let M be the hyperfinite II_1 -factor, and let N be the II_1 -factor generated by the left regular representation of a countable discrete free group G with two generators.

Then we shall show the following theorem.

THEOREM. *There is no subfactor in M which is $*$ -isomorphic to N .*

To prove the theorem, we shall proceed as follows.

Now suppose that there is a subfactor N_1 of M which is $*$ -isomorphic to N .

Let $\{\pi, \mathfrak{H}\}$ be the standard $*$ -representation of M on a Hilbert space $\mathfrak{H}$.

Let K be an $\aleph_0$ -dimensional Hilbert space, and let $\mathfrak{H} \otimes K$ be the tensor product of $\mathfrak{H}$ and K . We shall consider the W^* -representation $\{\pi \otimes I_K, \mathfrak{H} \otimes K\}$ of M on $\mathfrak{H} \otimes K$, where $\pi \otimes I_K$ is the amplification of π (cf. [1]) and I_K is the identity operator on K .

Received by the editors February 15, 1967.

¹ This paper was written with partial support from ONR Contract NR-551(57).

LEMMA 2. *The commutant $\{\pi(M) \otimes I_K\}' = \pi(M)' \otimes B(K)$ in $\mathfrak{H} \otimes K$ has the property P, where $B(K)$ is the W^* -algebra of all bounded operators on K .*

PROOF. $\pi(M)'$ has the property P (cf. [2]). Let a be an arbitrary element of the W^* -algebra $B(\mathfrak{H} \otimes K)$ of all bounded operators on the Hilbert space $\mathfrak{H} \otimes K$, and let $C_1(a)$ be the σ -weakly closed convex envelope of $\{u^*au \mid u \text{ (unitary)} \in I_{\mathfrak{H}} \otimes B(K)\}$, where $I_{\mathfrak{H}}$ is the identity operator on $\mathfrak{H}$. Then $C_1(a) \cap (I_{\mathfrak{H}} \otimes B(K))' = C_1(a) \cap (B(\mathfrak{H}) \otimes I_K) \neq (\emptyset)$, because $I_{\mathfrak{H}} \otimes B(K)$ is a type I factor so that there exists an increasing sequence of $2^n \times 2^n$ matrix algebras $\{M_n \mid n = 1, 2, 3, \dots\}$ such that the σ -weak closure of $\bigcup_{n=1}^{\infty} M_n = I_{\mathfrak{H}} \otimes B(K)$.

Next, take $a_0 \in C_1(a) \cap (B(\mathfrak{H}) \otimes I_K)$, and let $C_2(a_0)$ be the σ -weakly closed convex envelope of $\{u^*a_0u \mid u \text{ (unitary)} \in \pi(M)' \otimes I_K\}$. Then clearly $C_2(a_0) \cap \{\pi(M)' \otimes I_K\}' \cap B(\mathfrak{H}) \otimes I_K \neq (\emptyset)$, because $\pi(M)'$ has the property P. On the other hand, $C_2(a_0)$ is contained in the σ -weakly closed convex envelope of $\{u^*au \mid u \text{ (unitary)} \in \pi(M)' \otimes B(K)\}$. Hence $\pi(M)' \otimes B(K)$ has the property. This completes the proof.

The commutant $\{\pi(N_1) \otimes I_K\}'$ of $\pi(N_1) \otimes I_K = \pi(N_1)' \otimes B(K)$; hence it is a II_{∞} -factor.

Let ξ be a trace vector for M in $\mathfrak{H}$ and η be a nonzero vector in K .

Let $[\pi(N_1) \otimes I_K \xi \otimes \eta]$ be the closed subspace of $\mathfrak{H} \otimes K$ generated by $\pi(N_1) \otimes I_K \xi \otimes \eta$ and E' be the orthogonal projection of $\mathfrak{H} \otimes K$ onto $[\pi(N_1) \otimes I_K \xi \otimes \eta]$. Then the representation $x \rightarrow (\pi(x) \otimes I_K)E'(x \in N_1)$ of N_1 is standard; hence E' is a finite projection in $\{\pi(N_1) \otimes I_K\}'$.

Let $\{E'_n \mid n = 1, 2, \dots\}$ be a sequence of mutually orthogonal, equivalent projections in $\{\pi(N_1) \otimes I_K\}'$ such that $E'_n \sim E'$ and $\sum_{n=1}^{\infty} E'_n = I_{\mathfrak{H} \otimes K}$ where $I_{\mathfrak{H} \otimes K}$ is the identity operator on $\mathfrak{H} \otimes K$.

Now let G be the countable discrete free group with two generators, $L^2(G)$ the Hilbert space of all square integrable functions on G with respect to the Haar measure, $L^{\infty}(G)$ the algebra of all bounded functions on G .

Let $\{U, L^2(G)\}$ be the left regular representation of G on $L^2(G)$, and N be the W^* -algebra generated by $\{U(g) \mid g \in G\}$.

For $f \in L^{\infty}(G)$, we shall define a bounded operator T_f on $L^2(G)$ as follows:

$$T_f h = f o h \quad \text{for } h \in L^2(G).$$

Let $A = \{T_f \mid f \in L^{\infty}(G)\}$, then A is a commutative W^* -algebra on $L^2(G)$.

N on $L^2(G)$ is standard; hence N on $L^2(G)$ is spatial isomorphic to $\{\pi(N_1) \otimes I_K\}E'$ on $E' \mathfrak{H} \otimes K$.

Since $E'_n \sim E'$ and $\sum_{n=1}^{\infty} E'_n = I_{\mathfrak{H}} \otimes K$, we can express the W^* -algebra $\{\pi(N_1) \otimes I_K\}$ as follows:

$$\{\pi(N_1) \otimes I_K\} = \mathfrak{A} \otimes I_{K_1},$$

where $\mathfrak{A}$ is a W^* -algebra on a some Hilbert space $\mathfrak{H}_1$, and $\mathfrak{A}$ on $\mathfrak{H}_1$ is spatial isomorphic to N on $L^2(G)$, and K_1 is a Hilbert space. We shall identify $\mathfrak{A}$ with N by a spatial isomorphism.

Now we shall prove the theorem.

PROOF OF THEOREM. Now by Lemma 1, there exists a linear mapping P on $B(\mathfrak{H} \otimes K)$ to $\pi(N_1) \otimes I_K$ satisfying the properties (1)–(3).

Let τ be the trace on $\pi(N_1) \otimes I_K$.

For $f \in L^\infty(G)$, put $f_s(t) = f(s^{-1}t)$ for $s, t \in G$, then $T_{f_s}h(t) = f_s(t)h(t) = f(s^{-1}t)h(t) = U(s)T_fU(s^{-1})h(t)$ for $h \in L^2(G)$. Now we shall define a linear functional ϕ on $L^\infty(G)$ as follows: $\phi(f) = \tau(P(T_f \otimes I_{K_1}))$ for $f \in L^\infty(G)$. Then ϕ is a positive linear functional such that $\phi(1) = 1$.

Moreover,

$$\begin{aligned} \phi(f_s) &= \tau(P(U(s)T_fU(s^{-1}) \otimes I_{K_1})) \\ &= \tau(P(U(s) \otimes I_{K_1}T_f \otimes I_{K_1}U(s^{-1}) \otimes I_{K_1})) \\ &= \tau(U(s) \otimes I_{K_1}P(T_f \otimes I_{K_1})U(s^{-1}) \otimes I_{K_1}) \\ &= \tau(P(T_f \otimes I_{K_1})) \\ &= \phi(f) \quad \text{for } f \in L^\infty(G). \end{aligned}$$

Hence, ϕ will define a left invariant finitely additive measure on G . This is a contradiction (cf. [2]) and completes the proof.

Now, the following problem would be very interesting.

PROBLEM. Can we conclude that any II_1 -subfactor of the hyperfinite II_1 -factor is hyperfinite?

REFERENCES

1. J. Dixmier, "Les algèbres d'opérateurs l'espace hilbertien" in *Algèbres de von Neumann*, Gauthier-Villars, Paris, 1957.
2. J. Schwartz, *Two finite, non-hyperfinite, non-isomorphic factors*, Comm. Pure Appl. Math. 16 (1963), 19–26.
3. H. Umegaki, *Conditional expectation in an operator algebra*, Tohoku Math. J. 6 (1954), 177–181.

UNIVERSITY OF PENNSYLVANIA