THE ASYMPTOTIC BEHAVIOR OF A DECOMPOSITION DUE TO HARZHEIM¹

T. S. PITCHER

We will write $((x))$ for the fractional part of a real number x and $[x]$ for the greatest integer less than or equal to x so that $((x)) = x - [x]$. E. Harzheim has shown [2] the existence for each x in $[0, 1)$ of a sequence of nonnegative integers $\alpha_2(x), \dots, \alpha_n(x), \dots$ with

$$((x2^{\alpha_2(x)}3^{\alpha_3(x)} \dots n^{\alpha_n(x)})) < 1/n.$$

The α_n are determined inductively by setting $\alpha_n(x)$ equal to the unique integer k such that

$$\sum_{j=1}^k \frac{1}{n^j} \leq ((x2^{\alpha_2(x)} \dots (n-1)^{\alpha_{n-1}(x)})) < \sum_{j=1}^{k+1} \frac{1}{n^j}$$

(taking $\sum_{j=1}^0 = 0$). This is always possible since $\sum_{j=1}^{\infty} 1/n^j = 1/(n-1)$. Conversely, given any sequence $\alpha_2, \alpha_3, \dots$ of nonnegative integers there is an x in $[0, 1)$ with $\alpha_i(x) = \alpha_i$, $i = 2, 3, \dots$. We are concerned with the probabilistic behavior of the random variables α_n and x_n , $x_1(x) = x$, $x_n(x) = n((x2^{\alpha_2(x)} \dots n^{\alpha_n(x)}))$.

We will write λ for Lebesgue measure on $[0, 1)$. The terms independence, random variable, etc., will refer to this probability space.

LEMMA 1. *Each x_n is uniformly distributed on $[0, 1)$.*

PROOF. This is certainly true for x . If it is true for x_n then

$$\begin{aligned} \lambda\{[x \mid x_{n+1}(x) < a]\} \\ &= \lambda\left\{\bigcup_{j=0}^{\infty} [x \mid x_{n+1}(x) \leq a, \alpha_{n+1}(x) = j]\right\} \\ &= \sum_{j=0}^{\infty} \lambda\{[x \mid x_{n+1}(x) \leq a, \alpha_{n+1}(x) = j]\} \\ &= \sum_{j=0}^{\infty} \lambda\left\{\left[x \mid n \sum_{k=1}^j \left(\frac{1}{n+1}\right)^k \leq x_n < n \sum_{k=1}^j \left(\frac{1}{n+1}\right)^k + \frac{an}{(n+1)^j}\right]\right\} \\ &= \sum_{j=0}^{\infty} \frac{an}{(n+1)^j} = a. \end{aligned}$$

Received by the editors March 24, 1967.

¹ This research was supported in part by the National Science Foundation Grant GP 6216.

LEMMA 2.

$$\begin{aligned}\lambda\{[x \mid \alpha_n(x) = j]\} &= (n-1)/n^{j+1}, \quad j = 0, 1, \dots, \\ \lambda\{[x \mid \alpha_n(x) \geq j]\} &= 1/n^j, \quad j = 0, 1, \dots.\end{aligned}$$

PROOF.

$$\begin{aligned}\lambda\{[x \mid \alpha_n(x) = j]\} \\ &= \lambda\left\{[x \mid (n-1) \sum_{k=1}^j \frac{1}{n^k} \leq x_{n-1} < (n-1) \sum_{k=1}^{j+1} \frac{1}{n^k}]\right\} \\ &= \frac{n-1}{n^{j+1}}.\end{aligned}$$

The second formula is an immediate consequence of the first.

LEMMA 3. $\alpha_2(x) = \alpha_2$, $\alpha_3(x) = \alpha_3$, $\dots$, $\alpha_n(x) = \alpha_n$ if and only if

$$\begin{aligned}x &= \sum_{j=1}^{\alpha_2} \frac{1}{2^j} + 2^{-\alpha_2} \sum_{j=1}^{\alpha_3} \frac{1}{3^j} + \dots \\ &\quad + 2^{-\alpha_2} \dots (k-1)^{-\alpha_{k-1}} \sum_{j=1}^{\alpha_k} \frac{1}{k^j} + \dots \\ &\quad + 2^{-\alpha_2} \dots (n-1)^{-\alpha_{n-1}} \sum_{j=1}^{\alpha_n} \frac{1}{n^j} + 2^{-\alpha_2} \dots (n-1)^{-\alpha_{n-1}} \frac{\epsilon}{n-1}\end{aligned}$$

where $0 \leq \epsilon < (n-1)/n^{\alpha_{n-1}}$. In this case $x_n(x) = n^{\alpha_n} \epsilon / (n-1)$.

PROOF. For $n=2$ it is easily seen that $x = \sum_{j=1}^{\alpha_2} 1/2^j + \epsilon_2$ where $0 \leq \epsilon_2 < 1/2^{\alpha_2+1}$, and that

$$x_2 = 2((x \cdot 2^{\alpha_2})) = 2^{\alpha_2+1} \epsilon_2.$$

Assuming the formulas of the theorem to hold up to n and taking α_{n+1} to be any nonnegative integer we have

$$((x 2^{\alpha_2} \dots n^{\alpha_n})) = x_n(x)/n = n^{\alpha_n} \epsilon / (n-1)$$

so that $\alpha_{n+1}(x) = \alpha_{n+1}$ if and only if

$$\frac{n-1}{n^{\alpha_n}} \sum_{j=1}^{\alpha_{n+1}} \left(\frac{1}{n+1}\right)^j \leq \epsilon < \frac{n-1}{n^{\alpha_n}} \sum_{j=1}^{\alpha_{n+1}+1} \left(\frac{1}{n+1}\right)^j$$

or

$$\epsilon = \frac{n-1}{n^{\alpha_n}} \sum_{j=1}^{\alpha_{n+1}} \left(\frac{1}{n+1}\right)^j + \epsilon'$$

where

$$0 \leq \epsilon' < (n-1)/n^{\alpha_n}(1/(n+1))^{\alpha_{n+1}+1}.$$

Thus

$$x = \sum_{j=1}^{\alpha_2} \frac{1}{2^j} + \dots + 2^{-\alpha_2} \dots n^{-\alpha_n} \sum_{j=1}^{\alpha_{n+1}} \left(\frac{1}{n+1} \right)^j \\ + 2^{-\alpha_2} \dots n^{-\alpha_n} \frac{\epsilon_{n+1}}{n}$$

where

$$0 \leq \epsilon_{n+1} = n^{\alpha_n} \frac{n\epsilon'}{n-1} < \frac{n}{n+1} \left(\frac{1}{n+1} \right)^{\alpha_{n+1}}.$$

Also

$$x_{n+1}(x) = (n+1)((x2^{\alpha_2} \dots (n+1)^{\alpha_{n+1}})) \\ = (n+1)((n+1)^{\alpha_{n+1}\epsilon_{n+1}/n}) = (n+1)^{\alpha_{n+1}(n+1)/n\epsilon_{n+1}}.$$

THEOREM 1. *The α_n are independent and x_n is independent of $\alpha_1, \dots, \alpha_n$. The distributions of these random variables are given in Lemmas 1 and 2.*

PROOF. Let Ω be the subset of $[0, 1)$ where $x_n(x) \leq a$, $\alpha_2(x) = \alpha_2, \dots, \alpha_n(x) = \alpha_n$ and set

$$A = \sum_{j=1}^{\alpha_2} \frac{1}{2^j} + \dots + 2^{-\alpha_2} \dots (n-1)^{-\alpha_{n-1}} \sum_{j=1}^{\alpha_n} \frac{1}{n^j}.$$

Then

$$\lambda\{\Omega\} = \lambda\left\{ \left[x \mid x = A + 2^{-\alpha_2} \dots (n-1)^{\alpha_{n-1}} \frac{\epsilon}{n-1}, 0 \leq \epsilon \leq a \frac{n-1}{n} n^{-\alpha_n} \right] \right\} \\ = a \frac{1}{n} 2^{-\alpha_2} \dots n^{-\alpha_n} = a \prod_{j=2}^n \left(\frac{j-1}{j^{\alpha_j+1}} \right) \\ = \lambda\{[x \mid x_n(x) \leq a]\} \prod_{j=2}^n \lambda\{[x \mid \alpha_j(x) = \alpha_j]\}.$$

THEOREM 2. *For almost every x ,*

- (1) $\alpha_n(x)$ takes only the values 0 and 1 from some point on,
- (2) the value 1 occurs infinitely often,
- (3) for infinitely many n , $\alpha_k(x) = 0$ for $A_n \leq k < A_{n+1}$, A_n being any increasing sequence with $\sum_{n=2}^{\infty} A_n/A_{n+1} = \infty$.

PROOF.

$$\sum_{n=2}^{\infty} \lambda \{ [x \mid \alpha_n(x) \geq 2] \} = \sum_{n=2}^{\infty} \frac{1}{n^2} < \infty$$

so according to the Borel-Cantelli theorem the event $\alpha_n(x) \geq 2$ can occur only finitely often for almost every x . If only finitely many 1's occur then $\alpha_n(x) = 0$ from some point on and x must be rational. If we set

$$\Omega_n = [x \mid \alpha_k(x) = 0, A_n \leq k < A_{n+1}]$$

then the sets Ω_n are independent and

$$\lambda \{ \Omega_n \} = \prod_{k=A_n}^{A_{n+1}-1} \frac{k-1}{k} = \frac{A_n - 1}{A_{n+1} - 1}.$$

Thus $\sum_{n=0}^{\infty} \lambda \{ \Omega_n \} = \infty$, so by the Borel-Cantelli theorem almost every x is in infinitely many Ω_n .

If we set $Q_n(x) = 2^{\alpha_1} \cdots n^{\alpha_n}$ then, for some integer $P_n(x)$,

$$x = P_n(x)/Q_n(x) + (1/n)x_n/Q_n(x).$$

Our final theorem concerns the asymptotic distribution of x_n and Q_n .

THEOREM 3. *For every n*

$$x = P_n(x)/Q_n(x) + (1/n)x_n(x)/Q_n(x)$$

where

- (i) $P_n(x)$ is an integer,
- (ii) $Q_n(x)$ is an integer whose prime factors are all $\leq n$,
- (iii) $x_n(x)$ is uniformly distributed on $[0, 1)$ and is independent of $Q_n(x)$,

$$\begin{aligned} (iv) \quad Q_n(x) &= \left(\prod_{j=2}^n k^{1/(k-1)} \right) \\ &\quad \cdot \exp \left(\theta_n(x) \left[\sum_{j=2}^n (\log^2 k) \left(\frac{1}{k-1} + \frac{1}{(k-1)^2} \right) \right]^{1/2} \right) \end{aligned}$$

where, as $n \rightarrow \infty$,

$$\lambda \{ [x \mid \theta_n(x) \leq a] \} \rightarrow \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^a e^{-t^2/2} dt.$$

PROOF. Only (iv) needs to be proved. Straightforward calculations show

$$\begin{aligned}\int_0^1 \alpha_k(x) \log k dx &= \frac{\log k}{k-1}, \\ \int_0^1 \left(\alpha_k(x) - \frac{1}{k-1} \right)^2 \log^2 k dx &= (\log^2 k) \left(\frac{1}{k-1} + \left(\frac{1}{k-1} \right)^2 \right), \\ \int_0^1 \left| \alpha_k(x) - \frac{1}{k-1} \right|^3 \log^3 k dx &\leq C \frac{\log^3 k}{k-1}.\end{aligned}$$

Since

$$\tau_n^3 = \sum_{k=2}^n \int_0^1 \left| \alpha_k(x) - \frac{1}{k-1} \right|^3 \log^3 k dx \leq C_1 \log^4 n$$

and

$$\sigma_n^2 = \sum_{k=2}^n \int_0^1 \left| \alpha_k(x) - \frac{1}{k-1} \right|^2 \log^2 k dx \geq C_2 \log^3 n,$$

$\tau_n/\sigma_n \rightarrow 0$ as $n \rightarrow \infty$ so we can apply Liapounov's theorem with $\delta=1$ [1, Theorem 4.4, p. 144] to conclude that

$$\theta_n = \frac{1}{\sigma_n} \sum_{k=2}^n \left(\alpha_k(x) - \frac{1}{k-1} \right) \log k$$

has the asymptotic distribution required in (iv). This gives

$$\log Q_n = \sum_{k=2}^n \frac{\log k}{k-1} + \sigma_n \theta_n$$

and

$$Q_n = \left(\prod_{k=2}^n k^{1/(k-1)} \right) \exp(\sigma_n \theta_n).$$

BIBLIOGRAPHY

1. J. L. Doob, *Stochastic processes*, Wiley, New York, 1953.
2. E. Harzheim, *An elementary number theoretical remark*, Proc. Amer. Math. Soc. **17** (1966) 534-535.

UNIVERSITY OF SOUTHERN CALIFORNIA