A NOTE ON "NTH ROOTS OF OPERATORS"

M. FINKELSTEIN AND A. LEBOW

M. R. Embry has proved the following theorem [1]:

THEOREM [E]. "Let A and E be bounded operators on a Banach space, and suppose $\sigma(A) \cap \sigma(e^{2\pi i k/n} A) = \emptyset$ for $k=1, 2, \dots, n-1$. Then if E commutes with A^n , it commutes with A ."

This result can be obtained as a corollary to the following theorem: *Notation (as in [2]).* Let T be a bounded operator on a Banach space. Let $\mathfrak{F}(T)$ = the functions holomorphic on some neighborhood of $\sigma(T)$. Let $f \in \mathfrak{F}(T)$.

As usual,

$$f(T) = (2\pi i)^{-1} \int_c f(\lambda)(\lambda - T)^{-1} d\lambda.$$

THEOREM. *In order that T be representable as a function of $f(T)$ it is sufficient that (1) f is 1-1 on $\sigma(T)$, and (2) $f' \neq 0$ on $\sigma(T)$. Consequently if E is a bounded operator and commutes with $f(T)$ then it commutes with T .*

PROOF. (1) and (2) ensure that f is 1-1 on a neighborhood of $\sigma(T)$, for if not there exist $\{z_n\}$, $\{z'_n\}$, $z_n \neq z'_n$, such that $f(z_n) = f(z'_n)$ and these may be chosen so that $z_n \rightarrow z_0$, $z'_n \rightarrow z'_0$, $z_0 \in \sigma(T)$, $z'_0 \in \sigma(T)$. Now $z_0 = z'_0$, else (1) is violated. But then

$$0 = \frac{f(z_n) - f(z'_n)}{z_n - z'_n} \rightarrow f'(z_0)$$

which contradicts (2). If f is 1-1 on a neighborhood of $\sigma(T)$, then $f^{-1} \in \mathfrak{F}(f(T))$, and Theorem 12 [2, p. 570] yields $T = f^{-1}(f(T))$.

If f fails to satisfy (1) or (2), an E and T for which E commutes with $f(T)$ but not with T can be found in 2×2 matrices.

Note that z^n satisfies (1) and (2) precisely if the condition in Theorem E is satisfied.

REFERENCES

1. M. R. Embry, *Nth roots of operators*, Proc. Amer. Math. Soc. **19** (1968), 63-68.
2. N. Dunford and J. T. Schwartz, *Linear operators*. I, Interscience, New York, 1958.

UNIVERSITY OF CALIFORNIA, IRVINE

Received by the editors March 6, 1968.