NORM PRESERVING EXTENSIONS OF LINEAR TRANSFORMATIONS ON HILBERT SPACES

MICHAEL G. CRANDALL¹

Introduction. Let H be a Hilbert space and let D be a closed proper subspace of H . Let J_0 be a linear contraction on D to H . The problem of characterizing the contractions on all of H which extend J_0 is directly related to the extension problems for unbounded transformations posed and treated by M. G. Krein [2] and R. S. Phillips [3]. In §1 of this paper we establish the following solution of this problem: Let P be the orthogonal projection of H on D and $J_1 = J_0 P$. The contractions on all of H which extend J_0 are precisely the transformations of the form $J_1 + (I - J_1 J_1^*)^{1/2} B$, where B is any contraction which annihilates D . §2 is a short outline of the relationship of this result to the extension problem for dissipative transformations treated by Phillips in [3] and [4]. §3 contains some remarks concerning the case where J_0 is symmetric and one asks for the selfadjoint contractions which extend J_0 , which is essentially the situation treated by Krein [2]. In particular, we obtain the Krein extension of a symmetric contraction to a selfadjoint contraction by, we feel, a less elegant but more illuminating argument than Krein's original one.

1. The representation theorem. All transformations considered here are linear. The domain, range and null space of a transformation L will be denoted by $D(L)$, $R(L)$, and $N(L)$ respectively. Our goal is to prove the following theorem:

THEOREM I. *Let H be a Hilbert space. Let D be a closed proper subspace of H and let J_0 be a contraction on D to H . Let P be the orthogonal projection of H on D and set*

$$(1.1) \quad J_1 = J_0 P.$$

Then the contractions J on H to H which extend J_0 are precisely the transformations of the form

$$(1.2) \quad J = J_1 + (I - J_1 J_1^*)^{1/2} B$$

where B is any contraction on H which annihilates D .

REMARK. The correspondence (1.2) between contractions J which

Presented to the Society, June 29, 1968; received by the editors May 17, 1968.

¹ This report is supported in part by the National Science Foundation Grant GP 7458.

extend J_0 and contractions B which annihilate D becomes biunique if we further require that $R(B) \subset [R(T)]$, where we have set

$$(1.3) \quad T = (I - J_1 J_1^*)^{1/2}.$$

and $[R(T)]$ denotes the closure of $R(T)$.

PROOF OF THEOREM I. For convenience we set

$$(1.4) \quad E(J_0) = \{J: J \text{ is a contraction on all of } H \text{ and } J \text{ extends } J_0\}.$$

Let $J \in E(J_0)$. Clearly then $JP = J_1 = J_0 P$ and we have

$$(1.5) \quad J = JP + J(I - P) = J_1 + J(I - P) \quad \text{for } J \text{ in } E(J_0).$$

The decomposition (1.5) shows that every element of $E(J_0)$ has the form $J_1 + K$ where K is a contraction annihilating D . Conversely, any transformation of this form extends J_0 . Hence $E(J_0)$ consists of those sums $J_1 + K$, where K is a contraction annihilating D , which are contractions. The theorem is therefore an immediate consequence of the following lemma.

LEMMA I. *Let the assumptions of Theorem I be satisfied. Then, if K is a contraction on H which annihilates D , $J_1 + K$ is a contraction if and only if K has the form $K = TB$, where T is given by (1.3), for some contraction B which annihilates D .*

PROOF OF LEMMA I. An attempt to prove Lemma I directly would lead to some difficulties, however, passing to the adjoint allows a trivial proof. We first show the sufficiency. Let B be a contraction annihilating D . $J_1 + TB$ is a contraction if $J_1^* + B^*T$ is a contraction (note that T is selfadjoint). Moreover, $N(J_1) \supset D^\perp$ (the orthogonal complement of D in H) and $N(B) \supset D$, so $R(J_1^*)$ and $R(B^*)$ are orthogonal. Hence, since B^* is a contraction and since $\|Tw\|^2 = \|w\|^2 - \|J_1^*w\|^2$ for w in H by (1.3), where $\|\cdot\|$ is the norm in H , we have

$$\begin{aligned} \|(J_1^* + B^*T)w\|^2 &= \|J_1^*w\|^2 + \|B^*Tw\|^2 \leq \|J_1^*w\|^2 + \|Tw\|^2 \\ &= \|J_1^*w\|^2 + \|w\|^2 - \|J_1^*w\|^2 = \|w\|^2 \end{aligned}$$

for w in H , and $J_1^* + B^*T$ is a contraction.

Conversely, if K annihilates D and $J_1 + K$ is a contraction, then $R(J_1^*)$ and $R(K^*)$ are orthogonal and $J_1^* + K^*$ is a contraction, so we have

$$(1.6) \quad \|(J_1^* + K^*)w\|^2 = \|J_1^*w\|^2 + \|K^*w\|^2 \leq \|w\|^2 \quad \text{for } w \text{ in } H.$$

It follows from (1.6) that

$$(1.7) \quad \|K^*w\|^2 \leq \|Tw\|^2 \quad \text{for } w \text{ in } H.$$

According to (1.7) the map L on $R(T)$ to $R(K^*)$ defined by $LTw = K^*w$ is a contraction. We extend L to $[R(T)]$ by continuity and then to all of H by setting it equal to zero on $R(T)^\perp$. Then we have $LT = K^*$ and $K = TL^*$. Since $R(L) \subset R(K^*) \subset D^\perp$, $N(L^*) \supset D$ and, setting $B = L^*$, K has the form claimed by the lemma. This completes the proof of Lemma I and therefore Theorem I is also established.

REMARK. The proof of Theorem I may easily be adapted to provide a proof of the more general result stated below.

THEOREM I'. *Let H_1 and H_2 be Hilbert spaces. Let D be a closed proper subspace of H_1 and let J_0 be a bounded transformation on D to H_2 . Let P be the orthogonal projection of H_1 on D and set $J_1 = J_0P$. Then the bounded transformations on H_1 to H_2 of norm not exceeding $k\|J_0\|$, $k \geq 1$, which extend J_0 are precisely the transformations of the form $J_1 + (k^2\|J_0\|^2 - J_1J_1^*)^{1/2}B$, where B is a contraction on H_1 to H_2 which annihilates D .*

2. The extension problem for dissipative transformations. Let A be a transformation on $D(A) \subset H$ into H . A is called *dissipative* if

$$(2.1) \quad (u, Au) + (Au, u) \leq 0 \quad \text{for } u \text{ in } D(A).$$

If A is dissipative and has no proper dissipative extensions, it is called *maximal dissipative*. We consider the problem of finding all the maximal dissipative transformations which extend a given dissipative A . If A is dissipative then, in view of (2.1),

$$(2.2) \quad \begin{aligned} \|(I - A)u\|^2 &= \|u\|^2 + \|Au\|^2 - 2 \operatorname{Re}(u, Au) \\ &\geq \|u\|^2 + \|Au\|^2 \geq \|u\|^2 + \|Au\|^2 + 2 \operatorname{Re}(u, Au) \\ &= \|(I + A)u\|^2 \quad \text{for } u \text{ in } D(A). \end{aligned}$$

It follows from the first inequality of (2.2) that $I - A$ is one-to-one and that $R(I - A)$ is a closed subspace of H if A is a closed transformation, and from the inequality of the extreme members in (2.2) it follows that the transformation J_0 defined by

$$(2.3) \quad D(J_0) = R(I - A), \quad J_0(I - A)u = (I + A)u \quad \text{for } u \text{ in } D(A)$$

is a contraction. J_0 is called the Cayley transform of A . If $D(A)$ is dense in H , then there is a one-to-one correspondence between the maximal dissipative extensions of A and the elements of $E(J_0)$, that is the contractions which extend J_0 to H . Given J in $E(J_0)$ the corresponding maximal dissipative extension of A is the dissipative transformation A_J whose Cayley transform is J : It is defined by

$$(2.4) \quad D(A_J) = R(I + J), \quad A_J(I + J)u = (J - I)u \quad \text{for } u \text{ in } H.$$

For the straightforward proof of this see [3]. Theorem I, therefore, allows us to characterize the maximal dissipative extensions of a given dissipative transformation A . We proceed with this below.

Let A be a closed densely defined dissipative transformation and let J_0 be the Cayley transform of A . Let P be the orthogonal projection of H on $D(J_0) = R(I - A)$ and let $J_1 = J_0 P$. J_1 is determined by A as follows: $R(I - A)$ is closed and $R(I - A)^\perp = N(I - A^*)$. Every element w of H has a unique decomposition into a sum $w = (I - A)u + v$ with u in $D(A)$ and v in $N(I - A^*)$ and we have

$$(2.5) \quad J_1((I - A)u + v) = (I + A)u \text{ for } u \text{ in } D(A) \text{ and } v \text{ in } N(I - A^*).$$

J_1^* is related in a somewhat similar way to A^* . Since $N(I - A) = \{0\}$ and $R(I - A)$ is closed, $R(I - A^*) = H$. We claim

$$(2.6) \quad J_1^*(I - A^*)u = P(I + A^*)u \text{ for } u \text{ in } D(A^*).$$

In view of (2.5), the validity of (2.6) is equivalent to

$$\begin{aligned} ((I + A)u, (I - A^*)w) &= ((I - A)u + v, P(I + A^*)w) \\ &\text{for } u \text{ in } D(A), \quad w \text{ in } D(A^*) \quad \text{and} \quad v \text{ in } N(I - A^*), \end{aligned}$$

which is true in view of

$$\begin{aligned} &((I - A)u + v, P(I + A^*)w) \\ &= (P((I - A)u + v), (I + A^*)w) = ((I - A)u, (I + A^*)w) \end{aligned}$$

for u in $D(A)$, w in $D(A^*)$ and v in $N(I - A^*)$.

We have now, as an immediate corollary of Theorem I,

COROLLARY I. *Let A be a closed densely defined dissipative transformation. Then there is a one-to-one correspondence between contractions B on $N(I - A^*)$ to $[R(T)]$ where T is given by (1.3), J_1 by (2.5) and J_1^* by (2.6), and the maximal dissipative extensions of A . Given a contraction B on $N(I - A^*)$ to $[R(T)]$ the corresponding extension A_B of A is defined by*

$$\begin{aligned} D(A_B) &= D(A) + (I + TB)N(I - A^*), \\ A_B(u + (I + TB)v) &= Au + (TB - I)v \\ &\text{for } u \text{ in } D(A) \quad \text{and} \quad v \text{ in } N(I - A^*). \end{aligned}$$

REMARK 1. The main shortcoming of this result with regard to applications lies in the difficulty of computing the action of the transformation T given the transformation A . With further restrictions on the transformation A more convenient descriptions of its maximal dissipative extensions can be given. See M. G. Crandall and R. S. Phillips [1].

REMARK 2. A more general problem, including the extension problem for dissipative transformations, is treated by Phillips [3], [4] and [5]. In Theorem 5.3 of [4] Phillips gives a solution of this problem, the formulation of which involves completing a certain Hilbert space in a new metric. We remark that Theorem I' may be applied directly to this more general problem, giving a new solution which does not involve any "completions."

3. **The symmetric case.** We adopt here the notations of §1. We now assume that J_0 is symmetric, i.e.,

$$(3.1) \quad (J_0 u, v) = (u, J_0 v) \quad \text{for } u \text{ and } v \text{ in } D,$$

and ask for those J in $E(J_0)$ which are selfadjoint. See Krein [2] for the relationship of this question to the problem of characterizing the semibounded selfadjoint extensions of semibounded symmetric transformations. Let $J_1 = J_0 P$ as before. Clearly J_0 is symmetric if and only if PJ_1 is selfadjoint. Let $J = J_1 + TB$ be an element of $E(J_0)$; B is a contraction annihilating D and T is given by (1.3). It is not easy to characterize those B for which J is selfadjoint, but we can easily characterize those B for which J^* extends J_0 , and if J and J^* both extend J_0 then $(J + J^*)/2$ is a selfadjoint extension of J_0 . We let $P_1 = (I - P)$; P_1 is the orthogonal projection of H on $D^\perp$. Then

$$\begin{aligned} (J_1 + TB)^* &= ((P + P_1)(J_1 + TB)(P + P_1))^* \\ &= PJ_1 + PJ_1^* P_1 + B^* TP + B^* TP_1, \end{aligned}$$

because PJ_1 is selfadjoint and $J_1 P_1 = BP = 0$ by assumption, and it follows that $(J_1 + TB)^*$ extends J_0 if and only if

$$PJ_1 + B^* TP = J_1$$

or

$$(3.2) \quad B^* TP = P_1 J_1.$$

Equation (3.2) determines the action of B^* on $[TD]$. As noted below, (3.2) is consistent with the requirement that B^* be a contraction and Theorem I can be used to characterize those contractions B for which (3.2) is satisfied, which we leave to the reader. Here we restrict ourselves to noting that if L is defined by

$$(3.3) \quad \begin{aligned} L^* T u &= P_1 J_1 u & \text{for } u \text{ in } D \\ L^* v &= 0 & \text{for } v \text{ in } (TD)^\perp \end{aligned}$$

then L^* (and hence L) is a contraction for $J_1^* P = (PJ_1)^* = PJ_1$ implies

$$\begin{aligned}
\|Tu\|^2 &= \|u\|^2 - \|J_1^*u\|^2 = \|u\|^2 - \|J_1^*Pu\|^2 = \|u\|^2 - \|PJ_1u\|^2 \\
&= \|u\|^2 - \|J_1u\|^2 + \|P_1J_1u\|^2 \\
&\geq \|P_1J_1u\|^2 \quad \text{for } u \text{ in } D,
\end{aligned}$$

and $R(L^*) \subset D^\perp$ implies $N(L) \supset D$. By the above arguments it follows that $(J_1 + J_1^* + TL + L^*T)/2$ is a selfadjoint contraction which extends J_0 ; it is the extension found by Krein [2].

REMARK. If W is a bounded selfadjoint transformation on H and

$$(3.4) \quad J_0W = \varepsilon WJ_0 \quad \text{for } \varepsilon = 1 \text{ or } \varepsilon = -1,$$

it was noticed by Phillips [5, Lemma 3.1] that the Krein extension of J_0 will also satisfy (3.4) in place of J_0 . Our construction makes this obvious, for if (3.4) holds then W commutes with P and P_1 , so J_1 satisfies (3.4) and therefore J_1^* satisfies (3.4) and therefore $(I - J_1J_1^*)^{1/2}$ commutes with W and finally the transformations L^* and L defined by (3.3) satisfy (3.4). It follows that the Krein extension of J_0 will also satisfy (3.4).

REFERENCES

1. M. G. Crandall and R. S. Phillips, *On the extension problem for dissipative operators*, J. Functional Analysis 2 (1968), 147-176.
2. M. A. Krein, *The theory of self-adjoint extensions of semi-bounded Hermitian transformations and its applications*. I, Mat. Sb. 20 (62) (1947), 431-495. (Russian)
3. R. S. Phillips, *Dissipative operators and hyperbolic systems of partial differential equations*, Trans. Amer. Math. Soc. 90 (1959), 193-254.
4. ———, *On dissipative operators*, Van Nostrand Math. Studies Series (to appear).
5. ———, *The extension of dual subspaces invariant under an algebra*, Proc. International Sympos. on Linear Spaces, Israel, (1960), Jerusalem Academic Press, 1961, pp. 366-398.

STANFORD UNIVERSITY