ON THE EXISTENCE OF c -POINTS IN $\beta N \setminus N$

NEIL HINDMAN¹

R. S. Pierce has asked in [2] whether it is possible to show, without using the continuum hypothesis, that there are points of $\beta N \setminus N$ which lie simultaneously in the closure of three pairwise disjoint open sets. Such points are called, in his terminology, 3-points of $\beta N \setminus N$. In general, for any cardinal number n , and for any topological space X , a point x in X is called an n -point if it lies in the closure of each of n pairwise disjoint open subsets of X .

It is shown here in §1, as a corollary to a more general theorem, that, without appeal to the continuum hypothesis, not only are there 3-points in $\beta N \setminus N$, but in fact there are c -points in $\beta N \setminus N$. It is further shown in §2 that, if the continuum hypothesis is assumed, *each* point of $\beta N \setminus N$ is a c -point. These results seem particularly striking since $\beta N \setminus N$ is an F' -space, i.e., disjoint cozero sets in $\beta N \setminus N$ have disjoint closures. (A cozero set in a space X is a preimage by a continuous real valued function of an open set.) It is well known that if α is any cardinal greater than c then there do not exist α -points in $\beta N \setminus N$, since in fact, if $\mathcal{U}$ is a collection of pairwise disjoint open sets in $\beta N \setminus N$ then $|\mathcal{U}| \leq c$.

1. Results without the continuum hypothesis. Let A^p be the ultrafilter on a discrete space D associated with the point p in βD . The ultrafilter A^p on D is called uniform if each element of A^p has cardinality $|D|$. Define $\mu D = \{p \in \beta D : A^p \text{ is uniform}\}$. Note that $\mu N = \beta N \setminus N$.

We include here for completeness the following facts about βD for discrete D . Our notation and general point of view are those of the Gillman and Jerison textbook [1], to which the reader is referred for proofs and additional discussion. For each $Z \subset D$, we have $\text{cl}_{\beta D} Z = \{p \in \beta D : Z \in A^p\}$. Basic neighborhoods of a point p in $\beta D \setminus D$ are of the form $(\text{cl}_{\beta D} Z) \setminus D$ where $Z \in A^p$.

1.1 LEMMA. *Let D be a discrete space of cardinality m , where $m \geq \aleph_0$. Let $\mathcal{A}$ be an infinite collection of subsets of D with $|A| = m$ for each*

Received by the editors July 15, 1968.

¹ The author wishes to express his gratitude to R. S. Pierce for his encouragement, and to W. W. Comfort for calling the problem to his attention and for some helpful suggestions. The author is a National Science Foundation Trainee. He also gratefully acknowledges support received from the National Science Foundation via grant NSF-GP-8357.

A in $\mathcal{Q}$ and with $|A_1 \cap A_2| < m$ for each distinct pair of elements A_1, A_2 of $\mathcal{Q}$. Then there is a uniform ultrafilter A^p on D such that $|\{A \in \mathcal{Q}: |Z \cap A| = m\}| = |\mathcal{Q}|$ for each Z in A^p .

PROOF. For each A in $\mathcal{Q}$ choose x_A in $\text{cl}_{\beta D} A \cap \mu D$ and let $B = \{x_A: A \in \mathcal{Q}\}$. Now if $A_1 \neq A_2$ then $x_{A_1} \neq x_{A_2}$, since x_{A_1} and x_{A_2} are elements of μD and $|A_1 \cap A_2| < m$. Thus $|B| = |\mathcal{Q}|$. Now μD is closed in βD , hence is compact. (For if $p \in \beta D \setminus \mu D$ then there exists Z in A^p with $|Z| < m$, and $\text{cl}_{\beta D} Z$ is a neighborhood of p missing μD .) Clearly then there is a point p in μD such that each neighborhood of p contains $|\mathcal{Q}|$ elements of B . (For if not, pick a neighborhood of each point of μD which contains less than $|\mathcal{Q}|$ elements of B . From the existence of a finite subcover it follows immediately that $|B| < |\mathcal{Q}|$, a contradiction.) But then A^p is the desired ultrafilter, for if $|Z \cap A| < m$ then $x_A \notin \text{cl}_{\beta D} Z \cap \mu D$.

It follows from Theorem 7 of [3] that if m is a cardinal number such that $2^n \leq m$ for all $n < m$ and if D is a set of cardinality m , then there is a family $\mathcal{Q}$ of subsets of D , each of cardinality m , such that $|\mathcal{Q}| = 2^m$ and $|A_1 \cap A_2| < m$ for each pair A_1, A_2 of elements of $\mathcal{Q}$. Note that if the generalized continuum hypothesis holds then each infinite cardinal m has the above mentioned property. At any rate there are cofinally many cardinals with this property. (For let n_0 be any cardinal and define inductively $n_{i+1} = 2^{n_i}$. Then $m = \sup\{n_i: i \in \omega_0\}$ has the given property and is greater than n_0 .)

1.2 LEMMA. Let D be a discrete space of cardinality m where $2^n \leq m$ for all $n < m$. Let $\mathcal{Q}$ be a collection of subsets of D as given above. Let A^p be a uniform ultrafilter as given by Lemma 1.1, the elements Z_α of A^p being indexed by the ordinals less than 2^m . Then it is possible to choose, for each $\alpha < 2^m$, a subset X_α of Z_α so that $|X_\alpha| = m$ and so that $|X_\alpha \cap X_\gamma| < m$ whenever $\alpha < \gamma < 2^m$.

PROOF. For Z_1 pick A_1 in $\mathcal{Q}$ such that $|Z_1 \cap A_1| = m$. Assume that for $\sigma < \alpha$ we have chosen $A_\sigma \in \mathcal{Q}$ such that $|Z_\sigma \cap A_\sigma| = m$ and $A_\sigma \neq A_\gamma$ for all $\gamma < \sigma$. Now $|\{A \in \mathcal{Q}: |A \cap Z_\alpha| = m\}| = 2^m$ and $\alpha < 2^m$ so there is $A_\alpha \in \mathcal{Q}$ such that $|A_\alpha \cap Z_\alpha| = m$ and $A_\sigma \neq A_\alpha$ for all $\sigma < \alpha$. Let $X_\alpha = A_\alpha \cap Z_\alpha$.

1.3 THEOREM. Let D be a discrete space of cardinality m where $2^n \leq m$ for all $n < m$. Then there is a 2^m -point (i.e. a point in the closure of each of 2^m pairwise disjoint open sets) in μD .

PROOF. Let $\mathcal{Q}, A^p = \{Z_\alpha\}_{\alpha < 2^m}$ and $X_\alpha \subset Z_\alpha$ be as in Lemma 1.2. Now $|X_\alpha| = m$ so there is a collection $\{X_{\tau_\alpha}\}_{\tau_\alpha < 2^m}$ of subsets of X_α

such that $|X_{\tau\alpha}| = m$ for all $\tau < 2^m$ and such that $|X_{\tau\alpha} \cap X_{\sigma\alpha}| < m$ if $\tau \neq \sigma$. For each $\tau < 2^m$ let $U_\tau = (U_{\alpha < 2^m} \text{cl}_{\beta D} X_{\tau\alpha}) \cap \mu D$. Recall that $\text{cl}_{\beta D} X_{\tau\alpha}$ is open in βD , so that U_τ is open. We assert that the family $\{U_\tau\}_{\tau < 2^m}$ is as desired. That is, it is a pairwise disjoint family and the closure of each member contains p . Suppose first that $\sigma \neq \tau$ and $q \in U_\sigma \cap U_\tau$. Then there is a pair α_1, α_2 such that $q \in \text{cl}_{\beta D} X_{\sigma\alpha_1} \cap \text{cl}_{\beta D} X_{\tau\alpha_2} \cap \mu D$. If $\alpha_1 \neq \alpha_2$ then $X_{\sigma\alpha_1} \cap X_{\tau\alpha_2} \subset X_{\alpha_1} \cap X_{\alpha_2}$ and $|X_{\alpha_1} \cap X_{\alpha_2}| < m$. If $\alpha_1 = \alpha_2$ then $|X_{\sigma\alpha_1} \cap X_{\tau\alpha_2}| < m$. So in fact, in either case $|X_{\sigma\alpha_1} \cap X_{\tau\alpha_2}| < m$. But since $q \in \text{cl}_{\beta D} X_{\sigma\alpha_1} \cap \text{cl}_{\beta D} X_{\tau\alpha_2}$ we have that $X_{\sigma\alpha_1} \cap X_{\tau\alpha_2} \in A^q$, contrary to the fact $q \in \mu D$.

Finally let $Z_\alpha \in A^p$. Now $\text{cl}_{\beta D} Z_\alpha \cap \mu D$ is a basic neighborhood of p in μD so we need only show that $\text{cl}_{\beta D} Z_\alpha \cap \mu D \cap U_\tau \neq \emptyset$ for all $\tau < 2^m$. Since $|X_{\tau\alpha}| = m$ and $X_{\tau\alpha} \subset X_\alpha \subset Z_\alpha$ it follows that $|X_{\tau\alpha} \cap Z_\alpha| = m$, so there is $q \in \text{cl}_{\beta D} Z_\alpha \cap \text{cl}_{\beta D} X_{\tau\alpha} \cap \mu D \subset \text{cl}_{\beta D} Z_\alpha \cap U_\tau \cap \mu D \neq \emptyset$.

1.4 COROLLARY. *There is a c-point in $\beta N \setminus N$.*

PROOF. $2^n \leq \aleph_0$ for all $n < \aleph_0$ and $\beta N \setminus N = \mu N$.

2. Results assuming the continuum hypothesis. We show here that, if the continuum hypothesis is assumed, not only do there exist c-points of $\beta N \setminus N$ as shown above, but in fact *each* point of $\beta N \setminus N$ is a c-point of $\beta N \setminus N$.

2.1 LEMMA. *Assume the continuum hypothesis and let A^p be a free (i.e. uniform) ultrafilter on N . Order the elements of A^p by the ordinals less than ω_1 . Then there is a choice of $X_\alpha \subset Z_\alpha$ for each $Z_\alpha \in A^p$ such that $|X_\alpha| = \aleph_0$ and $|X_\alpha \cap X_\beta| < \aleph_0$ if $\alpha \neq \beta$.*

PROOF. Let X_1 be any infinite subset of Z_1 which is not in A^p . Let $\alpha < \omega_1$ and assume that for each $\sigma < \alpha$ we have chosen $X_\sigma \subset Z_\sigma$ so that $X_\sigma \notin A^p$ and $|X_\sigma \cap X_\gamma| < \aleph_0$ for all $\gamma < \sigma$. Let $B = \{\sigma < \alpha: |X_\sigma \cap Z_\alpha| = \aleph_0\}$. If B is finite then $(Z_\alpha \setminus \bigcup_{\sigma \in B} X_\sigma) \in A^p$ and we may let X_α be any infinite subset of $Z_\alpha \setminus \bigcup_{\sigma \in B} X_\sigma$ which is not in A^p . Then for $\sigma < \alpha$ we have $|X_\sigma \cap X_\alpha| < \aleph_0$. If B is infinite it is countable so we may write $B = \{\sigma_n\}_{n=1}^\infty$. Let $x_{\alpha k} \in (X_{\sigma_k} \cap Z_\alpha) \setminus \bigcup_{j < k} X_{\sigma_j}$. This is possible since $|X_{\sigma_k} \cap Z_\alpha| = \aleph_0$ and $|X_{\sigma_j} \cap X_{\sigma_k}| < \aleph_0$ for $j < k$. Let X_α be any infinite subset of $\{x_{\alpha k}\}_{k=1}^\infty$ which is not an element of A^p . Then $|X_\alpha \cap X_{\sigma_k}| \leq k < \aleph_0$, and if $\sigma \notin B$ then $|X_\sigma \cap X_\alpha| \leq |X_\sigma \cap Z_\alpha| < \aleph_0$.

2.2 THEOREM. *Assume the continuum hypothesis and let $p \in \beta N \setminus N$. Then p is a c-point of $\beta N \setminus N$.*

PROOF. Let $X_\alpha \subset Z_\alpha$ be as in Lemma 2.1 for each Z_α in A^p . The rest of the proof is the same as that of Theorem 1.3.

3. Remarks. It is an easy consequence of Theorem 1 of [3] that for any discrete space D of cardinality m where $m \geq \aleph_0$, μD does not have n -points for any $n > 2^m$ since in fact μD does not have n pairwise disjoint open sets.

It is not possible to directly generalize the proof of Lemma 2.1 to obtain the corresponding statement about μD for discrete spaces D of higher cardinality. Thus the following question remains open: Even assuming the generalized continuum hypothesis, is each point of D a $2^{|D|}$ -point for an infinite discrete space D ?

The author has also been unable to determine if the continuum hypothesis is in fact a necessary condition for Theorem 2.2.

REFERENCES

1. Leonard Gillman and Meyer Jerison, *Rings of continuous functions*, Van Nostrand, Princeton, N. J., 1960.
2. R. S. Pierce, *Modules over commutative regular rings*, Mem. Amer. Math. Soc. No. 70, (1967).
3. Alfred Tarski, *Sur la décomposition des ensembles en sous-ensembles presque disjoints*, Fund. Math. 12 (1928), 188–205.

WESLEYAN UNIVERSITY