SHORTER NOTES

The purpose of this department is to publish very short papers of an unusually elegant and polished character, for which there is no other outlet.

ON SEMIGROUPS NEAR THE IDENTITY

W. WILS

In a number of recent papers [1], [2], [3], and [4] the question of the existence of nontrivial semigroups of operators on a normed space "close to the identity" is considered. In this note a simple theorem is proved from which all known results readily follow.

It is a pleasure to thank R. Hirschfeld for mentioning the problem.

THEOREM. *Let G be a group and T an endomorphism of G . Suppose there exists a set F of homomorphisms from G into the additive group $\mathbb{C}$ such that:*

1. $f(T^n g) = o(n)$ for every $f \in F, g \in G$.
2. For every $0 \neq b \in G$, there is a $f_b \in F$ with

$$\limsup \frac{1}{n} \left| \sum_{k=0}^{n-1} f_b(T^k b) \right| > 0.$$

Then $Tg = g$ for all $g \in G$.

PROOF. Let $b = Tg - g$. For $f \in F$ we get

$$\frac{1}{n} \sum_{k=0}^{n-1} f(T^k b) = \frac{1}{n} (f(T^n g) - f(g))$$

and thus

$$\lim \frac{1}{n} \left(\sum_{k=0}^{n-1} f(T^k b) \right) = 0.$$

By 2 this implies $b = 0$. Q.E.D.

COROLLARY 1. *Let T be an additive operator on a locally convex space X , with topological dual X' , such that*

1. $\langle T^n x, x' \rangle = o(n)$ for every $x \in X, x' \in X'$.
2. For every $0 \neq x \in X$ there is an $x' \in X'$ with $\langle x, x' \rangle \neq 0$ and

$$\liminf \left| \frac{1}{n} \sum_{k=0}^{n-1} \langle T^k x, x' \rangle - \langle x, x' \rangle \right| < |\langle x, x' \rangle|.$$

Then $T = I$.

Received by the editors September 18, 1968.

PROOF. We apply the theorem to $X = G$, $F = X'$ and T .

We note that it is even not supposed that T is continuous.

COROLLARY 2. *Let T be a continuous operator on a normed space X such that*

1. $\|T^n\| = o(n)$.
2. $\liminf (1/n) \left\| \sum_{k=0}^{n-1} (T^k - I) \right\| < 1$.

Then $T = I$.

PROOF. Let $0 \neq x \in X$. There is a $0 \neq x' \in X'$ with $\langle x, x' \rangle = \|x\| \|x'\|$. Then

$$\liminf \left| \frac{1}{n} \sum_{k=0}^{n-1} \langle (T^k - I)x, x' \rangle \right| < \|x\| \|x'\| = \langle x, x' \rangle$$

so that Corollary 1 applies.

REFERENCES

1. R. H. Cox, *Matrices all of whose powers lie close to the identity*, Amer. Math. Monthly **73** (1966), 813.
2. R. Hirschfeld, *On semi-groups in Banach algebras close to the identity*, Proc. Japan Acad. **44** (1968), 55.
3. M. Nakamura and M. Yoshida, *On a generalization of a theorem of Cox*, Proc. Japan Acad. **43** (1967), 108–110.
4. L. J. Wallen, *On the magnitude of $x^n - 1$ in a normed algebra*, Proc. Amer. Math. Soc. **18** (1967), 956.

CATHOLIC UNIVERSITY OF NIJMEGEN, THE NETHERLANDS