ON A PROBLEM OF ERDÖS

MING-CHIT LIU

Erdős posed the following problem in [2, pp. 52–53]: Let $\{z_\nu\}$ be a sequence with $|z_\nu| = 1$ and put

$$B_k = \limsup_{m \rightarrow \infty} \left| \sum_{\nu=1}^m z_\nu^k \right|.$$

Is it true that $\limsup_{k \rightarrow \infty} B_k = \infty$?

In a subsequent paper [3] Erdős showed that there is an absolute constant C such that we always have $B_k/\log k > C$ infinitely often. He also conjectured that $B_k/k > C$ for infinitely many values of k . Recently, Clunie [1] proved that

$$\limsup_{k \rightarrow \infty} B_k/k^{1/2} > 0,$$

and there is a sequence $\{z_\nu\}$ such that $B_k \leq k$ ($k = 1, 2, \dots$).

In this paper we shall prove the following theorem.

THEOREM. *Let $\{z_\nu\}$ be any sequence with $|z_\nu| = 1$. If $\{z_\nu\}$ contains only a finite number of distinct elements, then for any $\delta > 0$,*

$$\limsup_{k \rightarrow \infty} B_k/k^{1-\delta} > \frac{1}{5}.$$

Let $z_\nu = \exp[i\phi_\nu]$ ($0 \leq \phi_\nu < 2\pi$). It is easy to see that for any sequence $\{z_\nu\}$ containing finite distinct elements there exists some $\epsilon > 0$ which depends on the sequence $\{z_\nu\}$ only, such that either

$$\phi_\nu - \phi_{\nu'} = 0 \quad \text{or} \quad |\phi_\nu - \phi_{\nu'}| \geq \epsilon > 0$$

for all positive integers ν, ν' . So in what follows we shall use ϵ to denote the maximum positive constant satisfying the above condition.

In order to prove our Theorem we need two lemmas.

LEMMA 1. *Let $\{\zeta_\nu\}$ be any sequence containing only a finite number of distinct elements with $|\zeta_\nu| = 1$ ($\nu = 1, 2, \dots$). Let n, k be any positive integers and put*

$$S_k = \sum_{\nu=1}^n \zeta_\nu^k.$$

Then

Received by the editors August 12, 1968.

$$\frac{nr^2}{(1-r^2)^2} - \frac{n^2}{(1-\cos^2 \epsilon)^2} \leq \sum_{k=1}^{\infty} |S_k|^2 k r^{2k} \quad (0 < r < 1).$$

PROOF. Let $z = re^{i\theta}$ ($0 < r < 1$),

$$P_n(z) = \sum_{\nu=1}^n \sum_{k=1}^{\infty} k^{1/2} (\zeta_{\nu, z})^k = \sum_{k=1}^{\infty} k^{1/2} S_k z^k.$$

We see that

$$\begin{aligned} \int_0^{2\pi} |P_n(z)|^2 d\theta &= \int_0^{2\pi} \sum_{\nu, \nu'=1}^n \sum_{k, k'=1}^{\infty} (kk')^{1/2} \zeta_{\nu} \bar{\zeta}_{\nu'} r^{k+k'} e^{i\theta(k-k')} d\theta \\ &= 2\pi \sum_{\nu, \nu'=1}^n \sum_{k=1}^{\infty} k (\zeta_{\nu} \bar{\zeta}_{\nu'} r^2)^k \\ &= 2\pi \sum_{\nu, \nu'=1}^n \frac{\zeta_{\nu} \bar{\zeta}_{\nu'} r^2}{(1 - \zeta_{\nu} \bar{\zeta}_{\nu'} r^2)^2} \\ &= \frac{2\pi nr^2}{(1-r^2)^2} + 2\pi \sum_{\nu \neq \nu'; 1 \leq \nu, \nu' \leq n} \frac{\zeta_{\nu} \bar{\zeta}_{\nu'} r^2}{(1 - \zeta_{\nu} \bar{\zeta}_{\nu'} r^2)^2}, \end{aligned} \quad (1)$$

and for $\phi_{\nu} \neq \phi_{\nu'}$,

$$\begin{aligned} \left| \frac{\zeta_{\nu} \bar{\zeta}_{\nu'} r^2}{(1 - \zeta_{\nu} \bar{\zeta}_{\nu'} r^2)^2} \right| &\leq \frac{1}{\inf_{0 < r < 1} |1 - \zeta_{\nu} \bar{\zeta}_{\nu'} r^2|^2} \\ &= \frac{1}{(1 - \cos^2 \epsilon)^2}. \end{aligned} \quad (2)$$

On the other hand, we have

$$\int_0^{2\pi} |P_n(z)|^2 d\theta = 2\pi \sum_{k=1}^{\infty} |S_k|^2 k r^{2k}. \quad (3)$$

It follows from (1), (2), (3) that

$$\sum_{k=1}^{\infty} |S_k|^2 k r^{2k} \geq \frac{nr^2}{(1-r^2)^2} - \frac{n^2}{(1-\cos^2 \epsilon)^2} \quad (0 < r < 1).$$

The proof of Lemma 1 is completed.

LEMMA 2. Let $\{z_{\nu}\}$ be any sequence of complex numbers. If $\limsup_{k \rightarrow \infty} B_k/k^{1-\delta} < C$, for some absolute constants $\delta, C > 0$, then there is an integer $k_1 > 0$ which depends on $\{z_{\nu}\}$, C and δ only, such that given integers n and $\mu(n)$, with $\mu(n) \geq k_1$, we can find n members of $\{z_{\nu}\} : \zeta_1, \zeta_2, \dots, \zeta_n$ satisfying

$$|S_k| = \left| \sum_{\nu=1}^n \zeta_\nu^k \right| < 2Ck^{1-\delta} \quad (k_1 \leq k \leq \mu(n)).$$

PROOF. From the given hypothesis there is an integer $k_1 > 0$ such that

$$B_k < Ck^{1-\delta} \quad (k_1 \leq k).$$

Since $B_k = \limsup_{m \rightarrow \infty} \left| \sum_{\nu=1}^m z_\nu^k \right|$, we see that for each $k \geq k_1$, there exists an integer $N(k) > 0$ such that

$$\left| \sum_{\nu=1}^m z_\nu^k \right| < Ck^{1-\delta} \quad (m > N(k)).$$

For given n choose an integer $N > 0$ so that

$$N \geq \max_{k_1 \leq k \leq \mu(n)} \{N(k)\}.$$

Putting $\zeta_\nu = z_{N+\nu}$ ($\nu = 1, 2, \dots, n$), we have

$$\begin{aligned} |S_k| &= |\zeta_1^k + \zeta_2^k + \dots + \zeta_n^k| \\ &= |z_{N+1}^k + z_{N+2}^k + \dots + z_{N+n}^k| \\ &= \left| \sum_{\nu=1}^{N+n} z_\nu^k - \sum_{\nu=1}^N z_\nu^k \right| \\ &\leq \left| \sum_{\nu=1}^{N+n} z_\nu^k \right| + \left| \sum_{\nu=1}^N z_\nu^k \right| \\ &< 2Ck^{1-\delta} \quad (k_1 \leq k \leq \mu(n)). \end{aligned}$$

This completes the proof of Lemma 2.

We come now to prove our Theorem. For the given sequence $\{z_\nu\}$, suppose $\limsup_{k \rightarrow \infty} B_k/k^{1-\delta} < C$. By virtue of Lemma 2, there exists a positive integer k_1 such that for any given n and $\mu(n)$, with $\mu(n) \geq k_1$, there are $\zeta_1, \zeta_2, \dots, \zeta_n \in \{z_\nu\}$ with

$$(4) \quad |S_k| < 2Ck^{1-\delta} \quad (k_1 \leq k \leq \mu(n)).$$

From Lemma 1 we have

$$\begin{aligned} (5) \quad \frac{-n^2}{(1 - \cos^2 \epsilon)^2} + \frac{n r^2}{(1 - r^2)^2} &\leq \sum_{k=1}^{\infty} |S_k|^2 k r^{2k} \\ &= \left(\sum_{k=1}^{k_1} + \sum_{k > k_1}^{\mu(n)} + \sum_{k > \mu(n)}^{\infty} \right) |S_k|^2 k r^{2k} \\ &= T_1 + T_2 + T_3, \text{ say.} \end{aligned}$$

Since $|S_k| \leq n$, we have

$$\begin{aligned} T_1 &= \sum_{k=1}^{k_1} |S_k|^2 k r^{2k} \leq n^2 \sum_{k=1}^{k_1} k < n^2 k_1, \\ (6) \quad T_3 &= \sum_{k > \mu(n)}^{\infty} |S_k|^2 k r^{2k} \leq n^2 \sum_{k > \mu(n)}^{\infty} k r^{2k} \\ &\leq n^2 \mu(n) \frac{r^{2\mu(n)}}{(1-r^2)^2}. \end{aligned}$$

Let A_k^α be the Cesàro numbers of order α ($0 < \alpha$), we have

$$\begin{aligned} \sum_{k=0}^{\infty} A_k^\alpha r^{2k} &= \frac{1}{(1-r^2)^{1+\alpha}} \quad (0 < r < 1), \\ A_k^\alpha &= \frac{k^\alpha}{\Gamma(1+\alpha)} \{1 + o(1)\} \quad \text{as } k \rightarrow \infty. \end{aligned}$$

See [4, Vol. I, pp. 76-77]. Assume that k_1 in (4) is large such that for every $k \geq k_1$ we have $A_k^{3-\delta} \Gamma(4) \geq k^{3-\delta}$. From (4) it follows that

$$\begin{aligned} (7) \quad T_2 &= \sum_{k > k_1}^{\mu(n)} |S_k|^2 k r^{2k} \leq 4C^2 \sum_{k > k_1}^{\mu(n)} k^{3-\delta} r^{2k} \\ &\leq 4C^2 \Gamma(4) \sum_{k=0}^{\infty} A_k^{3-\delta} r^{2k} = 24C^2 \frac{1}{(1-r^2)^{4-\delta}}. \end{aligned}$$

From (5), (6), (7) we have

$$(8) \quad \frac{-n^2}{(1 - \cos^2 \epsilon)^2} + \frac{n r^2}{(1 - r^2)^2} \leq n^2 k_1^2 + \frac{24C^2}{(1 - r^2)^{4-\delta}} + \frac{n^2 \mu(n) r^{2\mu(n)}}{(1 - r^2)^2}.$$

Put $1/(1-r^2) = n^{1/(2-\delta)}$ and $\mu(n) = n^2$. Then we have

$$\frac{n}{(1-r^2)^2} = \frac{1}{(1-r^2)^{4-\delta}} = n^{2+(\delta/(2-\delta))},$$

and

$$\frac{n^2 \mu(n) r^{2\mu(n)}}{(1-r^2)^2} < n^6 (1 - n^{-1/(2-\delta)})^{n^2} < n^6 e^{-n} = o(1),$$

as $n \rightarrow \infty$. Thus, taking $n \rightarrow \infty$, from (8) we have

$$1 \leq 24C^2.$$

This implies that

$$\limsup_{k \rightarrow \infty} B_k/k^{1-\delta} \geq 1/\sqrt{24} > 1/5 \quad (\delta > 0).$$

The proof is completed.

My thanks are due to the referee for some useful remarks and improvements in this paper.

REFERENCES

1. J. Clunie, *On a problem of Erdős*, J. London Math. Soc. **42** (1967), 133–136.
2. P. Erdős, *Problems and results on Diophantine approximations*, Compositio Math. **16** (1964), 52–65.
3. ———, *Some remarks on number theory*, Israel J. Math. **3** (1965), 6–12.
4. A. Zygmund, *Trigonometric series*, Cambridge Univ. Press, New York, 1959.

UNIVERSITY OF HONG KONG