GROUPS WHICH ARE COGROUPS

M. L. CURTIS AND J. DUGUNDJI

It is the purpose of this note to give an elementary proof that S^0 , S^1 , S^3 are the only spheres which can be made into topological groups. This was proved by Samelson in [3]. He showed that a compact Lie group which is a homotopy sphere must have rank 1, and then that a compact Lie group of rank 1 has dimension 1 or 3. We get the first part easily by showing that a compact Lie group which is an H -cogroup (e.g., a suspension) must have rank 1. The second part closely follows Samelson's proof. For basic facts about H -groups and H -cogroups the reader is referred to §§5 and 6 of Chapter I of Spanier [5].

If G is an H -group which is also an H -cogroup, then the set $[G, G]_0$ of pointed homotopy classes is an abelian group with its operation given equally by the H -cogroup structure on the domain G or the H -group structure on the range G [5, page 44]. From this the following lemma is immediate.

LEMMA 1. *If G is an H -group which is also an H -cogroup, $i: G \rightarrow G$ is the identity map and $\psi: G \rightarrow G$ is the squaring map $x \rightarrow x^2$, then*

$$[\psi] = 2[i] \in [G, G]_0.$$

Next we need a special case of a result due to Hopf [2].

LEMMA 2. *If G is a compact connected Lie group of rank r and ψ is the squaring map, then degree $\psi = 2^r$.*

PROOF. We will count the algebraic number of times a point $y \in G$ is covered. We may assume y is regular (i.e., lies in a unique maximal torus) because the set of singular (= not regular) points forms a set of dimension less than the dimension of G (indeed, the dimension is at most $\dim G - 3$; see [4, p. 19]).

Let T be the unique maximal torus containing y . Since $\dim T = r$ we easily find that y has exactly 2^r square roots in T . If z is any other square root of y , then z lies in some other maximal torus $T' \neq T$ (since maximal tori cover G). But then $y = z^2 \in T'$, contradicting the regularity of y . Thus there are exactly 2^r points $x \in G$ with $\psi(x) = y$.

We assert that all such x cover y with the same orientation. Given x with $x^2 = y$, let U be a small open neighborhood of x containing no other roots of y and such that U is star-shaped relative to x . Then

Received by the editors September 13, 1968.

$$\psi: U \rightarrow G$$

is a diffeomorphism (if U is small enough). Furthermore, if $T_x: U \rightarrow G$ is translation by x it is easy to get an *isotopy* between ψ and T_x on U . Since T_x preserves orientation, so does ψ . Hence a neighborhood of y is covered 2^r times with the same orientation. Hence the degree of ψ is 2^r .

From Lemmas 1 and 2 we get

THEOREM 1. *If a compact Lie group G is also an H -cogroup, then $\text{rank } G = 1$.*

COROLLARY. *If $n \geq 1$ and S^n can be made into a topological group, then S^n is a Lie group of rank 1.*

PROOF. A group manifold is a Lie group, and S^n is compact, so Theorem 1 applies.

THEOREM 2 (FOLLOWING SAMELSON). *If G is a compact Lie group of rank 1, and $\dim G = n > 1$, then $n = 3$.*

PROOF. Since G is compact we can get an inner product on its Lie algebra $\mathfrak{g}$ such that the adjoint representation

$$\text{Ad}: G \rightarrow GL(\mathfrak{g})$$

is such that each $\text{Ad}(g)$ is orthogonal; i.e., $\text{Ad}(G) \subset \mathcal{O}(\mathfrak{g})$ the orthogonal group on $\mathfrak{g}$.

Let $T = S^1$ be a maximal torus of G and let X be a unit vector in $\mathfrak{g}$ tangent to T . Define

$$\phi': G \rightarrow S^{n-1} \subset \mathfrak{g}$$

by $\phi'(g) = \text{Ad } gX$.

LEMMA. *ϕ' induces a homeomorphism $\phi: G/T \rightarrow S^{n-1}$.*

PROOF OF LEMMA. First off, ϕ' does induce ϕ since if $t \in T$

$$\phi'(gt) = \text{Ad}(gt)X = \text{Ad}(g)\text{Ad}(t)X = \text{Ad}(g)X = \phi'(g).$$

Second, ϕ is injective for if

$$\text{Ad } gX = \text{Ad } hX,$$

then $\text{Ad}(gh^{-1})X = X$. Thus $\text{Ad}(gh^{-1})$ leaves the one-dimensional Lie algebra $\mathcal{L}(T)$ of T elementwise fixed. Since T is maximal this implies $gh^{-1} \in T$.

Finally, G/T is a compact closed $n-1$ manifold and it follows that ϕ is surjective, so the Lemma is proved.

Since ϕ maps G/T onto S^{n-1} it follows that there exists $g_0 \in G$ such that $\text{Ad } g_0 X = -X$. This allows us to get a map $f: S^2 \rightarrow S^{n-1}$ as follows. Let $\omega: [0, 1] \rightarrow G$ be a path from the identity e to g_0 . Then since $T = S^1$ we have a map

$$F: S^1 \times I \rightarrow G$$

given by $F(t, \tau) = \omega(\tau)t\omega(\tau)^{-1}$. Then $F(S^1 \times 0) \subset S^1$ and $F(S^1 \times 1) \subset S^1$ so F induces a map $f: S^2 \rightarrow G/S^1 = S^{n-1}$. We assert that f is essential.

If f is nullhomotopic, we can cover a nullhomotopy in G to get a homotopy

$$\Phi: T \times I \rightarrow T$$

with $\Phi_0 = F(-, 0)$ the identity and $\Phi_1 = F(1, 1)$ sends t to t^{-1} . No such homotopy exists, so f is essential and we conclude that $S^{n-1} = S^2$ and $n = 3$.

THEOREM 3. *The n -sphere S^n ($n > 0$) can be made into a topological group if and only if $n = 1$ or $n = 3$.*

PROOF. Use Theorem 2 and the corollary to Theorem 1.

REMARK. $\text{SO}(3)$ is a compact Lie group of rank 1 and dimension 3, so that it could be a cogroup as far as the results above are concerned. But Hilton [1] has shown that a comultiplication on X implies X has Lusternik-Schnirelmann category ≤ 2 , whereas $\text{SO}(3) = P^3$ has category 4.

REFERENCES

1. P. J. Hilton, *On a generalization of nilpotency to semi-simplicial complexes*, Proc. London Math. Soc. **10** (1960), 604-622.
2. H. Hopf, *Über der Rang geschlossener Liescher Gruppen*, Comment. Math. Helv. **13** (1940-1941), 119-143.
3. H. Samelson, *Über die Sphären, die als Gruppenräume, auftreten*, Comment. Math. Helv. **13** (1940), 144-155.
4. ———, *Topology of Lie groups*, Bull. Amer. Math. Soc. **58** (1952), 2-37.
5. E. Spanier, *Algebraic topology*, McGraw-Hill, New York, 1966.

RICE UNIVERSITY AND
THE UNIVERSITY OF TEXAS