ISOMETRIES OF THE TRACE CLASS

BERNARD RUSSO¹

Let $\mathfrak{J}$ denote the Banach space of trace class operators on a complex Hilbert space H , in the norm $\|T\|_1 = \text{Tr}(|T|)$. The space $\mathfrak{J}$ is a two-sided ideal in the algebra $\mathfrak{L}$ of all bounded operators on H . See [4].

THEOREM. *If Φ is a linear isometry of the Banach space $\mathfrak{J}$ onto itself, then there exists a *-automorphism or a *-antiautomorphism α of $\mathfrak{L}$ and a unitary operator U in $\mathfrak{L}$ such that $\Phi(T) = \alpha(TU)$, (T in $\mathfrak{J}$).*

REMARK 1. The theorem provides a partial answer to [3, Remark 1, p. 231].

PROOF. The adjoint Φ' is a linear isometry of $\mathfrak{L}$ onto $\mathfrak{L}$ so by results of Kadison [2, Theorem 7, Corollary 11] has the form $\Phi'(A) = U\alpha(A)$ where α and U are as described in the statement of the theorem. It is elementary that $\Phi(T) = \Psi(TU)$ where $\Psi' = \alpha$. The proof will be complete if it is shown that α is the adjoint of α^{-1} (restricted to $\mathfrak{J}$). By the folk result [1, pp. 256, 9] it is sufficient to check this in the following two cases:

(i) $\alpha(A) = VA V^{-1}$ with V a fixed unitary operator; then $\langle T, \alpha(A) \rangle = \langle T, VA V^{-1} \rangle = \langle V^{-1}TV, A \rangle = \langle \alpha^{-1}(T), A \rangle$,

(ii) after the choice of an orthonormal basis, $\alpha(A)$ is the transposed matrix of A ; then $\langle T, \alpha(A) \rangle = \text{Tr}(T\alpha(A)) = \text{Tr}(\alpha(T)A) = \langle \alpha^{-1}(T), A \rangle$.

REMARK 2. A previous version of the above proof exploited a knowledge of the extreme points of the unit sphere of $\mathfrak{J}$. These were determined to be the partial isometries with initial (hence final) domain one-dimensional.

REFERENCES

1. J. Dixmier, *Les algèbres d'opérateurs dans l'espace hilbertien*, Gauthier-Villars, Paris, 1957.
2. R. V. Kadison, *Isometries of operator algebras*, Ann. of Math. (2) **54** (1951), 325-338.
3. B. Russo, *Isometries of L^p spaces associated with finite von Neumann algebras*, Bull. Amer. Math. Soc. **74** (1968), 228-232.
4. R. Schatten, *Norm ideals of completely continuous operators*, Springer-Verlag, Berlin, 1960.

UNIVERSITY OF CALIFORNIA, IRVINE

Received by the editors March 7, 1969.

¹ This research was supported by the National Science Foundation Grant GP-8291.