ON WU'S FORM OF THE FIRST MAIN THEOREM OF VALUE DISTRIBUTION

JOHN J. HIRSCHFELDER

In the First Main Theorem of value distribution for holomorphic maps into a compact Kähler manifold A , the central object is a differential form λ_a on A with a singularity at $a \in A$. The most satisfactory λ_a was given by Wu in [5]. His construction, using the Green function, has the advantage that λ_a is automatically invariant under the group of isometries of A ; more precisely, if $i: A \rightarrow A$ is an isometry with $i(a) = b$, then $i^* \lambda_b = \lambda_a$. This property is difficult to obtain with any other construction.

In this note we apply the technical machinery developed by Stoll and the author [1], [2], [3], [4] to Wu's λ_a to show that it is suitable for value distribution theory in arbitrary fiber dimension. We also show that the terms of the First Main Theorem are continuous functions.

v_f will denote the multiplicity of the holomorphic map f ; see [2]. O_k will denote any differential form of degree k whose coefficients are locally bounded; the subscript will usually be omitted. Of course it will never mean the same thing twice. On $\mathbb{C}^n$ set $v_1 = (i/2) \sum_{\mu=1}^n dz_\mu \wedge d\bar{z}_\mu$ and $v_1^k = v^k/k!$.

If M is a complex manifold and H an open subset of M , then $S = \bar{H} - H$ is a *boundary manifold* of H , provided that for every point of S there is a C^∞ coordinate system $(x_1, \dots, x_{2n})$ valid in a neighborhood U of the point, such that in U , S is described by $x_1 = 0$ and H by $x_1 < 0$. S will be considered as oriented to the exterior of H .

1. We collect the technical results which we will need.

LEMMA 1. Let $k \geq 0$, $0 \leq s \leq n$ and $0 \leq t \leq n$ be integers such that $s + t < 2n$. Let $\phi(w, z)$ be a differential form of bidegree (s, t) in $w \in \mathbb{C}^n$, dependent on a parameter $z \in \mathbb{C}^n$, whose coefficients are measurable and locally bounded. Let M be a complex manifold, and let $f: M \rightarrow \mathbb{C}^n$ be an open holomorphic map. Let K be a compact subset of M . Let $\chi(w, z)$ be a differential form in $w \in M$ dependent also on a parameter $z \in \mathbb{C}^n$, whose coefficients are measurable and locally bounded; then

$$F(z) = \int_{w \in K} [\log(1/|f(w) - z|)]^k |f(w) - z|^{-(s+t)} \\ \cdot f^* \phi(w, z) \wedge \chi(w, z)$$

Received by the editors April 29, 1969.

exists for each $z \in \mathbf{C}^n$, and is a measurable and locally bounded function on $\mathbf{C}^n$. If moreover ϕ is continuous off the diagonal and χ is continuous off $\{(w, z) | f(w) = z\}$, then F is continuous.

This is a restatement of Lemma 2.6 of [1].

Choose a function h of class C^∞ on $\mathbf{R}$ such that $0 \leq h(x) \leq 1$ for all x , $h(x) = 0$ if $x \leq 0$, and $h(x) = 1$ if $x \geq 1$. A constant $B > 0$ exists such that $2|h'(x)| \leq B$ for all x .

For $\rho > 0$ define

$$g_\rho(X) = h((2X - \rho)/\rho).$$

Then

- (a) $0 \leq g_\rho(X) \leq 1$,
- (b) $X |g'_\rho(X)| \leq B$,
- (c) $X \leq \rho/2 \Rightarrow g_\rho(X) = 0$,
- (d) $X \geq \rho \Rightarrow g_\rho(X) = 1$.

Now let $f: M \rightarrow \mathbf{C}^n$ be a holomorphic map. Define γ_ρ on M by $\gamma_\rho(w) = g_\rho(|f(w)|)$. Then $d\gamma_\rho = g'_\rho(|f|)((df|f) + (f|df))/2|f|$. Set $h_\rho = |f|g'_\rho(|f|)$ and $\eta = ((df|f) + (f|df))/2|f|$. Then the functions h_ρ are uniformly bounded, η is a locally bounded form, and

$$d\gamma_\rho = h_\rho \eta / |f|.$$

Define $L_\rho = \{w \in K | \rho/2 \leq \gamma_\rho(w) \leq \rho\}$.

LEMMA 2. *Hypotheses as in Lemma 1, let also a measurable function h_ρ be given for each $\rho > 0$, and suppose that these functions are uniformly bounded on K . Then*

$$\lim_{\rho \rightarrow 0} \int_{L_\rho} h_\rho [\log(1/|f|)]^k |f|^{-(s+t)} f^* \phi \wedge \chi = 0.$$

See Stoll [4, Lemma 1.7].

LEMMA 3. *Hypotheses as above, let ϕ be a locally bounded form of degree $2n-1$ on $\mathbf{C}^n$. Then*

$$\lim_{\rho \rightarrow 0} \int_K d\gamma_\rho \wedge |f|^{2-2n} [\log(1/|f|)]^k f^* \phi \wedge \chi = 0.$$

PROOF. The expression under the integral sign is

$$-(h_\rho/|f|) |f|^{2-2n} [\log(1/|f|)]^k f^* \phi \wedge (\eta \wedge \chi).$$

Now decompose ϕ into forms of bidegrees $(n, n-1)$ and $(n-1, n)$ and apply Lemma 2 twice. q.e.d.

LEMMA 4. *Hypotheses as above, and suppose also that H is an open subset of M such that $\overline{H} = K$ and that such that $S = \overline{H} - H$ is either empty or a smooth boundary manifold of H . Let T be the support of χ on S and suppose that $T \cap f^{-1}(0)$ is of measure zero on $f^{-1}(0)$. Then*

$$\begin{aligned} \lim_{\rho \rightarrow 0} \int_H d\gamma_\rho \wedge d^c \log(1/|f|) \wedge |f|^{2-2n} f^* v_n \wedge \chi \\ = - [2\pi^n / (n-1)!] \int_{H \cap f^{-1}(0)} v_f \chi. \end{aligned}$$

See Stoll [4, Lemma 1.8].

2. Let A be a compact Kähler manifold of dimension n . Denote its fundamental form by ω_1 , and set $\omega_k = \omega_1^k / k!$, $\omega = \omega_n$. We assume that the metric is normalized so that $\int_A \omega = 1$. Denote by $r(z, w)$ the distance between z and w .

Denote by $g(z, w)$ the Green function on $A \times A$. Then

$$\begin{aligned} (1) \quad g(z, w) &= \frac{1}{4}(n-2)! \pi^{-n} r(z, w)^{2-2n} + r(z, w)^{3-2n} O(z, w), \quad n > 1, \\ g(z, w) &= (1/2\pi) \log(1/r(z, w)) + O(z, w), \quad n = 1. \end{aligned}$$

There exists a constant C such that $g(z, w) + C$ is everywhere positive.

Define

$$(2) \quad \lambda_z(w) = (g(z, w) + C) \omega_{n-1}(w).$$

Then $dd^c \lambda_z = \omega$. (See Wu [5, I and II].)

Let $a \in A$, and let $(z_1, \dots, z_n)$ be a holomorphic coordinate system in a neighborhood U centered at a , such that in the underlying real coordinate system the metric tensor at a is represented by the identity matrix. Set

$$(3) \quad B(z, w) = |z - w| / r(z, w)$$

where $|z - w|$ is computed in the $(z_1, \dots, z_n)$ coordinate system. B is continuous and positive on $U \times U$ and $B(a, a) = 1$.

From (1), (2), and (3) we see

$$\begin{aligned} (4) \quad \lambda_z(w) &= |z - w|^{2-2n} O(z, w) v_{n-1}, \quad n > 1, \\ \lambda_z(w) &= (1/2\pi) \log(1/|z - w|) + O(z, w), \quad n = 1. \end{aligned}$$

From (1), (see Wu [5, II, §8] for justification)

$$\begin{aligned}
 d_w^c \lambda_z(w) &= [-(n-1)!/2\pi^n] r(z, w)^{1-2n} d_w^c r(z, w) \\
 &\quad + O_1(z, w) r(z, w)^{2-2n} \wedge \omega_{n-1}(w) \quad (\text{if } n > 1) \\
 (5) \qquad &= [[(n-1)!/2\pi^n] r(z, w)^{2-2n} d_w^c \log(1/r(z, w)) \\
 &\quad + O_1(z, w) r(z, w)^{2-2n}] \wedge \omega_{n-1}(w) \quad (\text{any } n).
 \end{aligned}$$

Now $\log(1/r(z, w)) = \log B(z, w) + \log(1/|z-w|)$ and $\log B(z, w)$ is locally bounded; hence

$$\begin{aligned}
 d_w^c \lambda_z(w) &= [[(n-1)!/2\pi^n] B(z, w)^{2n-2} r(z, w)^{2-2n} d_w^c \log(1/|z-w|) \\
 (6) \qquad &\quad + O_1(z, w) |z-w|^{2-2n}] \wedge \omega_{n-1}(w).
 \end{aligned}$$

Since $d^c \log(1/|z-w|) = -d^c |z-w|/|z-w|$, we have

$$(7) \qquad d^c \lambda_z(w) = O_{2n-1} |z-w|^{1-2n}.$$

Finally, look at $z=0$, i.e., at a :

$$\begin{aligned}
 d^c \log(1/|w|) \wedge (\omega_{n-1}(w) - v_{n-1}(w)) &= -d^c |w| \wedge (\omega_{n-1}(w) \\
 &\quad - v_{n-1}(w))/|w|.
 \end{aligned}$$

Since $\omega_{n-1}(a) = v_{n-1}(a)$, this is bounded. Thus

$$\begin{aligned}
 d^c \lambda_a(w) &= [(n-1)!/2\pi^n] B(a, w)^{2n-2} |w|^{2-2n} d^c \log(1/|w|) \wedge v_{n-1} \\
 (8) \qquad &\quad + O_{2n-1}(w) |w|^{2-2n}.
 \end{aligned}$$

LEMMA 5. Let $f: M \rightarrow A$ be open and holomorphic, K a compact subset of M , and χ continuous on K ; then

$$F(z) = \int_K f^* \lambda_z \wedge \chi$$

exists for each $z \in A$ and is continuous.

PROOF. The question is local on both K and A , so we may assume $f(K) \subset U$ and adopt the above notation. Apply (4) and Lemma 1. q.e.d.

LEMMA 6. In the same situation,

$$G(z) = \int_K f^* d^c \lambda_z \wedge \chi$$

exists and is continuous on A .

PROOF. Use (7) and Lemma 1.

3. THE RESIDUE THEOREM. *Let M be a complex manifold of dimension $m = n + q$. Let A be a compact Kähler manifold, and let $f: M \rightarrow A$ be an open holomorphic map. Let H be an open subset of M . Let $S = \overline{H} - H$ be empty or a boundary manifold of H , and suppose that $\overline{H}$ is compact. Let ψ be a continuous function on $\overline{H}$ satisfying a Lipschitz condition. (Then $d\psi$ exists almost everywhere.) Let χ be a differential form of bi-degree (q, q) and class C^1 on M .*

Then $d^c f^ \lambda_a \wedge d\psi \wedge \chi$ and $d^c f^* \lambda_a \wedge \psi \wedge d\chi$ are integrable over H for each $a \in A$.*

Also let T be the support of $\psi\chi$ on S , and assume that $T \cap f^{-1}(a)$ is a set of measure zero on $f^{-1}(a)$. Suppose that $d^c f^ \lambda_a \wedge \psi\chi$ is integrable on S . Then*

$$\begin{aligned} \int_S d^c f^* \lambda_a \wedge \psi\chi &= - \int_H d^c f^* \lambda_a \wedge d\psi \wedge \chi - \int_H d^c f^* \lambda_a \wedge \psi \wedge d\chi \\ &\quad + \int_H f^* \omega \wedge \psi\chi - \int_{H \cap f^{-1}(a)} \nu_f \psi\chi. \end{aligned}$$

PROOF. By Stokes' Theorem,

$$\begin{aligned} \int_S \gamma_\rho d^c f^* \lambda_a \wedge \psi\chi &= - \int_H \gamma_\rho d^c f^* \lambda_a \wedge d\psi \wedge \chi - \int_H \gamma_\rho d^c f^* \lambda_a \wedge \psi \wedge d\chi \\ &\quad + \int_H \gamma_\rho f^* \omega \wedge \psi\chi + \int_H d\gamma_\rho \wedge d^c f^* \lambda_a \wedge \psi\chi. \end{aligned}$$

We let $\rho \rightarrow 0$; from Lemma 6 we see that we need only show that the last integral has limit

$$- \int_{H \cap f^{-1}(a)} \nu_f \psi\chi.$$

By (8), it is

$$\begin{aligned} \int_H d\gamma_\rho \wedge [(n-1)!/2\pi^n] B(a, f)^{2n-2} |f|^{2-2n} \wedge d^c \log(1/|f|) \wedge \nu_{n-1} \wedge \psi\chi \\ + \int_H d\gamma_\rho f^* O_{2n-1} |f|^{2-2n} \wedge \psi\chi. \end{aligned}$$

By Lemma 3, the second term has limit zero. By Lemma 4, the first term has limit

$$- \int_{H \cap f^{-1}(a)} \nu_f \psi\chi B(a, a)^{2n-2}$$

but $B(a, a) = 1$. q.e.d.

SINGULAR STOKES THEOREM. *Let M, A, f, H, S , and χ be as above. Let ψ be a function of class C^2 on $\overline{H}$. Then $d^c f^* \lambda_a \wedge d\psi \wedge \chi, f^* \lambda_a \wedge dd^c \psi \wedge \chi$, and $f^* \lambda_a \wedge d^c \psi \wedge d\chi$ are integrable over H .*

Assume further that $f^ \lambda_a \wedge d^c \psi \wedge \chi$ is integrable over S . Then*

$$\begin{aligned} \int_S f^* \lambda_a \wedge d^c \psi \wedge \chi &= - \int_H d^c f^* \lambda_a \wedge d\psi \wedge \chi + \int_H f^* \lambda_a \wedge dd^c \psi \wedge \chi \\ &\quad - \int_H f^* \lambda_a \wedge d^c \psi \wedge d\chi. \end{aligned}$$

PROOF. Similar to that of the Residue Theorem. Apply Lemmas 5 and 6; then consider

$$\int_H d\gamma_\rho \wedge f^* \lambda_a \wedge d^c \psi \wedge \chi.$$

By (4) it is

$$\int_H d\gamma_\rho \wedge O_0 |f|^{2-2n} f^* v_{n-1} \wedge d^c \psi \wedge \chi$$

if $n > 1$ and

$$\int_H d\gamma_\rho \wedge [(1/2\pi) \log(1/|f|) + O_0] \wedge d^c \psi \wedge \chi$$

if $n = 1$. Now apply Lemma 3. q.e.d.

The so-called Unintegrated Form of the First Main Theorem is obtained by setting $\psi = 1$ and choosing χ so that $d\chi = 0$ in the Residue Theorem. We can now state the:

INTEGRATED FIRST MAIN THEOREM. *Let M, A, f and χ be as before, and assume $d\chi = 0$. Let G be an open subset of M with compact closure $\overline{G}$. Assume that $\Gamma = \overline{G} - G$ is a boundary manifold of G . Let g be open with $\overline{g} \subset G$, and assume that $\gamma = \overline{g} - g$ is a boundary manifold of g . Let ψ be a continuous, real-valued function on M , of class C^2 on $\overline{G} - g$, such that $\psi|_g = R$ a constant, $\psi|_{M-G} = 0$, and $0 \leq \psi(z) \leq R$ for all $z \in M$. (Such a ψ satisfies a Lipschitz condition on M .) Define*

$$m^{\#}(a) = - \int_{\Gamma} f^* \lambda_a \wedge d^c \psi \wedge \chi,$$

$$m^b(a) = - \int_{\gamma} f^* \lambda_a \wedge d^c \psi \wedge \chi,$$

$$N(a) = \int_{f^{-1}(a)} \nu_f \psi \chi,$$

$$T = \int_G f^* \omega \wedge \psi \chi,$$

$$\Delta(a) = \int_G f^* \lambda_a \wedge d^c d\psi \wedge \chi.$$

Then these are continuous functions on A , and

$$N + m^\# - m^b = T + \Delta.$$

Note that N , m , m^b , and T are always nonnegative; but in general we have information on Δ .

This theorem is obtained by applying the Residue Theorem to G , the Singular Stokes Theorem to $G - \bar{g}$, and eliminating the common term; see [1] for details. The continuity of Δ is a consequence of Lemma 5, and the continuity of N is a theorem of Stoll [3, Theorem 3.8]. Thus $m^\# - m^b$ is continuous. Now let h be a C^∞ function with compact support in G and which is 1 in a neighborhood of $\bar{g}$, and replace ψ by $h\psi$. We get another FMT with different (but still continuous) N , T , and Δ . The new $m^\#$ is zero but the new m^b is the same as the old. Hence the old m^b and $m^\#$ are continuous.

REFERENCES

1. J. Hirschfelder, *The first main theorem of value distribution in several variables*, Invent. Math. **8** (1969), 1–33.
2. W. Stoll, *The multiplicity of a holomorphic map*, Invent. Math. **2** (1966), 15–58.
3. ———, *The continuity of the fiber integral*, Math. Z. **95** (1967), 87–138.
4. ———, *A general first main theorem of value distribution theory*, Acta Math. **118** (1967), 111–191.
5. H. Wu, *Remarks on the first main theorem in equidistribution theory*. I, II, J. Differential Geometry **2** (1968), 197–202, 369–384.

UNIVERSITY OF WASHINGTON