GENERATING FUNCTIONS FOR JACOBI AND LAGUERRE POLYNOMIALS

H. M. SRIVASTAVA¹

Let v be a function of t defined by

$$(1) \quad v = t(1 + v)^{b+1}, \quad v(0) = 0.$$

Then it follows from Lagrange's expansion formula [6, Vol. I, p. 126, Ex. 212] that

$$(2) \quad \frac{(1 + v)^{a+1}}{1 - bv} = \sum_{n=0}^{\infty} \binom{a + (b+1)n}{n} t^n.$$

Making use of the formula (2), Carlitz [2] has proved that the Laguerre polynomial $L_n^{(a+bn)}(x)$, where

$$(3) \quad L_n^{(\alpha)}(x) = \sum_{k=0}^n (-1)^k \binom{\alpha + n}{n - k} \frac{x^k}{k!},$$

satisfies a generating relation in the form

$$(4) \quad \sum_{n=0}^{\infty} L_n^{(a+bn)}(x) t^n = \frac{(1 + v)^{a+1}}{1 - bv} \exp(-xv),$$

where v is given by (1) and a, b are arbitrary complex numbers. Note that the special case of (4) when b is an arbitrary integer was proved earlier by Brown [1].

In terms of the generalized hypergeometric function

$$(5) \quad {}_pF_q \left[\begin{matrix} \alpha_1, \dots, \alpha_p; \\ \beta_1, \dots, \beta_q; \end{matrix} x \right] = \sum_{n=0}^{\infty} \frac{\prod_{j=1}^p (\alpha_j)_n}{\prod_{j=1}^q (\beta_j)_n} \frac{x^n}{n!},$$

where

$$(6) \quad (\lambda)_n = \lambda(\lambda + 1)(\lambda + 2) \cdots (\lambda + n - 1), \quad n \geq 1, \quad (\lambda)_0 = 1,$$

the generating relation (4) assumes the form

Received by the editors March 22, 1969.

¹ Present address. Department of Mathematics, University of Victoria, Victoria, British Columbia, Canada.

$$\begin{aligned}
 (7) \quad \sum_{n=0}^{\infty} \binom{a + (b+1)n}{n} {}_1F_1 \left[\begin{matrix} -n; \\ x \end{matrix} \right] t^n \\
 = \frac{(1+v)^{a+1}}{1-bv} \exp(-xv).
 \end{aligned}$$

In (7) if we replace x by xz , multiply both sides by $z^{\lambda-1}$ and take their Laplace transforms with respect to the variable z , we shall readily obtain

$$\begin{aligned}
 (8) \quad \sum_{n=0}^{\infty} \binom{a + (b+1)n}{n} {}_2F_1 \left[\begin{matrix} -n, \lambda; \\ x \end{matrix} \right] t^n \\
 = \frac{(1+v)^{a+1}}{1-bv} (1+xv)^{-\lambda},
 \end{aligned}$$

where the binomial $(1+xv)^{-\lambda}$ may be written as an ${}_1F_0$.

The form of (8) suggests the existence of the general formula

$$\begin{aligned}
 (9) \quad \sum_{n=0}^{\infty} \binom{a + (b+1)n}{n} {}_{p+1}F_{q+1} \left[\begin{matrix} -n, \alpha_1, \dots, \alpha_p; \\ x \end{matrix} \right] t^n \\
 = \frac{(1+v)^{a+1}}{1-bv} {}_pF_q \left[\begin{matrix} \alpha_1, \dots, \alpha_p; \\ \beta_1, \dots, \beta_q; \end{matrix} -xv \right],
 \end{aligned}$$

where p, q are nonnegative integers, the α 's and a, b take general values, real or complex, and

$$(10) \quad \beta_j \neq 0, -1, -2, \dots, \quad j = 1, 2, \dots, q.$$

The derivation of (9) from (7) and (8) by the principle of multi-dimensional mathematical induction would require the Laplace and inverse Laplace transform techniques illustrated, for instance, by the author [7].

For a direct proof without using (7) and (8) we notice that, in view of the definition (5),

$$\begin{aligned}
& \sum_{n=0}^{\infty} \binom{a + (b+1)n}{n}_{p+1} F_{q+1} \left[\begin{matrix} -n, \alpha_1, \dots, \alpha_p; \\ 1 + a + bn, \beta_1, \dots, \beta_q; \end{matrix} x \right] t^n \\
&= \sum_{n=0}^{\infty} t^n \sum_{k=0}^n (-1)^k \binom{a + (b+1)n}{n-k} \frac{\prod_{j=1}^p (\alpha_j)_k}{\prod_{j=1}^q (\beta_j)_k} \frac{x^k}{k!} \\
&= \sum_{k=0}^{\infty} (-1)^k \frac{\prod_{j=1}^p (\alpha_j)_k}{\prod_{j=1}^q (\beta_j)_k} \frac{x^k}{k!} \sum_{n=k}^{\infty} \binom{a + (b+1)n}{n-k} t^n \\
&= \sum_{k=0}^{\infty} (-1)^k \frac{\prod_{j=1}^p (\alpha_j)_k}{\prod_{j=1}^q (\beta_j)_k} \frac{x^k t^k}{k!} \sum_{n=0}^{\infty} \binom{a + (b+1)k + (b+1)n}{n} t^n \\
&= \sum_{k=0}^{\infty} (-1)^k \frac{\prod_{j=1}^p (\alpha_j)_k}{\prod_{j=1}^q (\beta_j)_k} \frac{x^k t^k}{k!} \frac{(1+v)^{a+(b+1)k+1}}{1-bv},
\end{aligned}$$

by (2), and the formula (9) follows immediately. We can easily attribute a direct proof to the formula (8) which obviously corresponds to the special case $p=1$, $q=0$ of (9).

A similar generalization of Carlitz's formula [2, p. 827, Equation (16)] has the form

$$\begin{aligned}
(11) \quad & \sum_{n=0}^{\infty} \binom{-a - bn}{n}_{p+1} F_{q+1} \left[\begin{matrix} -n, \alpha_1, \dots, \alpha_p; \\ 1 - a - (b+1)n, \beta_1, \dots, \beta_q; \end{matrix} x \right] t^n \\
&= \frac{A(-t, a, b)}{1 - B(-t, b)} {}_pF_q \left[\begin{matrix} \alpha_1, \dots, \alpha_p; \\ \beta_1, \dots, \beta_q; \end{matrix} -\frac{xB(-t, b)}{1 - B(-t, b)} \right],
\end{aligned}$$

where, for convenience,

$$(12) \quad B(t, b) = - \sum_{n=1}^{\infty} \binom{(b+1)n}{n-1} \frac{t^n}{n}$$

and

$$(13) \quad A(t, a, b) = \frac{[1 - B(t, b)]^{a+1}}{1 + bB(t, b)}.$$

Indeed the formula (11) is obtainable from (9) by replacing a by $-a$ and b by $-(b+1)$.

It may be of interest to remark that for $b=0$ and $b=-1$ the formula (9) yields Chaundy's results (25) and (27) respectively (see [4, p. 62]). For $b=-\frac{1}{2}$, (9) reduces to the generating relation (7), p. 264 of Brown's recent paper.²

For the Jacobi polynomial defined by

$$(14) \quad P_n^{(\alpha, \beta)}(x) = \sum_{k=0}^n \binom{\alpha+n}{k} \binom{\beta+n}{n-k} \left(\frac{x-1}{2}\right)^{n-k} \left(\frac{x+1}{2}\right)^k,$$

it is easy to show from the identity (4.22.1) of [8, p. 63] that

$$(15) \quad P_n^{(\alpha-n, \beta-n)}(x) = \binom{n-\alpha-\beta-1}{n} \binom{1-x}{2}^n {}_2F_1 \left[\begin{matrix} -n, -\alpha; \\ -\alpha-\beta; \end{matrix} \middle| \frac{2}{1-x} \right],$$

and therefore (8) gives us the elegant generating function

$$(16) \quad \sum_{n=0}^{\infty} P_n^{(\alpha-n, \beta-(b+1)n)}(x) t^n = (1+w)^{-\alpha-\beta} (1+bw)^{-1} \left(1 + \frac{2w}{1-x}\right)^{\alpha},$$

where

$$(17) \quad w = \frac{1}{2}(1-x)t(1+w)^{b+1}.$$

Evidently (16) reduces to the known formula [3, p. 88]

$$(18) \quad \sum_{n=0}^{\infty} P_n^{(\alpha-n, \beta-n)}(x) t^n = [1 + \frac{1}{2}(x+1)t]^{\alpha} [1 + \frac{1}{2}(x-1)t]^{\beta}$$

when $b=0$, and for $b=-1$ it leads us to Feldheim's result [5, p.120]

$$(19) \quad \sum_{n=0}^{\infty} P_n^{(\alpha-n, \beta)}(x) t^n = (1+t)^{\alpha} [1 - \frac{1}{2}(x-1)t]^{-\alpha-\beta-1}.$$

Now from the definition (14) we readily have [8, p. 61]

$$(20) \quad P_n^{(\alpha, \beta)}(x) = \binom{\alpha+n}{n} {}_2F_1 \left[\begin{matrix} -n, 1+\alpha+\beta+n; \\ 1+\alpha; \end{matrix} \middle| \frac{1-x}{2} \right],$$

whence it follows at once that

² J. W. Brown, *New generating functions for classical polynomials*, Proc. Amer. Math. Soc. 20 (1969), 263-268.

$$(21) \quad P_n^{(\alpha+bn, \beta-(b+1)n)}(x) = \binom{\alpha + (b+1)n}{n} {}_2F_1 \left[\begin{matrix} -n, 1 + \alpha + \beta; \\ 1 + \alpha + bn; \end{matrix} \frac{1-x}{2} \right].$$

Consequently, (8) gives us another class of generating functions for the Jacobi polynomial in the form

$$(22) \quad \sum_{n=0}^{\infty} P_n^{(\alpha+bn, \beta-(b+1)n)}(x) t^n = (1+v)^{\alpha+1} (1-bv)^{-1} [1 - \tfrac{1}{2}(x-1)v]^{-\alpha-\beta-1},$$

where v is defined by (1) and b, α, β are unrestricted, in general.

For $b = -1$, (22) leads us again to Feldheim's formula (19); when $b = 0$, it reduces to the generating relation

$$(23) \quad \sum_{n=0}^{\infty} P_n^{(\alpha, \beta-n)}(x) t^n = (1-t)^{\beta} [1 - \tfrac{1}{2}(x+1)t]^{-\alpha-\beta-1}$$

also due to Feldheim [5, p. 120].

Finally, we remark that the special case $b = -\frac{1}{2}$ of our formula (22) corresponds to

$$(24) \quad \sum_{n=0}^{\infty} P_n^{(\alpha-n/2, \beta-n/2)}(x) t^n = [1+u(t)]^{\alpha+1} [1+\tfrac{1}{2}u(t)]^{-1} [1 - \tfrac{1}{2}(x-1)u(t)]^{-\alpha-\beta-1},$$

where

$$(25) \quad u(t) = \tfrac{1}{2}t[t + \sqrt{(t^2 + 4)}].$$

The formula (24) appears in Brown's recent paper referred to earlier.

ADDED IN PROOF. In a private communication to the author, Professor L. Carlitz suggests that following the method of proof of the formula (9) one can readily obtain its straightforward generalization in the form

$$(*) \quad \sum_{n=0}^{\infty} \binom{a + (b+1)n}{n} t^n \sum_{k=0}^n \frac{(-n)_k c_k}{(1+a+bn)_k} \frac{x^k}{k!} = \frac{(1+v)^{\alpha+1}}{1-bv} \sum_{k=0}^{\infty} c_k \frac{(-xv)^k}{k!},$$

where the c_k are arbitrary constants and v is defined by (1). It seems worthwhile to remark here that further extensions of (*) form the subject-matter of our discussion in a forthcoming paper.

REFERENCES

1. J. W. Brown, *On zero type sets of Laguerre polynomials*, Duke Math. J. **35** (1968), 821–823.
2. L. Carlitz, *Some generating functions for Laguerre polynomials*, Duke Math. J. **35** (1968), 825–827.
3. L. Carlitz, *A bilinear generating function for the Jacobi polynomials*, Boll. Un. Mat. Ital. (3) **18** (1963), 87–89.
4. T. W. Chaundy, *An extension of hypergeometric functions*, Quart. J. Math. Oxford Ser. **14** (1943), 55–78.
5. E. Feldheim, *Relations entre les polynomes de Jacobi, Laguerre et Hermite*, Acta Math. **74** (1941), 117–138.
6. G. Pólya and G. Szegő, *Aufgaben und Lehrsätze aus der Analysis*, Springer-Verlag, Berlin, 1925.
7. H. M. Srivastava, *An extension of the Hille-Hardy formula*, Math. Comp. **23** (1969), 305–311; see also Notices Amer. Math. Soc. **15** (1968), 634–635.
8. G. Szegő, *Orthogonal polynomials*, Amer. Math. Soc. Colloq. Publ., vol. 23, Amer. Math. Soc., Providence, R. I., 1939.

WEST VIRGINIA UNIVERSITY