ON DECOMPOSABLE OPERATORS

STEPHEN PLAFKER

Let E be a Banach space, and let $\mathcal{L}(E)$ be the set of continuous linear mappings of E into E . $T \in \mathcal{L}(E)$ has the *single-valued extension property* if, for any open subset Q of $\mathbb{C}$ and any analytic function f with domain Q and values in E such that $(\lambda - T)f(\lambda) = 0$ for $\lambda \in Q$, $f = 0$. In this case, for every $x \in E$, there exists a maximally defined analytic function φ_x with values in E such that $(\lambda - T)\varphi_x(\lambda) = x$ for all λ in the domain of φ_x . The complement of the domain of φ_x in $\mathbb{C}$ will be denoted by $\text{sp}_T(x)$. For any $F \subset \mathbb{C}$ let

$$\mathfrak{M}_T(F) = \{x \in E : \text{sp}_T(x) \subset F\}.$$

From here on E will be a fixed Banach space, and T will be a fixed element of $\mathcal{L}(E)$. The symbol N will denote the set $\{2, 3, \dots\}$.

A closed subspace D of E , invariant under T , is called *spectral maximal* if $Z \subset D$ for any other closed subspace Z of E invariant under T such that $\text{sp}(T|Z) \subset \text{sp}(T|D)$.

DEFINITION. Let $n \in N$. T is *n-decomposable* [resp. *strongly n-decomposable*] if, for any open cover $(G_i)_{1 \leq i \leq n}$ of $\mathbb{C}$ consisting of n (not necessarily distinct) sets, there exists a corresponding family $(D_i)_{1 \leq i \leq n}$ of spectral maximal spaces such that $\text{sp}(T|D_i) \subset G_i$ ($1 \leq i \leq n$) and $E = D_1 + \dots + D_n$ [resp. $D = D \cap D_1 + \dots + D \cap D_n$ for every spectral maximal space D]. T is [*strongly*] *decomposable* if T is [*strongly*] *n-decomposable* for every $n \in N$.

We will need the following facts. References [1] and [2] can be consulted for proofs.

(1) If T is *n-decomposable*, then T has the single-valued extension property and $\mathfrak{M}_T(F)$ is a spectral maximal space with $\text{sp}(T| \mathfrak{M}_T(F)) \subset F \cap \text{sp}(T)$ for every closed subset F of $\mathbb{C}$. If D is a spectral maximal space, $D = \mathfrak{M}_T(\text{sp}(T|D))$.

(2) T is *strongly n-decomposable* if and only if $T|D$ is *n-decomposable* for every spectral maximal space D .

THEOREM. If $T \in \mathcal{L}(E)$ is *strongly 2-decomposable*, then $\mathfrak{M}_T(G_1 \cup G_2) = \mathfrak{M}_T(G_1) + \mathfrak{M}_T(G_2)$ for any open subsets G_1 and G_2 of $\mathbb{C}$.

PROOF. It is enough to show that $\mathfrak{M}_T(G_1 \cup G_2) \subset \mathfrak{M}_T(G_1) + \mathfrak{M}_T(G_2)$. To this end let $x \in \mathfrak{M}_T(G_1 \cup G_2)$. Then $D = \mathfrak{M}_T(\text{sp}_T(x))$ is spectral maximal, $T|D$ is 2-decomposable and, since $\text{sp}(T|D) \subset \text{sp}_T(x)$,

Received by the editors June 4, 1969.

$\{G_1, G_2\}$ is an open cover of $\text{sp}(T|D)$. Therefore there exist spectral maximal spaces D_1 and D_2 of $T|D$ such that $\text{sp}(T|D_i) \subset G_i$ ($i = 1, 2$) and $D = D_1 + D_2 \subset \mathfrak{M}_T(G_1) + \mathfrak{M}_T(G_2)$. Since $x \in D$, the result follows.

COROLLARY. *Every strongly 2-decomposable operator T is strongly decomposable.*

PROOF. By (2) above it is enough to show that T is decomposable. Let $(G_i)_{1 \leq i \leq n}$ be an open cover of $\text{sp}(T)$, and let $(H_i)_{1 \leq i \leq n}$ be an open cover of $\text{sp}(T)$ such that $\overline{H_i} \subset G_i$ for every $1 \leq i \leq n$. For each $1 \leq i \leq n$, $D_i = \mathfrak{M}_T(\overline{H_i})$ is spectral maximal, $\text{sp}(T|D_i) \subset \overline{H_i} \subset G_i$, and

$$\begin{aligned} E &= \mathfrak{M}_T(H_1 \cup \cdots \cup H_n) \\ &= \mathfrak{M}_T(H_1) + \cdots + \mathfrak{M}_T(H_n) \subset D_1 + \cdots + D_n \subset E. \end{aligned}$$

(This last step is justified by the above theorem.)

OPEN QUESTIONS. 1. Is every 2-decomposable operator strongly 2-decomposable?

2. Is every 2-decomposable operator decomposable?

An affirmative answer to Question 1 implies affirmative answers to Question 2 and Open Question (b) of [2, p. 217], which in turn provides an interesting characterization of decomposable operators (by the above theorem). The operators introduced by the author [3], which include operators studied by Colojoara, Dunford, Foias, Kantorovitz, and Maeda, are strongly decomposable.

REFERENCES

1. C. Apostol, *Restrictions and quotients of decomposable operators in a Banach space*, Rev. Roumaine Math. Pures Appl. **13** (1968), 147–150. MR 37 #6786.
2. I. Colojoara and C. Foias, *Theory of generalized spectral operators*, Gordon and Breach, New York, 1968.
3. S. Plafker, *Spectral representations for a general class of operators on a locally convex space*, Illinois J. Math. **13** (1969), 573–582.

TULANE UNIVERSITY