A NOTE ON CONVEX AND BAZILEVIČ FUNCTIONS

M. NUNOKAWA

1. Let $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ be regular, univalent in $|z| < 1$ and map $|z| < 1$ onto a domain which is starlike with respect to the origin. Then we call $f(z)$ a starlike function. It is well known that a function $f(z)$ is starlike in $|z| < 1$ if and only if

$$\operatorname{Re} \frac{zf'(z)}{f(z)} > 0 \quad \text{in } |z| < 1.$$

Let $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ be regular, univalent in $|z| < 1$ and map $|z| < 1$ onto a convex domain. Then we call $f(z)$ a convex function.

It is well known that a regular function $f(z)$ is convex if and only if

$$1 + \operatorname{Re} \frac{zf''(z)}{f'(z)} > 0 \quad \text{in } |z| < 1.$$

Every convex function is a starlike function [4].

Let $L(r)$ denote the length of the closed curve $C(r)$ which is the image of the circle $|z| = r < 1$ under the mapping $w = f(z)$ and $A(r)$ the area enclosed by $C(r)$.

Recently Thomas [6], [8] has shown that if $f(z)$ is a starlike function, then

$$(1) \quad L(r) \leq 2(\pi A(r))^{1/2} \left(1 + \log \frac{1+r}{1-r} \right).$$

In this note, for the convex functions we obtain a stronger result than (1).

THEOREM 1. *Let $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ be a convex function. Then we have*

$$L(r) = O \left(A(r) \log \frac{1}{1-r} \right)^{1/2} \quad \text{as } r \rightarrow 1.$$

PROOF. Since $f(z)$ is a univalent function we can get

Received by the editors May 26, 1969.

$$\begin{aligned}
L(r) &= \int_0^{2\pi} |zf'(z)| d\theta \leq \int_0^{2\pi} \int_0^r |zf''(z) + f'(z)| d\rho d\theta \\
&= \int_0^r \int_0^{2\pi} \left| f'(z) \left(1 + \frac{zf''(z)}{f'(z)} \right) \right| d\theta d\rho \\
&= \int_0^{r_1} \int_0^{2\pi} \left| f'(z) \left(1 + \frac{zf''(z)}{f'(z)} \right) \right| d\theta d\rho \\
&\quad + \int_{r_1}^r \int_0^{2\pi} \left| f'(z) \left(1 + \frac{zf''(z)}{f'(z)} \right) \right| d\theta d\rho
\end{aligned}$$

where r_1 is a fixed constant and $0 < r_1 < r < 1$.

Therefore we have

$$\begin{aligned}
L(r) &\leq C + \left(\int_{r_1}^r \int_0^{2\pi} \rho |f'(z)|^2 d\theta d\rho \right)^{1/2} \left(\int_{r_1}^r \int_0^{2\pi} \frac{1}{\rho} \left| 1 + \frac{zf''(z)}{f'(z)} \right|^2 d\theta d\rho \right)^{1/2} \\
&\leq C + \left(\int_0^r \int_0^{2\pi} \rho |f'(z)|^2 d\theta d\rho \right)^{1/2} \left(\frac{1}{r_1} \int_0^r \int_0^{2\pi} \left| 1 + \frac{zf''(z)}{f'(z)} \right|^2 d\theta d\rho \right)^{1/2} \\
&\leq C + \left(\frac{A(r)}{r_1} \right)^{1/2} \left(\int_0^r \int_0^{2\pi} \left| 1 + \frac{zf''(z)}{f'(z)} \right|^2 d\theta d\rho \right)^{1/2}
\end{aligned}$$

where C is a bounded constant.

On the other hand, it is well known that (see for instance [3, p. 294])

$$\int_0^r \int_0^{2\pi} \left| 1 + \frac{zf''(z)}{f'(z)} \right|^2 d\theta d\rho \leq 4\pi \log \frac{1+r}{1-r}.$$

This completes our proof and a question arises whether there is a positive constant α and a convex function $f(z)$ for which

$$L(r) \geq \alpha \left(A(r) \log \frac{1}{1-r} \right)^{1/2} \quad \text{as } r \rightarrow 1.$$

I can not give an answer for this question.

2. A function $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ that is regular in $|z| < 1$ is called a Bazilevič function of type β , if there exists a starlike function $g(z)$ and $\beta > 0$ such that

$$\operatorname{Re} \frac{zf'(z)}{f(z)^{1-\beta} g(z)^\beta} > 0 \quad \text{in } |z| < 1.$$

Bazilevič [1] has shown that each such function is univalent in $|z| < 1$. Every starlike function is a Bazilevič function of type β .

Then the following theorems have been obtained in [6], [7] and [2], [5].

THEOREM A. *Let $f(z)$ be a Bazilevič function of type β , $0 < \beta \leq 1$ and let*

$$M(r) = \max_{|z|=r} |f(re^{i\theta})| \leq (1-r)^{-\alpha} \quad 0 < \alpha \leq 2.$$

Then

$$(2) \quad L(r) = O(1-r)^{-\alpha} \quad \text{as } r \rightarrow 1.$$

THEOREM B. *Let $f(z)$ be a Bazilevič function of type β , $\arg f(z)$ be a function of bounded variation on $|z|=r < 1$ and let*

$$M(r) = \max_{|z|=r} |f(re^{i\theta})| \leq (1-r)^{-\alpha} \quad 0 < \alpha \leq 2.$$

Then we have

$$(3) \quad L(r) = O(1-r)^{-\alpha} \quad \text{as } r \rightarrow 1.$$

THEOREM 2. *The results of Theorem A and Theorem B are sharp for each α , $0 < \alpha \leq 2$ and therefore O in (2) and (3) can not be replaced by o .*

PROOF. It is easily verified that the function

$$f(z) = \frac{z}{(1-z)^\alpha}, \quad 0 < \alpha \leq 2$$

is a starlike function [2, p. 216] and

$$M(r) \leq (1-r)^{-\alpha}.$$

Applying the theorem of Fejér and Riesz to $f(z) = z/(1-z)^\alpha$ we have

$$\begin{aligned} L(r) &= \int_0^{2\pi} |zf'(z)| d\theta = \int_0^{2\pi} |z| \left| \frac{1-z+\alpha z}{(1-z)^{\alpha+1}} \right| d\theta \\ &\geq 2r \int_{-r}^r \frac{1-\rho+\alpha\rho}{(1-\rho)^{\alpha+1}} d\rho \\ &= O(1-r)^{-\alpha} \quad \text{as } r \rightarrow 1. \end{aligned}$$

This completes our proof.

The author would like to acknowledge helpful comments made by the referee.

REFERENCES

1. I. E. Bazilevič, *On a case of integrability in quadratures of the Loewner-Kufarev equation*, Mat. Sb. **37** (79) (1955), 471–476. (Russian) MR **17**, 356.
2. Ch. Pommerenke, *On starlike and convex functions*, J. London Math. Soc. **37** (1962), 209–224. MR **25** #1279.
3. ———, *On starlike and close-to-convex functions*, Proc. London Math. Soc. (3) **13** (1963), 290–304. MR **26** #2597.
4. Z. Nehari, *Conformal mapping*, McGraw-Hill, New York, 1952. MR **13**, 640.
5. M. Nunokawa, *On Bazilevič and convex functions*, Trans. Amer. Math. Soc. **143** (1969), 337–341.
6. D. K. Thomas, *On starlike and close-to-convex univalent functions*, J. London Math. Soc. **42** (1967), 427–435. MR **35** #6812.
7. ———, *On Bazilevič functions*, Trans. Amer. Math. Soc. **132** (1968), 353–361. MR **36** #5330.
8. ———, *A note on starlike functions*, J. London Math. Soc. **43** (1968), 703–706. MR **37** #4248.

GUNMA UNIVERSITY, JAPAN