ON THE IDEAL STRUCTURE OF THE ALGEBRA OF RADIAL FUNCTIONS

ALAN SCHWARTZ¹

ABSTRACT. Let L denote the convolution Banach algebra of integrable functions defined on $\mathbf{R}^n$ and let L_r consist of the sub-algebra of radial functions. If I is a closed ideal of L , the zero-set of I is defined by $Z(I) = \{y | \hat{f}(y) = 0 \text{ for all } f \in I\}$ where $\hat{f}$ is the Fourier transform of f . The following theorem is proved. If I_1 and I_2 are closed ideals of L_r such that $I_1 \subset I_2$ ($\subset$ denotes proper inclusion) then there is a closed ideal I such that $I_1 \subset I \subset I_2$.

Let n be a fixed positive integer, and let L denote the Banach algebra of integrable functions defined on $\mathbf{R}^n$ with the usual norm and convolution. (The practice of identifying two functions which agree almost everywhere will be followed.) A function f defined on $\mathbf{R}^n$ is said to be *radial* if $f(\mathbf{x}) = \phi(|\mathbf{x}|)$ for some function ϕ defined on $[0, \infty)$ and for almost every $\mathbf{x}$ in $\mathbf{R}^n$; L_r will denote the space of radial functions contained in L . A function in L is radial if and only if its Fourier transform is a radial function (see [1, pp. 69–79]), so L_r is a Banach algebra. If I is a closed ideal of L or of L_r , let

$$Z(I) = \{y | \hat{f}(y) = 0 \text{ for every } f \in I\}.$$

$Z(I)$ is called the zero-set of I .

Helson showed in [2] that if I_1 and I_2 are closed ideals of L such that $Z(I_1) = Z(I_2)$ and $I_1 \subset I_2$ ($\subset$ denotes proper inclusion), then there is a closed ideal I such that $I_1 \subset I \subset I_2$. In the present paper Helson's theorem will be used to prove the following:

THEOREM. *If I_1 and I_2 are closed ideals of L_r such that $Z(I_1) = Z(I_2)$ and $I_1 \subset I_2$, then there is a closed ideal I of L_r such that $I_1 \subset I \subset I_2$.*

The proof of the theorem will be given later; it is necessary, first, to examine how L_r sits in L .

Let $d\mu$ be the positive measure of unit mass distributed uniformly on the hypersphere $S = \{\mathbf{x} | \mathbf{x} \in \mathbf{R}^n \text{ and } |\mathbf{x}| = 1\}$, and set

Received by the editors September 5, 1969.

AMS 1969 subject classifications. Primary 4240, 4258.

Key words and phrases. Convolution algebra, Fourier transform, ideal structure, radial functions, zero-sets.

¹ Supported by an Assistant Professor Research Grant at the University of Missouri, St. Louis.

$$(1) \quad f_r(x) = \int f(|x|y) d\mu(y).$$

The integral must exist for almost every x by Fubini's theorem since $\mathbb{R}^n$ can be thought of as a product of two measure spaces: one being S with the measure $d\mu$ and the other being $[0, \infty)$ with the measure $c\rho^{n-1}d\rho$ where $d\rho$ is Lebesgue measure and c is the surface area of S . It also follows from Fubini's theorem that f_r is in L . Define

$$L_0 = \{f \mid f \in L \text{ and } f_r(x) = 0 \text{ for almost every } x \text{ in } \mathbb{R}^n\};$$

finally let $f_0(x) = f(x) - f_r(x)$. The following lemmas list some properties of L_r , L_0 , f_r , and f_0 .

LEMMA 1. *The map $f \rightarrow f_r$ is a continuous projection with unit norm, hence its null space L_0 is closed and so $L = L_0 \oplus L_r$.*

The proof of Lemma 1 follows from the easily verified facts that $\|f_r\| \leq \|f\|$ and that $f = f_r$ if f is radial.

A thorough discussion of this decomposition can be found in [3].

LEMMA 2. *f is contained in L_0 if and only if*

$$(2) \quad \int \hat{f}(\rho y) d\mu(y) = 0 \quad (\rho > 0).$$

PROOF. Application of Fubini's theorem yields

$$\begin{aligned} \int \hat{f}(\rho y) d\mu(y) &= \int_{\mathbb{R}^n} f(x) dx \left\{ \int \exp(ix \cdot \rho y) d\mu(y) \right\} \\ &= \int_{\mathbb{R}^n} f(x) K(x) dx, \end{aligned}$$

where $K(x)$ is the value of the inner integral. $K(x)$ is a radial function because μ is a weak limit of radial functions and $K(x)$ is the Fourier-Stieltjes transform of μ , or see [1, pp. 69-79]. Conversion of the last integral into hyperspherical coordinates yields (2).

To prove the converse, suppose (2) holds for some f in L . Then

$$\int \hat{f}_r(|x|y) d\mu(y) + \int \hat{f}_0(|x|y) d\mu(y) = 0.$$

The second integral vanishes by the first part of this lemma since f_0 is in L_0 , and the value of the first integral is $\hat{f}_r(x)$. Thus $\hat{f}_r = 0$, so $f_r = 0$ and hence f is in L_0 .

LEMMA 3. *The convolution of a function in L_0 and a function in L_r is contained in L_0 .*

PROOF. Suppose f is in L_r and g is in L_0 . Then for each x in $\mathbb{R}^n$

$$\begin{aligned}\int (f * g)(|x|y) d\mu(y) &= \int \hat{f}(|x|y) \hat{g}(|x|y) d\mu(y) \\ &= \hat{f}(x) \int \hat{g}(|x|y) d\mu(y) \\ &= 0\end{aligned}$$

by Lemma 2.

LEMMA 4. *Let I be a closed ideal of L_r and let K be the closed ideal of L generated by I . Then $I = K \cap L_r$ and $Z(I) = Z(K)$.*

PROOF. I is contained in K , hence in $K \cap L_r$. The fact that $K \cap L_r$ is contained in I will follow from the stronger fact that if f is in K , then f_r is in I . Suppose

$$(3) \quad f = h + \sum_{i=1}^m h_i * g_i \quad (h \in I, h_i \in I, g_i \in L; i = 1, 2, \dots, m).$$

Then $f = h + \sum_{i=1}^m h_i * (g_i)_r + \sum_{i=1}^m h_i * (g_i)_0$. The second sum is contained in L_0 by Lemma 3 and the first sum is contained in I because I is an ideal of L_r . Finally, the first sum plus h is f_r by Lemma 1; hence if f has the form (3) then f_r is in I . If f is any function in K , there is a sequence $\{f_k\}$ of finite linear combinations of the form of (3) such that f_k converges to f in L . The transformation of f into f_r is continuous on L so $\{(f_k)_r\}$ converges to f_r . Since I is closed, it must contain f_r .

Finally $Z(K) \subseteq Z(I)$ because $I \subseteq K$ and $Z(K) = Z(I)$ since finite linear combinations of the form of (3) are dense in K .

PROOF OF THEOREM. Let K_1 and K_2 be the closed ideals of L generated by I_1 and I_2 respectively. Then $K_1 \subset K_2$ by Lemma 4 because $K_1 \cap L_r = I_1 \subset I_2 = K_2 \cap L_r$, and

$$Z(K_1) = Z(I_1) = Z(I_2) = Z(K_2).$$

By Helson's theorem there must be a closed ideal K such that $K_1 \subset K \subset K_2$. Since K_2 is the ideal generated by I_2 , it follows that $K \cap L_r \subset I_2$. The inclusion $I_1 \subset K \cap L_r$ is not immediate. Suppose there is no closed ideal K of L such that

$$(4) \quad I_1 \subset K \cap L_r \subset I_2,$$

then define $\mathcal{K}$ to be the collection of all closed ideals of L such that

$$K_1 \subset K \subset K_2 \quad \text{and} \quad K \cap L_r = I_1.$$

Let $\mathcal{K}$ be ordered by inclusion and let K^* be the union of all the ideals in a maximal chain of $\mathcal{K}$.

K^* is contained in $\mathcal{K}$. To see this let J be the closure in L of K^* . If f is in $K^* \cap L_r$, then f is in $K \cap L_r$ for some K in $\mathcal{K}$ so f is in I_1 ; thus $K^* \cap L_r = I_1$ so $J \cap L_r = I_1$. Since $J \cap L_r = I_1$, it follows that $J \subset K_2$. Since K^* is a union of elements of $\mathcal{K}$ it follows that $K_1 \subset J \subset K_2$. Thus J is in $\mathcal{K}$ and so $K^* = J$ by the construction of K^* ; hence, K^* is in $\mathcal{K}$. It also follows that $Z(K^*) = Z(K_2)$ because K^* lies between K_1 and K_2 .

Helson's theorem can now be invoked to guarantee the existence of an ideal K^{**} such that $K^* \subset K^{**} \subset K_2$. Since $K^{**} \subset K_2$ it follows that $K^{**} \cap L_r \subset I_2$, and the proper inclusion $I_1 \subset K^{**} \cap L_r$ holds by the construction of K^* . Thus K^{**} contradicts our assumption that no ideal of L satisfies (4).

REFERENCES

1. S. Bochner and K. Chandrasekharan, *Fourier transforms*, Ann. of Math. Studies, no. 19, Princeton Univ. Press, Princeton, N. J., 1949. MR 11, 173.
2. H. Helson, *On the ideal structure of group algebras*, Ark. Mat. 2 (1952), 83–86. MR 14, 246.
3. E. M. Stein and G. Weiss, *An introduction to Fourier analysis in Euclidean spaces*, Princeton Univ. Press, Princeton, N. J. (to appear).
4. Sadahiro Saeki, *An elementary proof of a theorem of Henry Helson*, Tôhoku Math. J. (2) 20 (1968), 244–247. MR 37 #6694.

UNIVERSITY OF MISSOURI, ST. LOUIS, MISSOURI 63121