SQUARE INTEGRABLE SOLUTIONS OF $Ly=f(t, y)$

ANTON ZETTL

ABSTRACT. Let L be an ordinary linear differential operator and L^+ its formal adjoint. It is shown that, under suitable conditions on f , all solutions of $Ly=f(t, y)$ are in $L^2(0, \infty)$ provided that all those of $Ly=0$ and $L^+y=0$ are.

1. Introduction. For L a regular ordinary linear differential or quasi-differential operator of any order we show, under suitable conditions on f , that all solutions of

$$(1) \quad Ly = f(t, y)$$

are in $L^2(a, \infty)$, provided all solutions of

$$(2) \quad Ly = 0$$

and

$$(3) \quad L^+y = 0$$

are, where L^+ is the formal adjoint of L .

The case $Ly=L^+y=(ry')'+qy$ is a recent result of J. S. Bradley [4]. The linear case with $Ly=L^+y=y''+qy$ and $f(t, y)=h(t)y$ with $h(t)$ bounded is due to Bellman [3]. See also Halvorsen [6].

2. Notation and preliminaries. We consider first the classical operators L and L^+ defined by

$$Ly = y^{(n)} + p_{n-1}y^{(n-1)} + \dots + p_1y' + p_0y,$$

$$L^+z = (\dots (((-1)^nz' + (-1)^{n-1}p_{n-1}z)' + (-1)^{n-2}p_{n-2}z)' \dots - p_1z)' + p_0z$$

where the p_i 's for $i=0, \dots, n-1$ are continuous real valued functions on $[a, \infty)$. This form of the adjoint equation is used to avoid smoothness assumptions on the coefficients. Denote by S and S^+ the solution spaces of $Ly=0$ and $L^+y=0$, respectively. Let y_i for $i=1, \dots, n$ be the solutions of (2) determined by the initial conditions

$$y_i^{(j)}(a) = \delta_{i,j+1} \quad \text{for } j = 0, \dots, n-1.$$

Received by the editors January 16, 1970.

AMS 1969 subject classifications. Primary 3453; Secondary 4760.

Key words and phrases. Square integrable solutions, differential operators, Gronwall inequality, variation of parameters.

Throughout the paper δ will denote the Kronecker δ . Define $D_0^+ z = z$, $D_i^+ z = (D_{i-1}^+ z)' + (-1)^i p_{n-i} z$ for $i = 1, \dots, n-1$.

Let z_i for $i = 1, \dots, n$ be the solutions of (3) determined by

$$D_j^+ z_i(a) = (-1)^{n+i} \delta_{i, n-j} \quad \text{for } j = 0, \dots, n-1.$$

The proof of the main theorem will be based on two lemmas which we now state.

LEMMA 1. Suppose $f \in C[a, \infty)$. The solution y of $Ly = f$ satisfying $y^j(a) = \alpha_{j+1}$, for $j = 0, \dots, n-1$, is given by

$$(4) \quad y(t) = \sum_{i=1}^n \alpha_i y_i(t) + \int_a^t \sum_{i=1}^n y_i(t) z_i(s) f(s) ds.$$

This representation of solutions of nonhomogeneous equations differs from that found in most textbooks. So, although Lemma 1 follows from Theorems 2 and 3 in [11], we sketch a proof here for the sake of completeness.

PROOF. Let

$$(5) \quad Y' = FY$$

be the usual vector-matrix formulation of equation (2), i.e. Y is the column vector $[y, y', \dots, y^{n-1}]$ and F is the matrix having all 1's in the super diagonal, $-p_0, -p_1, \dots, -p_{n-1}$ in the bottom row from left to right and zeros elsewhere. Note that equation (3) is equivalent to the vector-matrix equation

$$(6) \quad Z' = F^+ Z$$

where Z is the column vector $[D_0^+ z, D_1^+ z, \dots, D_{n-1}^+ z]$ and $F^+ = -J^{-1} F^* J$ with $J = ((-1)^i \delta_{i, n+1-j})$. Denote by Y_u the unique matrix solution of (5) satisfying $Y(u) = I$ and by Z_u the unique matrix solution of (6) satisfying $Z(u) = I$ where I is the $n \times n$ identity matrix.

Let $M(t, u) = Y_u(t) = (M_{ij}(t, u))$ and $N(t, u) = Z_u(t) = (N_{ij}(t, u))$ for $t \geq a, u \geq a$. Noting that $M(t, s) = M(t, u)M(u, s)$ one can readily verify that

$$(7) \quad M(t, u) = J^{-1} N^*(u, t) J \quad \text{for } t \geq a, u \geq a.$$

From the variation of constants formula it follows that

$$(8) \quad y(t) = \sum_{i=1}^n M_{1i}(t, a) \alpha_i + \int_a^t M_{1n}(t, u) f(u) du.$$

As a consequence of (8) and the definitions of M , N , y_i and z_i we can obtain that

$$\begin{aligned} M_{1n}(t, s) &= \sum_{i=1}^n M_{1i}(t, a) M_{in}(a, s) \\ &= \sum_{i=1}^n M_{1i}(t, a) (-1)^{i+n} N_{1i}(s, a) = \sum_{i=1}^n y_i(t) z_i(s) \end{aligned}$$

and (4) follows.

The second lemma is a very special case of a generalization of the well-known Gronwall inequality due to H. E. Gollwitzer [5] (also see Willett [9] and Willett and Wong [10]).

LEMMA 2. *Let u , ϕ , g , h be nonnegative continuous functions on $[a, \infty)$ and suppose that*

$$u(t) \leq \phi(t) + g(t) \left(\int_a^t u^2(s) h(s) ds \right)^{1/2} \quad \text{for } t \geq a.$$

Then

$$u(t) \leq \phi(t) + g(t) \left\{ \int_a^t 2\phi^2(s) h(s) \exp \left[\int_s^t 2g(x) h(x) dx \right] ds \right\}^{1/2} \quad \text{for } t \geq a.$$

3. The main theorem. Suppose there exist functions h and k such that

$$(9) \quad |f(t, y)| \leq k(t) + h(t)y \quad \text{for } t \geq a, \quad -\infty < y < \infty.$$

THEOREM 1. *Suppose $S \cup S^+ \subset L^2(a, \infty)$ and $kz \in L^1(a, \infty)$ for all $z \in S^+$ and either*

- (i) $|h|^{1/2}|y| \in L^2(a, \infty)$ for all $y \in S \cup S^+$ or
- (ii) $hz \in L^2(a, \infty)$ for all $z \in S^+$.

Then all solutions of (1) are in $L^2(a, \infty)$.

PROOF. We prove first that all solutions of (1) are in $L^2(a, \infty)$ if (i) holds. Note that (9) implies all solutions are defined on $[a, \infty)$. Let y be a solution of (1).

By Lemma 1

$$y(t) = \sum_{i=1}^n \alpha_i y_i(t) + \int_a^t \sum_{i=1}^n y_i(t) z_i(s) f(s, y(s)) ds.$$

Hence

$$|y(t)| \leq \sum_{i=1}^n |\alpha_i| |y_i(t)| + \sum_{i=1}^n |y_i(t)| \int_a^t |z_i(s)| (|k(s)| + |h(s)y(s)|) ds.$$

Letting $C_i = \int_a^\infty |z_i k|$ and the Schwarz inequality we obtain

$$(10) \quad |y(t)| \leq \sum_{i=1}^n (C_i + |\alpha_i|) |y_i(t)| + \sum_{i=1}^n |y_i(t)| \left(\int_a^t z_i^2 |h| \int_a^t y^2 |h| \right)^{1/2}.$$

Let $D_i = (\int_a^\infty z_i^2 |h|)^{1/2}$, then $|y(t)| \leq \phi(t) + g(t) (\int_a^t y^2 |h|)^{1/2}$ where $\phi(t) = \sum_{i=1}^n (C_i + |\alpha_i|) |y_i(t)|$ and $g(t) = \sum_{i=1}^n D_i |y_i(t)|$. It follows from Lemma 2 that

$$(11) \quad |y(t)| \leq \phi(t) + g(t) \left\{ \int_a^t 2\phi^2(s) |h(s)| \cdot \exp \left[\int_a^t 2g^2(x) |h(x)| dx \right] ds \right\}^{1/2}.$$

Since $\int_a^\infty \phi^2(s) |h(s)| ds$ and $\int_a^\infty g^2(x) |h(x)| dx$ are both finite, we conclude that $|y|$ is bounded by a linear combination of ϕ and g and hence y is in $L^2(a, \infty)$. If (ii) holds, the proof is the same up to (10). In this case (10) becomes

$$|y(t)| \leq \sum_{i=1}^n (C_i + |\alpha_i|) |y_i(t)| + \sum_{i=1}^n |y_i(t)| \left(\int_a^t z_i^2 h^2 \int_a^t y^2 \right)^{1/2}.$$

Letting $D_i = (\int_a^\infty z_i^2 h^2)^{1/2}$ and proceeding as before (11) gets replaced by

$$|y(t)| \leq \phi(t) + g(t) \left\{ \int_a^t 2\phi^2(s) \exp \left[\int_a^t 2g^2(x) dx \right] ds \right\}^{1/2},$$

and the conclusion follows similarly.

4. Remarks. (1) Theorem 1 applied to the linear equation $Ly = hy$ yields: If $S \cup S^+ \subset L^2(a, \infty)$ and either $|h|^{1/2}y \in L^2(a, \infty)$ for all y in $S \cup S^+$ or $hz \in L^2(a, \infty)$ for all $z \in S^+$, then all solutions of $Ly = hy$ are in $L^2(a, \infty)$.

(2) The hypothesis on h is clearly satisfied if either h is bounded or $h \in L^1(a, \infty)$.

(3) Theorem 1 holds for general quasi-differential operators such as l and l^+ studied in [11]. Equivalent versions of these were considered by Shinn [8] and include, in particular, those used in [7] and [1]. See also J. H. Barrett [2] and many others. The argument given above can be readily extended to l and l^+ without any major modifications. In particular, Theorem 1 applies to operators of the type

$$Ly = (p_n y^n)^n + (p_{n-1} y^{n-1})^{n-1} + \cdots + p_0 y$$

which include the Sturm operator $(p_1 y')' + p_0 y$.

(4) Condition (9) is essentially the same as the one used by Bradley [4].

BIBLIOGRAPHY

1. N. I. Ahiezer and I. M. Glazman, *Theory of linear operators in Hilbert space*, GITTL, Moscow, 1950; English transl., Ungar, New York, 1961. MR 13, 358.
2. J. H. Barrett, *Oscillation theory of ordinary linear differential equations*, *Advances in Math.* 3 (1969), 415-509.
3. Richard Bellman, *A stability property of solutions of linear differential equations*, *Duke Math. J.* 11 (1944), 513-516. MR 6, 66.
4. J. S. Bradley, *Comparison theorems for the square integrability of solutions of $(r(t)y')' + q(t)y = f(t, y)$* (private communication).
5. Herman Gollwitzer, *A note on a functional inequality*, *Proc. Amer. Math. Soc.* (to appear).
6. Sigbjorn Halvorsen, *On the quadratic integrability of solutions of $d^2x/dt^2 + f(t)x = 0$* , *Math. Scand.* 14 (1964), 111-119. MR 29 #6102.
7. M. A. Naĭmark, *Linear differential operators*, GITTL, Moscow, 1954; English transl., Ungar, New York, 1967; German transl., Akademie-Verlag, Berlin, 1960. MR 16, 702; MR 35 #6885.
8. D. Shinn, *Existence theorems for the quasi-differential equation of the n^{th} order*, *C. R. Acad. Sci. USSR* 18 (1938), 515-518.
9. D. Willett, *Nonlinear vector integral equations as contraction mappings*, *Arch. Rational Mech. Anal.* 15 (1964), 79-86. MR 28 #2417.
10. D. Willett and J. S. W. Wong, *On the discrete analogues of some generalizations of Gronwall's inequality*, *Monatsh. Math.* 69 (1965), 362-367. MR 32 #2644.
11. Anton Zettl, *Quasi-differential operators in Hilbert space*, MRC Technical Summary Report #854, February, 1968.

NORTHERN ILLINOIS UNIVERSITY, DEKALB, ILLINOIS 60115