A SHORT PROOF OF BERGER'S CURVATURE TENSOR ESTIMATES

HERMANN KARCHER

ABSTRACT. A simple proof of Berger's estimate of the curvature tensor components in terms of the pinching is given and the result is extended to nonorthonormal arguments of the curvature tensor.

Let M be a Riemannian manifold with scalar product $(\ ; \)$ and curvature tensor $R: TM \times TM \rightarrow TM$. The sectional curvatures of M at p are given by

(1) $K(u, v) = (R(u, v)v; u) \cdot ((u; u)(v; v) - (u; v)^2)^{-1}$, for independent $u, v \in M_p$.

Put

(2) $\delta = \min_{u, v \in M_p} K(u, v)$, $\Delta = \max_{u, v \in M_p} K(u, v)$, $\bar{K} = \max(|\delta|, |\Delta|)$.

$(\Delta - \delta) \cdot \bar{K}^{-1}$ is the pinching of M .

Berger's result [1], [2] (originally for $\delta \geq 0$, but this is not needed as Tsagas [4] observed) is

(3) $| (R(u, v)v; w) | \leq \frac{1}{2}(\Delta - \delta)$ for orthonormal $u, v, w \in M_p$,

(4) $| (R(u, v)x; w) | \leq \frac{2}{3}(\Delta - \delta)$ for orthonormal $u, v, w, x \in M_p$.

The constants $\frac{1}{2}$ and $\frac{2}{3}$ cannot be improved. Apply Theorem 3.2.7 of [2, p. 92] to CP^n .

PROOF OF (3) AND (4). We use the curvature tensor identities

(5) $4(R(u, v)v; w) = (R(u + w, v)v; u + w) - (R(u - w, v)v; u - w)$,

(6) $6(R(u, v)x; w) = (R(u, v + x)v + x; w) - (R(u, v - x)v - x; w) - (R(v, u + x)u + x; w) + (R(v, u - x)u - x; w)$,

which are immediate from the linearity and symmetries of R . (5) and (6) express $(R(u, v)x; w)$ in a simpler way in terms of sectional curvatures than is given in [3, p. 93].

For orthonormal u, v, w we have $(u \pm w)^2 = 2$, hence (3) is clear from (5), (1), (2). If u, v, w, x are orthonormal then (3) applies to the four terms on the right of (6); together with $(v \pm x)^2 = (u \pm x)^2 = 2$ we have (4).

To formulate the extensions of (3) and (4) we interpret the end-points of three unit vectors $u, v, w \in M_p$ as vertices of a spherical tri-

Received by the editors February 26, 1970.

AMS 1969 subject classifications. Primary 5370, 5342.

Key words and phrases. Curvature tensor identities, sectional curvature, pinching, curvature tensor norm, estimation of curvatures tensor components.

angle. With the notation $(u; v)^2 = \cos^2(u, v) = 1 - \sin^2(u, v)$ the spherical cosine formula reads

$$\cos(u, w) = \cos(u, v) \cdot \cos(v, w) + \sin(u, v) \cdot \sin(v, w) \cdot \cos \angle uvw.$$

THEOREM. *The following generalization of (3) and (4) holds (for unit vectors $u, v, w, x \in M_p$)*

$$(7) \quad \begin{aligned} |R(u, v)x; w| &\leq |\sin(u, v) \cdot \sin(v, w)| \\ &\quad \cdot \left\{ \frac{1}{2}(\Delta - \delta) |\sin \angle uvw| + \bar{K} \cdot |\cos \angle uvw| \right\} \\ &\leq (\bar{K}^2 + \frac{1}{4}(\Delta - \delta)^2)^{1/2}, \end{aligned}$$

where $\cos \angle uvw = 0$ if $w \perp u, v$.

$$(8) \quad |(R(u, v)x; w)| \leq |\sin(u, v) \cdot \sin(x, w)| \cdot \frac{2}{3}(\Delta - \delta), \quad \text{for } w \perp u, v.$$

$$(9) \quad |(R(u, v)x; w)| \leq (\bar{K}^2 + (25/36)(\Delta - \delta)^2)^{1/2},$$

hence, if $\|R\|$ denotes the norm of R as quadrilinear map then $\|R\| \leq (\bar{K}^2 + (25/36)(\Delta - \delta)^2)^{1/2}$.

PROOF. Put $y = u - v(u; v)$, $z = w - v(w; v)$, then $|y| = |\sin(u, v)|$, $|z| = |\sin(w, v)|$ and $v \perp y, z$. Therefore we can apply (3) to the first term on the right of

$$\begin{aligned} (R(u, v)x; w) &= \left(R(y, v)x; z - \frac{y}{|y|^2} (y; z) \right) \\ &\quad + \left(R(y, v)x; \frac{y}{|y|^2} (y; z) \right) \end{aligned}$$

and get the first part of (7) since—by the spherical cosine formula—

$$|\cos(y, z)| = \frac{|(y; z)|}{|y||z|} = |\cos \angle uvw|$$

and

$$\begin{aligned} |y| \cdot \left| z - \frac{y}{|y|^2} (y; z) \right| &= |y||z| |\sin(y, z)| \\ &= |\sin(u, v) \cdot \sin(v, w) \cdot \sin \angle uvw|. \end{aligned}$$

The second part of (7) follows from $|A \cdot \sin \alpha| + |B \cdot \cos \alpha| \leq (A^2 + B^2)^{1/2}$. The sin-factors in (8) are trivial since with

$$u' = (u - v(u; v)) \cdot |\sin(u, v)|^{-1}, \quad x' = (x - w(x; w)) \cdot |\sin(x, w)|^{-1}$$

we have

$$(R(u, v)x; w) = |\sin(u, v) \cdot \sin(x, w)| \cdot (R(u', v)x'; w).$$

We apply (7) to the four terms on the right of (6) (with u', x' instead of u, x) and since $w \perp u', v, x'$ all the $\bar{K}$ -cos-terms vanish. Hence (8) follows using $(v+x')^2 + (v-x')^2 + (u'+x')^2 + (u'-x')^2 = 8$. In (9) we can assume $u \perp v, x \perp w$ (replace u, x by u', x'). Put $y = u - x(u; x)$, $z = v - x(v; x)$, $a = u(v; x) - v(u; x)$. Note $a \perp x$ and $y(v; x) - v(u; x) = a - x(u; x)(v; x)$. Then

$$\begin{aligned}(R(u, v)x; w) &= (R(y, z)x; w) + (R(y, x(v; x))x; w) + (R(x(u; x), v)x; w) \\ &= (R(y, z)x; w) + (R(a, x)x; w).\end{aligned}$$

From (8) we have

$$\begin{aligned}|(R(y, z)x; w)| &\leq \frac{2}{3}(\Delta - \delta) |y| |z| |\sin(y, z)| \\ &= \frac{2}{3}(\Delta - \delta) |\sin(u, x) \cdot \sin(v, x) \cdot \sin \angle uxv|,\end{aligned}$$

and from (7) we have

$$|(R(a, x)x; w)| \leq |a| \cdot (\bar{K}^2 + \frac{1}{4}(\Delta - \delta)^2)^{1/2}.$$

Observe $|a|^2 = (u; x)^2 + (v; x)^2 \leq |x|^2 = 1$ and—from the spherical cosine formula for the triangle u, x, v —

$$|\sin(u, x) \cdot \sin(v, x) \cdot \sin \angle uxv|^2 = 1 - |a|^2.$$

Then (9) follows again from $|A \cdot \sin \alpha| + |B \cdot \cos \alpha| \leq (A^2 + B^2)^{1/2}$.

REFERENCES

1. M. Berger, *Sur quelques variétés riemanniennes suffisamment pincées*, Bull. Soc. Math. France **88** (1960), 57–71. MR **24** #A3606.
2. S. I. Goldberg, *Curvature and homology*, Pure and Appl. Math., vol. XI, Academic Press, New York, 1962. MR **25** #2537.
3. D. Gromoll, W. Klingenberg and W. Meyer, *Riemannsche Geometrie im Grossen*, Lecture Notes in Math., no. 55, Springer-Verlag, Berlin and New York, 1968. MR **37** #4751.
4. G. Tsagas, *A relation between Killing tensor fields and negative pinched Riemannian manifolds*, Proc. Amer. Math. Soc. **22** (1969), 476–478. MR **39** #3429.

MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN, BONN, GERMANY