ON THE OPERATIONAL CALCULUS FOR GROUPS OF OPERATORS¹

SHMUEL KANTOROVITZ

ABSTRACT. We study some compactness properties of the natural operational calculus for groups of operators.

1. Let $\{T(t) | t \text{ real}\}$ be a strongly continuous group of operators on a Banach space X . Let M_T be the class of all complex Borel measures m on the real line R such that

$$\|m\|_T = \int \|T(t)\| d|m|(t) < \infty.$$

M_T is a Banach algebra for the usual operations with measures and for the norm $\|\cdot\|_T$. The image A_T of M_T under the Fourier transform $m \rightarrow \hat{m}$ is a Banach algebra of bounded continuous complex functions on R , with the usual pointwise operations and with the norm $\|\hat{m}\|_T = \|m\|_T$.

If m_t denotes the delta measure at t and $f_t(s) = \hat{m}_t(s) = \exp(its)$ ($t, s \in R$), then $m_t \in M_T$ ($f_t \in A_T$) and

$$(1) \quad \|f_t\|_T = \|m_t\|_T = \|T(t)\|, \quad t \in R.$$

Let F be a Banach algebra of complex functions on R which contains the functions f_t ($t \in R$). An F -operational calculus for $T(\cdot)$ is a continuous representation of F on X which sends f_t to $T(t)$ for all $t \in R$.

For $f \in A_T$, define the operator $\tau(f)$ on X by

$$(2) \quad \tau(f)x = \int T(t)x dm(t) \quad (x \in X, f = \hat{m} \in A_T).$$

τ is an A_T -operational calculus for $T(\cdot)$ with $\|\tau\| = 1$ (cf. (1)); we call it the *natural operational calculus* for $T(\cdot)$.

The operational calculus may be localized as follows. For $0 \neq x \in X$,

Received by the editors March 3, 1970.

AMS 1969 subject classifications. Primary 4730; Secondary 4750.

Key words and phrases. Group of operators, operational calculus, compact operator, compactly imbedded, regular vectors, spectral operator, quasi-nilpotent operator.

¹ Research supported by NSF Grant GP 8289.

let A_{T_x} denote the Banach space of all Fourier transforms of complex Borel measures m on R for which

$$\|m\|_{T_x} = \int \|T(t)x\|_X |m|(t) dt < \infty,$$

with the norm $\|\cdot\|_{T_x}$. The map $\tau(\cdot)x: A_{T_x} \rightarrow X$ defined by (2) is a continuous linear mapping of norm 1, such that $\tau(f_i)x = T(t)x$ and $\tau(fg)x = \tau(f)\tau(g)x$ ($f \in A_T$, $g \in A_{T_x}$).

DEFINITION 1. $x \in X$ is (strongly) regular for $T(\cdot)$ if the ratio $\|T(\cdot)x\|/\|T(\cdot)\|$ has a positive (infimum) $\limsup$ as $|t| \rightarrow \infty$.

THEOREM 1. Let $S(\cdot)$ and $T(\cdot)$ be strongly continuous groups of operators, and let $x \in X$. Then

- (i) $A_S \subset A_T$ iff $\|T(\cdot)\|/\|S(\cdot)\|$ is bounded.
- (ii) $A_{T_x} = A_T$ iff x is strongly regular for $T(\cdot)$.

PROOF. Let C'_0 be the dual of the space C_0 of all complex continuous functions with compact support on R (with the usual topology). If $h \in C_0$, $Q = \text{supp } h$, and $K = \sup_Q \|T(\cdot)\|^{-1}$ (finite!), then

$$\left| \int h dm \right| \leq K \sup |h| \cdot \|m\|_T,$$

so that M_T is topologically contained in C'_0 . Now if $A_S \subset A_T$, then $M_S \subset M_T$, and the latter (hence the former) inclusion is topological, since both spaces are Banach spaces topologically contained in C'_0 (cf. the Closed Graph Theorem). Since $f_i \in A_S$, it follows that

$$\|T(t)\| = \|f_i\|_T \leq B \|f_i\|_S = B \|S(t)\|, \quad t \in R.$$

This proves the nontrivial part of (i), and (ii) is proved by the same argument.

2. The group $T(\cdot)$ is bounded away from zero if $\inf \|T(\cdot)\| > 0$, and temperate if $\|T(t)\| = O(|t|^k)$ as $|t| \rightarrow \infty$, for some nonnegative integer k . Let

$$w^\pm = \lim_{t \rightarrow \pm \infty} t^{-1} \log \|T(t)\|$$

(cf. [2, p. 244]). The spectral norm of $T(t)$ is $\exp[tw^+]$ for $t \geq 0$ and $\exp[tw^-]$ for $t < 0$; this implies the equivalence of the following statements:

- 1. $T(\cdot)$ is bounded away from zero;
- 2. $\|T(\cdot)\| \geq 1$;
- 3. $w^- \leq 0 \leq w^+$.

In particular, *temperate groups are bounded away from 0.*

For $Q \subset R$ compact, denote by $A_T(Q)$ the closed subspace of A_T consisting of all $f \in A_T$ with support in Q . If $f = \hat{m} \in A_T(Q)$, then $dm(t) = \hat{f}(-t)dt/2\pi$, and therefore

$$(3) \quad \|f\|_T = \int \|T(t)\| |\hat{f}(-t)| dt/2\pi, \quad f \in A_T(Q).$$

THEOREM 2. *Let $S(\cdot)$ and $T(\cdot)$ be strongly continuous groups of operators. If $S(\cdot)$ is bounded away from 0 and $\|T(t)\|/\|S(t)\| \rightarrow 0$ as $|t| \rightarrow \infty$, then, for each compact $Q \subset R$ and $x \in X$, $A_S(Q)$ is compactly imbedded in A_T and the restrictions $\tau|_{A_S(Q)}$ and $\tau(\cdot)x|_{A_S(Q)}$ are compact operators. Conversely, if $S(\cdot)$ and $T(\cdot)$ are temperate and $A_S(Q)$ is compactly imbedded in A_T for some compact Q with nonvoid interior, then $\|T(t)\|/\|S(t)\| \rightarrow 0$ as $|t| \rightarrow \infty$.*

PROOF. 1. Let $S(\cdot)$ be bounded away from 0 (hence $\|S(\cdot)\| \geq 1$), and let $\|T(t)\|/\|S(t)\| \rightarrow 0$ as $|t| \rightarrow \infty$. Given $\epsilon > 0$, choose $N > 0$ such that $\|T(t)\|/\|S(t)\| < \epsilon$ for $|t| > N$. Let $Q \subset R$ be compact, and let $\{\hat{h}'_n\}$ be in the closed unit ball of $A_S(Q)$. By (3), $\{\hat{h}'_n\}$ is in the closed 2π -ball of $L^1(R)$. Therefore, a subsequence $\{\hat{h}_n\}$ converges weak* in $(L^\infty)^*$. In particular, for each $s \in R$

$$\frac{1}{2\pi} \int e^{-its} \hat{h}_n(t) dt = h_n(s)$$

converge. Since

$$|h_n(s)| \leq \frac{1}{2\pi} \|\hat{h}_n\|_{L^1} \leq 1 \quad \text{and} \quad \hat{h}_n(t) = \int_Q e^{its} h_n(s) ds,$$

it follows by Dominated Convergence that $\hat{h}_n$ converge pointwise on R and $|\hat{h}_n| \leq |Q|$ (the Lebesgue measure of Q). Now, by (3),

$$\|h_n - h_m\|_T \leq \int_{-N}^N \|T(t)\| |\hat{h}_n(-t) - \hat{h}_m(-t)| dt/2\pi + \epsilon \|h_n - h_m\|_S.$$

The first term on the right tends to 0 as $n, m \rightarrow \infty$ (Dominated Convergence), and the second term is $\leq 2\epsilon$. Hence $\{h_n\}$ is Cauchy in A_T , and the first part of the theorem follows.

2. Let $S(\cdot)$ and $T(\cdot)$ be temperate, and suppose $A_S(Q)$ is compactly imbedded in A_T for some compact Q with nonvoid interior. There exists a nonzero C^∞ function g with support in Q . Let $g_t = g f_t / \|S(t)\|$ ($t \in R$). Then $g_t \in A_S(Q)$, and by (3),

$$\begin{aligned}
\|g_t\|_S &= \int \|S(s)\| |\hat{g}(t-s)| \|S(t)\|^{-1} ds / 2\pi \\
&= \int \|S(t+u)\| \|S(t)\|^{-1} |\hat{g}(-u)| du / 2\pi \\
&\leq \int \|S(u)\| |\hat{g}(-u)| du / 2\pi = \|g\|_S \quad (t \in R).
\end{aligned}$$

Therefore, the set $\{g_t | t \in R\}$ has limit points in A_T , which are also limit points in the topology induced on A_T by the weak* topology on $L^\infty(R)$ (because $\|T(\cdot)\| \geq 1$). However, $g_t \rightarrow 0$ (as $|t| \rightarrow \infty$) in the latter topology, since for each $h \in L^1(R)$,

$$\left| \int h(s) g_t(s) ds \right| = |(\hat{hg})(t)| / \|S(t)\| \leq |(\hat{hg})(t)| \rightarrow 0$$

by the Riemann-Lebesgue lemma. Therefore $g_t \rightarrow 0$ in A_T as $|t| \rightarrow \infty$. Now

$$\begin{aligned}
\|g_t\|_T &= \int \|T(s)\| |\hat{g}(t-s)| ds / (2\pi \|S(t)\|) \\
&\geq (\|T(t)\| / \|S(t)\|) \int \|T(t-s)\|^{-1} |\hat{g}(t-s)| ds / 2\pi \\
&= c \|T(t)\| / \|S(t)\|,
\end{aligned}$$

where $c = \int \|T(u)\|^{-1} |\hat{g}(u)| du / 2\pi$ is a positive constant. Since $\|g_t\|_T \rightarrow 0$, the result follows.

The same arguments (with the estimate $\|T(s)x\| \geq \|T(t)x\| \cdot \|T(t-s)\|^{-1}$ replacing the inequality $\|T(s)\| \geq \|T(t)\| \|T(t-s)\|^{-1}$ in the last part of the proof) prove the following.

THEOREM 3. *Let $T(\cdot)$ be a strongly continuous group of operators, and let $x \in X$. If $T(\cdot)$ is bounded away from zero and x is nonregular, then $A_T(Q)$ is compactly imbedded in A_{Tx} for each compact $Q \subset R$. Conversely, if $T(\cdot)$ is temperate and $A_T(Q)$ is compactly imbedded in A_{Tx} for some compact set Q with nonvoid interior, then x is nonregular.*

COROLLARY. *Let $T(\cdot)$ be a temperate strongly continuous group of operators. Then x is a regular vector for $T(\cdot)$ iff the injection $A_T(Q) \rightarrow A_{Tx}$ is not compact for any compact set Q with nonvoid interior.*

3. Remarks. The existence of regular vectors seems to be a difficult question in general. Observe that if $S(\cdot)$ and $N(\cdot)$ are commuting

groups of operators, and $S(\cdot)$ is uniformly bounded, then the groups $T(\cdot) = S(\cdot)N(\cdot)$ and $N(\cdot)$ have the same (strongly) regular vectors. This follows from the estimate (with $K = \sup_{\mathbb{R}} \|S(\cdot)\|$):

$$\|T(t)x\|/\|T(t)\| \geq K^{-1}\|S(t)N(t)x\|/\|N(t)\| \geq K^{-2}\|N(t)x\|/\|N(t)\|.$$

In particular, every nonzero vector is strongly regular for a uniformly bounded group.

PROPOSITION. *If T is a real spectral operator of finite type, then the group $T(t) = e^{itT}$ possesses regular vectors.*

PROOF. Let $T = S + N$ be the canonical decomposition of T (cf. [1]). Then $T(\cdot) = S(\cdot)N(\cdot)$, where $S(t) = e^{itS}$ and $N(t) = e^{itN}$ are commuting groups and $S(\cdot)$ is uniformly bounded. By the preceding remark, it suffices to show that $N(\cdot)$ possesses regular vectors. Let k be the nonnegative integer defined by the requirements $N^k \neq 0$ and $N^{k+1} = 0$. Then

$$(it)^{-k}N(t) = \sum_{j=0}^{k-1} (it)^{j-k}N^j/j! + N^k/k! \rightarrow N^k/k!$$

as $|t| \rightarrow \infty$, in the uniform operator topology. Therefore,

$$\begin{aligned} \lim_{|t| \rightarrow \infty} \|N(t)x\|/\|N(t)\| &= \lim_{|t| \rightarrow \infty} \|(it)^{-k}N(t)x\|/\|(it)^{-k}N(t)\| \\ &= \|N^kx\|/\|N^k\|, \end{aligned}$$

so that each vector not in the null space of N^k is regular for $N(\cdot)$.

CONJECTURE. Every temperate group of operators possesses regular vectors.

The following example of R. Beals shows that the "finite type" hypothesis cannot be omitted in the above proposition.

For $n = 1, 2, \dots$, let N_n be the $n \times n$ matrix with $1/n$ in the first superdiagonal and 0 elsewhere, and consider $N = \sum \bigoplus N_n$ as an operator in l_1 . Since $N_n^n = 0$ and $\|N_n^k\| = n^{-k}$ for $k < n$, we have $N^k = \sum_{n > k} \bigoplus N_n^k$ and $\|N^k\| = \sup_{n > k} n^{-k} = (k+1)^{-k}$. Hence N is quasi-nilpotent. Let $N(t) = e^{tN} = \sum \bigoplus \exp(tN_n)$. One calculates that $\|\exp(tN_n)\| = \sum_{k < n} (t/n)^k/k!$. Since

$$\|N(t)\| = \sup_n \|\exp(tN_n)\|,$$

it follows that for each fixed n ,

$$(1) \quad \|\exp(tN_n)\|/\|N(t)\| \leq \|\exp(tN_n)\|/\|\exp(tN_{n+1})\| \rightarrow 0$$

as $t \rightarrow \infty$.

Let $P_n: (x_1, x_2, \dots) \rightarrow (x_1, \dots, x_{m(n)}, 0, 0, \dots)$, where $m(n) = 1 + \dots + n$. For any $x \in l_1$ and $n = 1, 2, \dots$,

$$\begin{aligned} \|N(t)x\|/\|N(t)\| &\leq \|N(t)P_n x\|/\|N(t)\| + \|N(t)(I - P_n)x\|/\|N(t)\| \\ &\leq \|\exp(tN_n)\| \|x\|/\|N(t)\| + \|(I - P_n)x\|, \end{aligned}$$

and therefore,

$$\limsup_{t \rightarrow \infty} \|N(t)x\|/\|N(t)\| \leq \|(I - P_n)x\|,$$

by (1). Letting $n \rightarrow \infty$, we obtain that the $\limsup$ is 0. Similarly, the $\limsup$ as $t \rightarrow -\infty$ is 0. Since x was arbitrary, this shows that $N(\cdot)$ has no regular vectors.

REFERENCES

1. N. Dunford, *A survey of the theory of spectral operators*, Bull. Amer. Math. Soc. **64** (1958), 217-274. MR 21 #3616.
2. E. Hille and R. S. Phillips, *Functional analysis and semi-groups*, rev. ed., Amer. Math. Soc. Colloq. Publ., vol. 31, Amer. Math. Soc., Providence, R. I., 1957. MR 19, 664.

UNIVERSITY OF ILLINOIS AT CHICAGO CIRCLE, CHICAGO, ILLINOIS 60680