ON THE STRUCTURE OF A FINITE SOLVABLE K -GROUP

MARSHALL KOTZEN

ABSTRACT. In this note we investigate the structure of a finite solvable K -group. It is proved that a finite group G is a solvable K -group if and only if G is a subdirect product of a finite collection of solvable K -groups H_i such that each H_i is isomorphic to a subgroup of G , and each H_i possesses a unique minimal normal subgroup.

This note eliminates several errors in [1]. The statement and the proof of Theorem 1 need correction. The statement of Theorem 1 should read “ $\dots$ collection of solvable K -groups $\dots$ ” rather than “ $\dots$ collection of K -groups $\dots$,” while its proof is invalid. In this note we will consider only finite groups with the notation and terminology in [1] assumed to be known.

The following results by Zacher [4] will be useful:

(1) A finite solvable group G is a K -group if and only if G contains a series of normal subgroups $1 = N_0 < N_1 < \dots < N_r = G$ such that each N_{i+1}/N_i is a maximal normal nilpotent subgroup of G/N_i and $\Phi(G/N_i) = 1$ for $i = 0, 1, \dots, r-1$.

(2) Each homomorphic image of a solvable K -group is a K -group.

THEOREM 1. *A group G is a solvable K -group if and only if G is a subdirect product of a finite collection of solvable K -groups H_i such that each H_i is isomorphic to a subgroup of G , and each H_i possesses a unique minimal normal subgroup.*

PROOF. Considering a solvable group G , we know that G is a subdirect product of a finite collection of solvable groups H_i such that for all i , H_i has a unique minimal normal subgroup. Adding the condition that G is a K -group, we denote by A the kernel of the projection of G onto H_i for some i . In particular, using the normality of A in the K -group G , it must follow that there exists a subgroup B of G such that $G = AB$ with $A \cap B = 1$. In particular, we observe that $B \cong H_i$, i.e. the

Received by the editors November 18, 1969.

AMS 1969 subject classifications. Primary 2000; Secondary 2025, 2027, 2052.

Key words and phrases. Solvable group, K -group, subdirect product, unique minimal normal subgroup, series, maximal normal nilpotent subgroup, Frattini subgroup, Fitting group, elementary abelian, G splits over $F(G)$, completely reducible, direct product.

Copyright © 1971, American Mathematical Society

direct factors in the subdirect product are isomorphic to subgroups of G . Applying (2) we note that H_i is a K -group. Thus we have established the necessity of the conditions of the theorem.

The sufficiency of the conditions of the above theorem follows by induction on the order of G . Suppose the result holds for all groups of order less than G . We denote the projection of G on H_i by π_i and the kernel of π_i by A_i , and note that $\bigcap_{i \in I} A_i = 1$. Since H_i is a K -group, $\Phi(H_i) = 1$, and since $\Phi(G)\pi_i \leq \Phi(H_i)$ for all $i \in I$, then $\Phi(G) \leq \bigcap_{i \in I} A_i = 1$. Thus for each subgroup N normal in G , $\Phi(N) = 1$ (see [2]). But this implies that the Fitting group $F(G)$ is elementary abelian, and that G splits over $F(G)$. That is, there exists a subgroup C such that $G = F(G) \cdot C$ with $F(G) \cap C = 1$. Note that G is clearly solvable, and we are assuming $G \neq 1$; thus $\Phi(G) = 1 < F(G)$. Thus the order of C is less than the order of G . Since each H_i is a solvable K -group, $F(H_i)$ is completely reducible. Thus $F(H_i)$ must be the unique minimal normal subgroup of H_i for each i . Now we note that $H_i = (F(G)\pi_i)(C\pi_i)$. If $F(G)\pi_i \cap C\pi_i = 1$, then $H_i/F(G)\pi_i \cong C\pi_i$. Thus $C\pi_i$ is a solvable K -group. If $F(G)\pi_i \cap C\pi_i = B_i \neq 1$, then $B_i \leq F(G)\pi_i \leq F(H_i)$, and $B_i \neq 1$ implies $B_i = F(H_i)$. Furthermore since $F(G)\pi_i \leq F(H_i)$, it follows that $F(G)\pi_i \leq C\pi_i = H_i$. (I.e. since $F(G)\pi_i \leq F(H_i) = B_i$, and $B_i \leq C\pi_i$; $F(G)\pi_i \leq C\pi_i$. Furthermore since $H_i = (F(G)\pi_i)(C\pi_i)$, $g \in H_i$ implies $g = fk$ where $f \in F(G)\pi_i$, $k \in C\pi_i$. Thus $f \in F(H_i)$, $k \in C\pi_i$ implies $f \in B_i \leq C\pi_i$, and $k \in C\pi_i$ and all this implies $fk = g \in C\pi_i$.) Again $C\pi_i$ is a solvable K -group. Since for each $i \in I$, $C\pi_i$ is a solvable K -group, it follows that each $C\pi_i$ is a subdirect product of solvable K -groups each having precisely one minimal normal subgroup. If we combine the H_i for which $C\pi_i = H_i$ together with the direct factors in the subdirect product associated with those $C\pi_i$ for which $C\pi_i \neq H_i$, a direct product can be formed for which C is a subdirect product of solvable K -groups containing precisely one minimal normal subgroup where we use composition of mappings where necessary. By the inductive hypothesis C is a solvable K -group. But this implies that in $G = F(G) \cdot C$ there exists a series of subgroups $1 = N_0 < N_1 < \dots < N_r = G$ such that N_i/N_{i-1} is the maximal normal nilpotent subgroup of G/N_{i-1} and $\Phi(G/N_{i-1}) = 1$. Now using Zacher's result (1) above, we note that G is a K -group.

I thank Professor Bechtell for suggesting this paper to me.

BIBLIOGRAPHY

1. H. Bechtell, *A note on finite solvable K-groups*, Proc. Amer. Math. Soc. 17 (1966), 1447-1450. MR 35 #4302.

2. W. Gaschütz, *Über die Φ -Untergruppe endlichen Gruppen*, Math. Z. **58** (1953), 160–170. MR 15, 285.
3. M. Suzuki, *Structure of a group and the structure of its lattice of subgroups*, Ergebnisse der Math., Heft 10, Springer-Verlag, Berlin, 1956. MR 18, 715.
4. G. Zacher, *Caratterizzazione dei gruppi risolubili d'ordine finite complementati*, Rend. Sem. Mat. Univ. Padova **22** (1953), 113–122. MR 15, 286; MR 16, 1336.

STATE COLLEGE AT WORCESTER, WORCESTER, MASSACHUSETTS 01602