A NOTE ON EXTENSIONS OF FREE GROUPS BY TORSION GROUPS¹

PAUL HILL

ABSTRACT. Under the assumption of the continuum hypothesis, a torsion free group G is constructed that is the extension of a free group F by a reduced primary group T such that G cannot be embedded in a product of Z 's. In particular, G is not free. This settles a question raised by Griffith.

In a paper just published, Griffith [3] has shown that if the torsion free abelian group G is an extension of a free group F by a torsion group T then G itself must be free provided that T is weak S_p -projective for some reduced cotorsion functor S_p . In particular, this is the case if T is a subgroup of a totally projective primary group. Specializing much further, one has that G is free if G is the extension of a free group F by any countable, reduced primary group T . In [3] Griffith raised, and left unsettled, the following question: does the above result in fact hold for an arbitrary (uncountable) reduced, primary group T ? In this note, we answer the question negatively. Unfortunately, our proof requires the continuum hypothesis, but we suspect that our result itself does not. At the very least we may save one from the futile effort of trying to generalize Griffith's theorem, without additional hypotheses, to an arbitrary reduced primary T .

Our notation is more or less standard; in particular, c denotes the cardinality of the continuum and Ω denotes the first uncountable ordinal. If B is a direct sum of cyclic p -groups, $\overline{B}$ denotes the torsion completion of B . In this connection, recall (from [2]) that any primary group T without elements of infinite height is isomorphic to some pure subgroup between B and $\overline{B}$, where B is a basic subgroup of T . $\sum_{n < \omega} C(p^n)$ is called the standard basic p -group.

THEOREM 1. *Let B be any unbounded basic p -group, that is, let B be any unbounded primary group that can be written as a direct sum of cyclic groups. If the continuum hypothesis is assumed, then there exists a torsion free group G that is not free but has a free subgroup F such that $G/F = \overline{B}$, the torsion completion of B .*

Received by the editors May 7, 1970.

AMS subject classifications. Primary 2030; Secondary 2048, 2051.

Key words and phrases. Torsion free abelian group, direct limits, free groups, extensions of groups, free groups by torsion groups.

¹ This research was partially supported by NSF Grant GP-12318.

Copyright © 1971, American Mathematical Society

PROOF. Given B , we shall construct a group G with the desired properties. By Lemma 50.1 in [2], it is enough to construct a nonfree group G with a free subgroup F such that G/F is any primary group without elements of infinite height having B as a basic subgroup. Furthermore, it suffices to do this for a countable B , for once we obtain such a group G we can add to it an appropriate free group and get

$$\bar{B} + \bar{C} = \overline{B + C}$$

as a homomorphic image, still with free kernel, where C is arbitrary. Thus let B denote an unbounded, countable p -group.

The key to the construction of G is to do more than required; we shall actually construct G such that it does not even have Z as a direct summand. In particular, G will not be separable, for it is necessarily $\aleph_1$ -free [3]. Before we begin the construction of G , let G_α denote, for each countable ordinal α , the free group of countably infinite rank and let B_α denote an isomorphic copy of B . We shall obtain G as a direct limit of the G_α 's. For simplicity of notation, we shall merely identify, in the appropriate manner, G_α in G_β if $\alpha < \beta < \Omega$ and let $G = \bigcup_{\alpha < \Omega} G_\alpha$. Thus the construction of G is accomplished essentially by the decision of how we shall identify G_α in $G_{\alpha+1}$ for each countable α . Since we do not want G to have a homomorphism onto Z , the choice of the identification of G_α in $G_{\alpha+1}$ will be heavily influenced by our desire to place obstruction to the extension to $G_{\alpha+1}$ of a map from G_α to Z . For each countable ordinal α , denote by E_α the set of epimorphisms of G_α onto (a fixed) Z . There are exactly c such epimorphisms. In view of the continuum hypothesis, the elements of E_α are in one-to-one correspondence with the countable ordinals; let

$$E_\alpha = \{ \pi_{\alpha,0}, \pi_{\alpha,1}, \dots, \pi_{\alpha,\beta}, \dots \}, \quad \beta < \Omega.$$

Set $E = \bigcup_{\alpha < \Omega} E_\alpha$ and put E into one-to-one correspondence with the countable ordinals with a function $f: E \rightarrow [0, \Omega)$ such that $f(\pi_{\alpha,\beta}) \geq \alpha$ for all $\alpha < \Omega$.

To begin the construction of G , let ϕ_0 be an epimorphism of G_0 onto B_0 with kernel K_0 ,

$$K_0 \rightarrow G_0 \rightarrow B_0.$$

Let $\gamma < \Omega$. Suppose that we have an embedding of G_α into G_β , an embedding of B_α into B_β , and that we have an epimorphism ϕ_α from G_α onto B_α if $\alpha \leq \beta < \gamma$ such that the following conditions are satisfied:

(1) G_β/G_α is p -divisible if $\alpha < \beta < \gamma$. (2) ϕ_β is an extension of ϕ_α and the kernel of ϕ_α is a direct summand of the kernel of ϕ_β if $\alpha < \beta < \gamma$.

(3) The map $\pi_{\alpha, \beta}: G_{\alpha} \rightarrow Z$ cannot be extended to a homomorphism from G_{λ} into Z whenever $f(\pi_{\alpha, \beta}) + 1 = \lambda < \gamma$. (4) B_{α} is a basic subgroup (in the set-theoretic sense) of B_{β} if $\alpha < \beta < \gamma$.

We wish to embed the G_{α} 's into G_{γ} , the B_{α} 's into B_{γ} , and to produce an epimorphism ϕ_{γ} from G_{γ} onto B_{γ} in such a way that conditions (1)–(4) remain valid, respectively, for $\beta \leq \gamma$ or $\lambda \leq \gamma$. As usual, there are two cases depending on whether γ is a limit ordinal or not.

Case 1. γ is a limit. Set $B_{\gamma} = \bigcup_{\alpha < \gamma} B_{\alpha}$ and set $G_{\gamma} = \bigcup_{\alpha < \gamma} G_{\alpha}$; we remark that B_{γ} and G_{γ} are already structurally determined, but the union of the B_{α} 's is necessarily isomorphic to B_{γ} in view of condition (4) and the union of the G_{α} 's is by Griffith's theorem [3] necessarily isomorphic to G_{γ} . Thus these identifications are admissible, and the induction obviously survives if we put $\phi_{\gamma} = \sup \{\phi_{\alpha}\}$. Note that the extension of (3) to $\lambda \leq \gamma$ is vacuous because γ is a limit ordinal.

Case 2. $\gamma - 1$ exists. Let α and β be the countable ordinals such that $f(\pi_{\alpha, \beta}) = \gamma - 1$; recall that $f(\pi_{\alpha, \beta}) \geq \alpha$, so $\alpha \leq \gamma - 1$. We know that $G_{\gamma-1} = \sum_{i < \omega} \{x_i\}$ is free and that $B_{\gamma-1} = \sum_{i < \omega} \{a_i\}$ is a direct sum of cyclic p -groups. According to [1], we can choose these decompositions such that $\phi_{\gamma-1}(x_i) = a_i$ for each i . As we already observed, there is no loss of generality in assuming that the order of a_{i+1} is greater than the order of a_i for each i —the same argument that was used in the reduction to a countable B applies. Hereafter, we shall make this assumption. In what follows, if $\pi_{\alpha, \beta}$ cannot be extended to a homomorphism from $G_{\gamma-1}$ into Z , then its obstruction has already been obtained and the induction step simplifies. In particular, in that case, the nonnegative integer N can be chosen arbitrarily. Note that if $\pi_{\alpha, \beta}$ does extend to $G_{\gamma-1}$ then its extension is unique because $G_{\gamma-1}/G_{\alpha}$ is p -divisible (and Z is p -reduced). We shall continue to denote this unique extension by $\pi_{\alpha, \beta}$, so assume that $\pi_{\alpha, \beta}: G_{\gamma-1} \rightarrow Z$. Let N be the smallest nonnegative integer such that $\pi_{\alpha, \beta}$ maps $\sum_{i \leq N} \{x_i\}$ onto Z . By the choice of N , we have $\pi_{\alpha, \beta}(x_N) \neq 0$. A crucial point is that we can assume that $\pi_{\alpha, \beta}(x_i) = 0$ if $i > N$ and maintain the condition that $\phi_{\gamma-1}(x_i) = a_i$ for each i because of the increasing orders of the a_i 's. Denote the order of a_i by $p^{e(i)}$ and set $e(i+1) = e(i) + d(i)$. Choose a prime $q \neq p$ such that $\pi_{\alpha, \beta}(x_N)$ has height zero in Z at the prime q . Set $G_{\gamma} = \sum_{i < \omega} \{y_i\}$ and embed $G_{\gamma-1}$ into G_{γ} according to the following relations:

$$\begin{aligned}
 x_i &= y_i & \text{if } i < N, \\
 (*) \quad x_N &= qy_N + p^{d(N)}y_{N+1}, \\
 x_i &= y_i + p^{d(i)}y_{i+1} & \text{if } i > N.
 \end{aligned}$$

Likewise, we set $B_\gamma = \sum_{i < \omega} \{b_i\}$ where b_i has order $p^{e(i)}$, and we embed $B_{\gamma-1}$ into B_γ by the relations:

$$\begin{aligned}
 a_i &= b_i & \text{if } i < N, \\
 a_N &= qb_N + p^{d(N)}b_{N+1}, \\
 a_i &= b_i + p^{d(i)}b_{i+1} & \text{if } i > N.
 \end{aligned}
 \tag{**}$$

It is well known and straightforward to verify that (**) embeds $B_{\gamma-1}$ into B_γ as a basic subgroup, so condition (4) holds for $\beta \leq \gamma$. An immediate consequence of (*) is that $G_\gamma/G_{\gamma-1}$ is p -divisible. Thus condition (1) is satisfied for $\beta \leq \gamma$. Clearly, the map $\phi_\gamma: y_i \rightarrow b_i$ extends $\phi_{\gamma-1}: x_i \rightarrow a_i$ because of the consistency of (*) and (**). Furthermore, the kernel K_γ of ϕ_γ is simply $\sum \{p^{e(i)}y_i\}$, whereas the kernel $K_{\gamma-1}$ of $\phi_{\gamma-1}$ is

$$\sum_{i < N} \{p^{e(i)}y_i\} + \{qp^{e(N)}y_N + p^{e(N+1)}y_{N+1}\} + \sum_{i > N} \{p^{e(i)}y_i + p^{e(i+1)}y_{i+1}\}.$$

Observe that $K_\gamma = \{p^{e(N)}y_N\} + K_{\gamma-1}$. Therefore, condition (2) is retained for $\beta \leq \gamma$. In order to show that condition (3) remains valid for $\lambda \leq \gamma$, we need to show only that $\pi_{\alpha, \beta}$ cannot be extended to a homomorphism of G_γ into Z . Suppose that $\pi_{\alpha, \beta}$ extends to G_γ . Then $\pi_{\alpha, \beta}(y_{N+1}) = 0$ since $x_i = y_i + p^{d(i)}y_{i+1}$ for all $i > N$ and since $\pi_{\alpha, \beta}(x_i) = 0$ for all $i > N$. Thus $\pi_{\alpha, \beta}(x_N) = q\pi_{\alpha, \beta}(y_N)$ which contradicts the fact that $\pi_{\alpha, \beta}(x_N)$ has zero q -height in Z . This completes the verification of the induction hypotheses.

Set $G = \bigcup_{\alpha < \Omega} G_\alpha$, $C = \bigcup_{\alpha < \Omega} B_\alpha$ and let $\phi = \sup \{\phi_\alpha\}_{\alpha < \Omega}$. Then ϕ is a homomorphism from G onto C and its kernel K is free (because K_α splits out of K_β if $\alpha < \beta$ and K_α is free for each α). Obviously, condition (3) implies that G admits no homomorphism onto Z . Condition (4) implies that C has no elements of infinite height and that B is a basic subgroup of C .

For the convenience of possible future reference, we record what we have proved as it relates to the standard basic p -group.

THEOREM 2. *Let B be the standard basic p -group. If the continuum hypothesis is assumed, then there exists a torsion free group G that does not have Z as a homomorphic image but has a free subgroup F such that G/F is between B and $\overline{B}$.*

COROLLARY. *If the continuum hypothesis is assumed, there exists a (slender and $\aleph_1$ -free) torsion free group G that is the extension of a free group F by a reduced primary group T such that G cannot be embedded in a product of Z 's.*

REFERENCES

1. J. M. Cohen and H. Gluck, *Stacked bases for modules*, Bull. Amer. Math. Soc. **75** (1969), 978–979.
2. L. Fuchs, *Abelian groups*, Akad. Kiadó, Budapest, 1958; republished by Internat. Series of Monographs on Pure and Appl. Math., Pergamon Press, New York, 1960. MR 21 #5672; MR 22 #2644.
3. P. Griffith, *Extensions of free groups by torsion groups*, Proc. Amer. Math. Soc. **24** (1970), 677–679.

FLORIDA STATE UNIVERSITY, TALLAHASSEE, FLORIDA 32306