FACTORIZATION OF DIFFERENTIAL OPERATORS

ANTON ZETTL

ABSTRACT. A necessary and sufficient condition for a differential operator of order n to be factorable into a product of operators of orders $n-r$ and r , for any $0 < r < n$, is given.

Consider differential operators of the form

$$(*) \quad Ly = s_0 y^{(n)} + s_1 y^{(n-1)} + \dots + s_n y$$

with s_i continuous for $i=0, \dots, n$ and $s_0(t) \neq 0$.

THEOREM. Suppose L is an operator of order n of type $(*)$ and $0 < r < n$. Then there exist operators P and Q both of type $(*)$ of orders $n-r$ and r respectively such that $L=PQ$ on some interval I if and only if there exist r linearly independent solutions $y_1, \dots, y_r$ of $Ly=0$ satisfying the condition that the Wronskian $W_r = W_r(y_1 \dots y_r) \neq 0$ on I .

PROOF. Suppose $L=PQ$ where Q has order r . Since any solution of $Qy=0$ is also a solution of $Ly=0$, we can choose any r linearly independent solutions of $Qy=0$ and condition $W_r \neq 0$ will be satisfied.

On the other hand, assume $Ly_i=0$ for $i=1, \dots, r$ and $W_r(y_1 \dots y_r) \neq 0$. Let

$$Qy = \det \begin{bmatrix} y_1 & \dots & y_r & y \\ y'_1 & \dots & y'_r & y' \\ \vdots & & \vdots & \vdots \\ y_1^{(r)} & \dots & y_r^{(r)} & y^{(r)} \end{bmatrix} = q_0 y^{(r)} + q_1 y^{(r-1)} + \dots + q_r y.$$

Note that $q_0 = W_r(y_1 \dots y_r) \neq 0$ and $q_i \in C^{n+1-r-i}$. A direct computation shows that the coefficients p_i for $i=0, \dots, n-r$ can be chosen successively so that the coefficients of $y^{(n)}, y^{(n-1)}, \dots, y^{(n-r)}$ in the product PQy where $Py = p_0 y^{n-r} + \dots + p_{n-r} y$ are $s_0, s_1, \dots, s_r$, respectively. Hence the operator $N=L-PQ$ is of order less than r . But $y_1, \dots, y_r$ are linearly independent solutions of $Ny=0$. Hence $N=0$ and $L=PQ$.

Although the above theorem may have been known to Ince—see

Received by the editors July 20, 1970.

AMS 1970 subject classifications. Primary 34A30, 34A05; Secondary 34A01.

Key words and phrases. Ordinary differential equations, factoring differential operators, Frobenius factorization, Pólya's property " W ", Wronskians, linear homogeneous differential equations.

the discussion on pp. 119 and 126–127 in [1]—the author has not seen an explicit statement of it anywhere in the literature.

The conditions $W_r \neq 0$ for $r=1, \dots, n-1$ are known to be necessary and sufficient for the factorability of an operator of type (*) into a “product” of first order operators—see [2].

REFERENCES

1. E. L. Ince, *Ordinary differential equations*, Dover, New York, 1944. MR 6, 65.
2. G. Pólya, *On the mean value theorem corresponding to a given linear homogeneous differential equation*, Trans. Amer. Math. Soc. 24 (1922), 312–324.

NORTHERN ILLINOIS UNIVERSITY, DE KALB, ILLINOIS 60115