SHORTER NOTES

The purpose of this department is to publish very short papers of an unusually elegant and polished character, for which there is no other outlet.

SERIES CONVERGENCE ON BOOLEAN ALGEBRAS

MILOSLAV NOSAL¹

ABSTRACT. Necessary and sufficient conditions for convergence of an infinite series $\sum_{n=1}^{\infty} a_n$ of elements of an abstract boolean σ -algebra is that $\lim_{n \rightarrow \infty} a_n = 0$.

We will base our investigation on the monograph [1] by Carathéodory, and define an abstract boolean σ -algebra $\mathcal{B}$ as an idempotent ring with a unit which is σ -complete (see also [3]). Let us denote addition by $\sum$, multiplication by $\cap$ and the corresponding join by $\cup$.

Further, if $x_n \in \mathcal{B}$, $n = 1, 2, 3, \dots$, then $x^\sim = \limsup x_n = \bigcap_{n=1}^{\infty} \bigcup_{i=n}^{\infty} x_i$, $x_\sim = \liminf x_n = \bigcup_{n=1}^{\infty} \bigcap_{i=n}^{\infty} x_i$, $x = \lim_{n \rightarrow \infty} x_n \Leftrightarrow_{\text{def}} x^\sim = x_\sim = x$.

Finally we define $\sum_{n=1}^{\infty} x_n = \lim_{n \rightarrow \infty} \sum_{i=1}^n x_i$.

THEOREM. Let $\mathcal{B}$ be an abstract boolean σ -algebra, $a_n \in \mathcal{B}$ for $n = 1, 2, 3, \dots$. Then $\sum_{n=1}^{\infty} a_n$ exists if and only if $\lim_{n \rightarrow \infty} a_n = 0$.

PROOF. The following statements are consequences of properties of the operations $\sum$, $\cup$, $\cap$ and can be found proven for example in §§1–32 of [1]. Necessity of the condition is an obvious consequence of continuity of $\sum$. Let $a'_n = 1 + a_n$, $d_n = \sum_{i=1}^n a_i$. Then

$$\begin{aligned} 1 &= \limsup a_n + \liminf a'_n \\ &= 0 + \liminf a'_n = \bigcup_{n=1}^{\infty} \bigcap_{i=n}^{\infty} a'_i; \\ \limsup d_n &= 1 \cap \limsup d_n = \left(\bigcup_{n=1}^{\infty} \bigcap_{i=n}^{\infty} a'_i \right) \cap \left(\bigcap_{m=1}^{\infty} \bigcup_{j=1}^{\infty} d_j \right) \\ &= \bigcup_{n=1}^{\infty} \bigcap_{i=n}^{\infty} \bigcap_{m=1}^{\infty} \bigcup_{j=m}^{\infty} (a'_i \cap d_j) \subset \bigcup_{n=1}^{\infty} \bigcap_{i=n}^{\infty} \bigcup_{j=n}^{\infty} (a'_i \cap d_j). \end{aligned}$$

Received by the editors August 25, 1970 and, in revised form, November 24, 1970.
AMS 1970 subject classifications. Primary 02J05, 06A40, 06A70; Secondary 02K30.

Key words and phrases. Boolean rings, boolean algebras, convergence on boolean algebras, infinite series, series convergence.

¹ This work was partially supported by the National Research Council of Canada, grant no. A 7347.

$$\begin{aligned}
\bigcap_{i=n}^{\infty} \bigcup_{j=n}^{\infty} (a'_i \cap d_j) &= \bigcup_{j=n}^{\infty} \bigcap_{i=j+1}^{\infty} a'_i \cap \left[\left(\sum_{k=1}^{n-1} a_k \right) \cap \left(\bigcap_{i=n}^j a'_i \right) \right. \\
&\quad \left. + \left(\sum_{k=n}^j a_k \right) \cap \left(\bigcap_{i=n}^j a'_i \right) \right] \\
&= \bigcup_{j=n}^{\infty} \bigcap_{i=j+1}^{\infty} a'_i \cap \left[d_{n-1} \cap \bigcap_{i=n}^j a'_i + \sum_{k=n}^j \bigcap_{i=n}^j (a_k \cap a'_i) \right] \\
&= d_{n-1} \cap \bigcup_{j=n}^{\infty} \bigcap_{i=n}^{\infty} a'_i = d_{n-1} \cap \bigcap_{i=n}^{\infty} a'_i
\end{aligned}$$

because $\sum_{k=n}^j \bigcap_{i=n}^j (a'_i \cap a_k) = 0$.

For $k \geq n$,

$$\begin{aligned}
d_k \cap \bigcap_{i=n}^{\infty} \bigcup_{j=n}^{\infty} (a'_i \cap d_j) &= d_k \cap d_{n-1} \cap \bigcap_{i=n}^{\infty} a'_i \\
&= d_{n-1} \cap \left(d_{n-1} + \sum_{l=n}^k a_l \right) \cap \bigcap_{i=n}^{\infty} a'_i \\
&= d_{n-1} \cap \bigcap_{i=k+1}^{\infty} a'_i \cap \left(d_{n-1} \cap \bigcap_{i=n}^k a'_i + \left(\sum_{l=n}^k a_l \right) \cap \left(\bigcap_{i=n}^k a'_i \right) \right) \\
&= d_{n-1} \cap \bigcap_{i=n}^{\infty} a'_i.
\end{aligned}$$

Therefore $\bigcap_{i=n}^{\infty} \bigcup_{j=n}^{\infty} (a'_i \cap d_j) \subset d_k$ for every $k \geq n$. Finally we have

$$\limsup d_n \subset \bigcup_{n=1}^{\infty} \bigcap_{i=n}^{\infty} \bigcup_{j=n}^{\infty} (a'_i \cap d_j) \subset \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} d_k = \liminf d_n. \quad \text{Q.E.D.}$$

COMMENT. Apparently existence of the unit in $\mathcal{B}$ is not substantial. All we need is the existence of an upper bound m for any sequence of elements a_n in $\mathcal{B}$.

ACKNOWLEDGEMENT. I am very indebted to Norbert Sauer for inspiring discussions.

REFERENCES

1. C. Carathéodory, *Algebraic theory of measure and integration*, Chelsea, New York, 1963. MR 26 #6340.
2. P. R. Halmos, *Measure theory*, Van Nostrand, Princeton, N.J., 1950. MR 11, 504.
3. ———, *Lectures on Boolean algebras*, Van Nostrand Math. Studies, no. 1, Van Nostrand, Princeton, N. J., 1963. MR 29 #4713.
4. R. Sikorski, *Boolean algebras*, 3rd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 25, Springer-Verlag, Berlin, 1969. MR 39 #4053.

UNIVERSITY OF CALGARY, CALGARY, ALBERTA, CANADA