ON p -CLOSE-TO-STAR FUNCTIONS OF ORDER α

HASSOON S. AL-AMIRI¹

ABSTRACT. In a recent paper M. Finkelstein has used an iterated form of Schwarz's lemma to obtain growth estimates for convex functions and starlike functions of order zero. These estimates involve the second coefficient. W. E. Chase has announced results about the growth estimates and also about a radius of univalence and starlikeness for p -close-to-star functions of order zero. The author generalizes the above results to functions of order α , $0 \leq \alpha \leq 1$.

1. Introduction. Let $C(\alpha, p)$ denote the class of functions analytic in the unit disc E ($|z| < 1$) and of the form

$$c(z) = z + b_2 z^2 + \cdots,$$

such that $\operatorname{Re}\{zc''(z)/c'(z) + 1\} > \alpha$, $z \in E$, where $|b_2| = p$ and $0 \leq \alpha \leq 1$. A function $c(z) \in C(\alpha, p)$ is said to be convex of order α . The functions in $C(\alpha, p)$ are univalent in E and $C(0, p)$ is called convex. Note that $0 \leq p \leq 1$, and if $c(z)$ is in $C(1, p)$, then $c(z) = z$.

Let $S(\alpha, p)$ denote the class of functions analytic in E and of the form

$$s(z) = z + a_2 z^2 + \cdots,$$

such that $\operatorname{Re}\{zs'(z)/s(z)\} > \alpha$, $z \in E$, where $|a_2| = 2p$ and $0 \leq \alpha \leq 1$. A function $s(z) \in S(\alpha, p)$ is said to be starlike of order α . The functions in $S(\alpha, p)$ are univalent in E and $S(0, p)$ is called starlike. Note that $0 \leq p \leq 1$, and if $s(z)$ is in $S(1, p)$, then $s(z) = z$.

Let $S_p(\alpha)$ denote the class of functions analytic in E and of the form

$$g(z) = z + d_2 z^2 + \cdots,$$

such that $\operatorname{Re}\{g(z)/s(z)\} > 0$ for $z \in E$ and for some $s(z) \in S(\alpha, p)$. A function $g(z) \in S_p(\alpha)$ is said to be p -close-to-star of order α . Functions in $S_p(\alpha)$ need not be univalent in E as can be seen by $g(z) = (z^2 + z)/(1 - z)$, with $s(z) = z$. A function in $S_p(0)$ is called p -close-to-star.

Received by the editors May 11, 1970.

AMS 1968 subject classifications. Primary 3040, 3042; Secondary 3050, 3065.

Key words and phrases. Univalent functions, convex functions of order α , starlike functions of order α , p -close-to-star functions of order α , Schwarz lemma, functions of positive real part, radius of univalence, radius of starlikeness.

¹Supported in part by the Research Faculty Grant at Bowling Green State University.

In [2], M. Finkelstein has obtained growth estimates of $|c(z)|$, $|c'(z)|$, $|c''(z)|$ and $|s(z)|$, where $c(z)$ and $s(z)$ are functions in $C(0, p)$ and $S(0, p)$, respectively. In [1], W. Chase has obtained, among other things, the radius of univalence and starlikeness of functions in the class $S_p(0)$.

The purpose of this paper is to generalize the results of [1] and [2] to functions in $C(\alpha, p)$, $S(\alpha, p)$ and $S_p(\alpha)$.

2. **The classes $C(\alpha, p)$ and $S(\alpha, p)$.** An iterated form of Schwarz's lemma is used which we state next.

LEMMA A. Let $f(z) = a_1 z + \dots$ be an analytic map of the unit disc into itself. Then $|a_1| \leq 1$ and

$$|f(re^{i\theta})| \leq r(r + |a_1|)/(1 + |a_1|r).$$

Equality holds at some point $z \neq 0$ if and only if

$$f(z) = e^{-it}z(z + ae^{it})/(1 + \bar{a}e^{-it}z), \quad t \geq 0.$$

REMARK 1. With an appropriate rotation and with no loss of generalities, the second coefficients in $C(\alpha, p)$, $S(\alpha, p)$ may be replaced by p and $2p$, respectively. Also, as a consequence of Lemma A, the order is related to p by $\alpha + p \leq 1$, as will be seen in the proof of Theorem 1. Thus in the sequel we assume $\alpha + p \leq 1$. The trivial cases where $p = 0$ and $\alpha = 1$ are omitted for the obvious reason.

REMARK 2. We could not exhibit extremal functions for Theorems 1, 2 and 3 in the general case since the functions are in an integral form of a complicated nature. However, for the special case where $\alpha = 0$, the extremal functions are those offered in [2].

THEOREM 1. Let $c(z) = z + pz^2 + \dots$ be in $C(\alpha, p)$, $p > 0$. Then

$$(1) \quad |c'(re^{i\theta})| \leq \left(\frac{1}{1-r^2}\right)^{1-\alpha} \left(\frac{1+r}{1-r}\right)^p,$$

where $z = re^{i\theta}$ is in E . Also $p + \alpha \leq 1$.

PROOF. For $p > 0$, let

$$h(z) = (1/(1-\alpha))(zc''(z)/c'(z)) = (2p/(1-\alpha))z + \dots$$

Then

$$\operatorname{Re}\{(1/(1-\alpha))zc''(z)/c'(z) + 1\} = (1/(1-\alpha))\operatorname{Re}\{zc''(z)/c'(z) + 1 - \alpha\} > 0.$$

So, there exists an analytic function $K(z)$ such that

$$(2) \quad 1 + h(z) = (1 + K(z))/(1 - K(z)),$$

where $K(0)=0$, $|K(z)| \leq 1$ for $z \in E$. Note that

$$K(z) = (p/(1-\alpha))z + \dots$$

Applying Lemma A to $K(z)$, we get $p+\alpha \leq 1$, and

$$(3) \quad |K(z)| \leq r((1-\alpha)r + p)/(1-\alpha + pr).$$

From (2) and (3), it follows that

$$|h(z)| = |2K/(1-K)| \leq 2r((1-\alpha)r + p)/(1-\alpha)(1-r^2).$$

Hence,

$$(4) \quad |zc''(z)/c'(z)| \leq 2r((1-\alpha)r + p)/(1-r^2).$$

The above result shows that

$$\operatorname{Re}\{zc''(z)/c'(z)\} \leq |zc''(z)/c'(z)| \leq 2r((1-\alpha)r + p)/(1-r^2).$$

But

$$\begin{aligned} \operatorname{Re}\{zc''(z)/c'(z)\} &= \operatorname{Re}\{(\partial/\partial \log z) \log c'(z)\} \\ &= r(\partial/\partial r) \log |c'(z)|, \end{aligned}$$

so

$$(5) \quad (\partial/\partial r) \log |c'(z)| \leq 2((1-\alpha)r + p)/(1-r^2).$$

On integrating (5) with respect to r , and then exponentiating we get (1). For the extremal function, see Remark 2.

THEOREM 2. *Let $c(z)$ be a function in $C(\alpha, p)$, $p > 0$. Then*

$$(6) \quad |c'(z)| \geq 1/(1 + 2pr/(1-\alpha) + r^2)^{1-\alpha},$$

for $z \in E$.

PROOF. From (2) it follows that

$$\begin{aligned} 1 + \operatorname{Re}\{h(z)\} &= \operatorname{Re}\{(1+K)/(1-K)\} = (1-K\bar{K})/|1-K|^2 \\ &\geq (1-|K|)/(1+|K|). \end{aligned}$$

Applying (3) to the above, we obtain

$$\operatorname{Re}\{h(z)\} \geq -2r((1-\alpha)r + p)/((1-\alpha)(1+r^2) + 2pr).$$

Hence

$$\begin{aligned} (1-\alpha) \operatorname{Re}\{h(z)\} &= \operatorname{Re}\{zc''/c'\} \\ &\geq -2(1-\alpha)r((1-\alpha)r + p) \div ((1-\alpha)(1+r^2) + 2pr). \end{aligned}$$

Thus

$$(\partial/\partial r) \log |c'| \geq -2((1-\alpha)r+p)/(1+2pr/(1-\alpha)+r^2).$$

Again, on integrating the above with respect to r and then exponentiating, we obtain (6). For the extremal function, see Remark 2.

THEOREM 3. Let $c(z)$ be a function in $C(\alpha, p)$, $p > 0$. Then

$$(7) \quad |c''(z)| \leq [2((1-\alpha)r+p)/(1-r^2)^{2-\alpha}][(1+r)/(1-r)]^p,$$

for $z \in E$.

PROOF. Inequality (7) follows readily from (1) and (4). For the extremal function, see Remark 2.

THEOREM 4. Let $s(z) = z + 2pz^2 + \dots$ be a function in $S(\alpha, p)$, $p > 0$. Then

$$(8) \quad \begin{aligned} (i) \quad & |s(z)| \leq (r/(1-r^2)^{1-\alpha})((1+r)/(1-r))^p \\ (ii) \quad & |s(z)| \geq r/(1+2pr/(1-\alpha)+r^2)^{1-\alpha}. \end{aligned}$$

Equality holds for part (i) for the function

$$s(z) = (z/(1-z^2)^{1-\alpha})((1+z)/(1-z))^p,$$

and equality holds for part (ii) for the function

$$s(z) = z/(1-2pz/(1-\alpha)+z^2)^{1-\alpha}.$$

PROOF. Set $c(z) = \int_0^z (s(t)/t)dt = z + pz^2 + \dots$. Then

$$(9) \quad zc'(z) = s(z).$$

From (9), it follows that $z(c''/c') + 1 = zs'/s$. Hence $s(z)$ is in $S(\alpha, p)$ if and only if $c(z)$ is in $C(\alpha, p)$. Now applying (1) to (9), we obtain (8-i), while on applying (6) to (9), we obtain (8-ii).

Now to show that the extremal functions are in $S(\alpha, p)$. For

$$s(z) = (z/(1-z^2)^{1-\alpha})((1+z)/(1-z))^p = z + 2pz + \dots,$$

we have

$$zs'(z)/s(z) = 1 + (p+\alpha-1)z/(1+z) + (p-\alpha+1)z/(1-z).$$

Since $p+\alpha-1 \leq 0$, $p-\alpha+1 \geq 0$, $\operatorname{Re}\{z/(1+z)\} < \frac{1}{2}$ and $\operatorname{Re}\{z/(1-z)\} > -\frac{1}{2}$ for z in the unit disc, it follows that

$$\operatorname{Re}\{zs'/s\} > 1 + \frac{1}{2}(p+\alpha-1) - \frac{1}{2}(p-\alpha+1) = \alpha.$$

For $s(z) = z/(1-2pz/(1-\alpha)+z^2)^{1-\alpha} = z + 2pz + \dots$, we have

$$zs'/s = (1-2b\alpha z + (2\alpha-1)z^2)/(1-2bz+z^2),$$

where $b = p/(1-\alpha) \leq 1$. The right-hand side of the above can be

written in the form, $2\alpha - 1 + 2(1 - \alpha)/(1 + z(z - b)/(1 - bz))$. We note that $|(z - b)/(1 - bz)| \leq 1$ and thus $\operatorname{Re}\{1/(1 + z(z - b)/(1 - bz))\} \geq \frac{1}{2}$. Hence $\operatorname{Re}\{zs'/s\} \geq 2\alpha - 1 + 2(1 - \alpha)\frac{1}{2} = \alpha$.

THEOREM 5. *Let $s(z)$ be a function in $S(\alpha, p)$, $p > 0$. Then*

$$(10) \quad |s'(z)| \leq [(1 - 2\alpha)r^2 + 2pr + 1]/(1 - r^2)^{2-\alpha} [(1 + r)/(1 - r)]^p.$$

The extremal function is that of Theorem 4, part (i).

PROOF. On differentiating (9), we obtain $s'(z) = zc''(z) + c'(z)$. Thus $|s'(z)| \leq r|c''| + |c'|$. Applying (1) and (7) to the above will yield (10).

3. The class $S_p(\alpha)$.

THEOREM 6. *Let $g(z) = z + d_2z^2 + \dots$ be a function in $S_p(\alpha)$, $p > 0$. Then*

$$(11) \quad \begin{aligned} \text{(i)} \quad & |g(z)| \leq [r/(1 - r^2)^{1-\alpha}] [(1 + r)/(1 - r)]^{p+1}, \\ \text{(ii)} \quad & |g(z)| \geq r(1 - r)/(1 + r)(1 + 2pr/(1 - \alpha) + r^2)^{1-\alpha}. \end{aligned}$$

The extremal function, for part (i), is

$$g(z) = [z/(1 - z^2)^{1-\alpha}] [(1 + z)/(1 - z)]^{p+1},$$

with $s(z) = [z/(1 - z^2)^{1-\alpha}] [(1 + z)/(1 - z)]^p$, while for part (ii), the extremal function is

$$g(z) = [z/(1 - 2pz/(1 - \alpha) + z^2)^{1-\alpha}] [(1 + z)/(1 - z)]$$

with $s(z) = z/(1 - 2pz/(1 - \alpha) + z^2)^{1-\alpha}$.

PROOF. Since $g(z)$ is in $S_p(\alpha)$, there exists an $s(z) \in S(\alpha, p)$ such that

$$(12) \quad g(z) = s(z)q(z),$$

where $q(z)$ is an analytic function whose real part is positive for $z \in E$, and such that $q(0) = 0$, $q'(0) = 1$. It is well known that

$$(13) \quad (1 - r)/(1 + r) \leq |q(z)| \leq (1 + r)/(1 - r),$$

see, e.g., [4, p. 169].

On applying (8) and (13) to (12) one gets (11).

Now to show that the extremal functions are in $S_p(\alpha)$. This follows easily from the fact that in both cases $s(z)$ is in $S(\alpha, p)$ as was seen at the end of the proof of Theorem 4, and $\operatorname{Re}\{g/s\} = \operatorname{Re}\{(1 + z)/(1 - z)\} > 0$, for z in the unit disc.

THEOREM 7. *Let $g(z)$ be a function in $S_p(\alpha)$, $p > 0$. Then $g(z)$ has radius of univalence and radius of starlikeness of at least $R_p(\alpha)$, where $R_p(\alpha)$ is the least positive root of*

$$(14) \quad (1 - \alpha)(2\alpha - 1)r^4 + 2(1 + p\alpha - \alpha)r^3 + 2((1 - \alpha)^2 + 2p)r^2 + 2(1 - p\alpha - \alpha)r - 1 + \alpha = 0.$$

PROOF. From (9), by differentiation and then equating the real parts, we get $\operatorname{Re}\{zs'/s\} = \operatorname{Re}\{z(c''/c') + 1\}$. We have shown in the proof of Theorem 2 that

$$\operatorname{Re}\{zc''/c'\} \geq -2(1 - \alpha)r((1 - \alpha)r + p)/((1 - \alpha)(1 + r^2) + 2pr).$$

So

$$(15) \quad \frac{\operatorname{Re}\{zs'/s\}}{\geq [(1 - \alpha)(2\alpha - 1)r^2 + 2pr\alpha + 1 - \alpha]/[(1 - \alpha)(1 + r^2) + 2pr]}.$$

MacGregor [3] shows $|q'/q| \leq 2/(1 - r^2)$ for functions analytic in the unit disc and of positive real part there. Hence we have

$$(16) \quad \operatorname{Re}\{zq'/q\} \geq -2r/(1 - r^2)$$

for such functions.

Since $g(z)$ is a p -close-to-star of order α , there exists an $s(z) \in S(\alpha, p)$ such that $g(z) = s(z)q(z)$, where $q(z)$ is a function that we have just described above. This gives

$$(17) \quad \operatorname{Re}\{zg'(z)/g(z)\} = \operatorname{Re}\{zs'(z)/s(z)\} + \operatorname{Re}\{zq'(z)/q(z)\}.$$

Now, applying (15) and (16) to (17), we have

$$\operatorname{Re}\{zg'/g\} \geq [(1 - \alpha)(2\alpha - 1)r^2 + 2p\alpha r + 1 - \alpha]/[(1 - \alpha)(1 + r^2) + 2pr] - 2r/(1 - r^2).$$

Set $\operatorname{Re}\{zg'/g\} = 0$, we obtain (14). Thus the radius of univalence and of starlikeness is at least as large as the smallest positive number satisfying (14). We note that, if $\alpha = 0$, then $R_p(0)$ is given by $r^4 - 2r^3 - 2(1 + 2p)r^2 - 2r + 1 = 0$, which is Chase's.

The author would like to thank the referee for his valuable suggestions.

REFERENCES

1. Warren E. Chase, p -close-to-star-functions, Notices Amer. Math. Soc. **17** (1970), 172. Abstract #672-315.
2. Mark Finkelstein, Growth estimates of convex functions, Proc. Amer. Math. Soc. **18** (1967), 412-418. MR **35** #5598.
3. T. H. MacGregor, Functions whose derivative has a positive real part, Trans. Amer. Math. Soc. **104** (1962), 532-537. MR **25** #4090.
4. Zeev Nehari, Conformal mapping, McGraw-Hill, New York, 1952. MR **13**, 640.

BOWLING GREEN STATE UNIVERSITY, BOWLING GREEN, OHIO 43402