IRREDUCIBLE LIE ALGEBRAS OF INFINITE TYPE

ROBERT LEE WILSON¹

ABSTRACT. Let V be a finite dimensional vector space over an algebraically closed field of characteristic $\neq 2, 3, 5$. It is shown that if $L \subseteq \mathfrak{gl}(V)$ is an irreducible Lie algebra of infinite type then either $L = \mathfrak{gl}(V)$, $L = \mathfrak{sl}(V)$, $\dim V = 2r \geq 4$ and $L = \mathfrak{sp}(V)$, $\dim V = 2r \geq 4$ and $L = \mathfrak{csp}(V)$, or there exists $A \in L$ such that $\text{ad } A \neq 0 = (\text{ad } A)^2$. As a corollary we obtain E. Cartan's classification of the irreducible Lie algebras of infinite type over $\mathbb{C}$.

Let L be a Lie algebra of linear transformations of a vector space V . For each nonnegative integer n the n th Cartan prolongation, L_n , is defined inductively by $L_0 = L$ and

$$L_n = \{ \phi \in \text{Hom}(V, L_{n-1}) \mid y(x\phi) = x(y\phi) \text{ for all } x, y \in V \}$$

for $n \geq 1$. If $L_n \neq (0)$ for all $n \geq 0$ then L is said to be of *infinite type*. The main result of this paper is:

THEOREM 1. *Let V be a finite dimensional vector space over an algebraically closed field of characteristic $\neq 2, 3, 5$. If L is an irreducible Lie algebra of infinite type then either $L = \mathfrak{gl}(V)$, $L = \mathfrak{sl}(V)$, $\dim V = 2r \geq 4$ and $L = \mathfrak{sp}(V)$, $\dim V = 2r \geq 4$ and $L = \mathfrak{csp}(V)$, or there exists an $A \in L$ such that $\text{ad } A \neq 0 = (\text{ad } A)^2$.*

Now it is easily seen (as in [7]) that if $(\text{ad } A)^2 = 0$ then A belongs to the radical of the Killing form. If $\Phi = \mathbb{C}$ and L is reductive this implies that $\text{ad } A = 0$. Furthermore, it is known (Theorem 1 of [4]) that an irreducible Lie algebra over $\mathbb{C}$ is of infinite type if and only if $L_2 \neq (0)$. Thus Theorem 1 implies the following theorem of E. Cartan [1]:

THEOREM 2. *Let V be a finite dimensional vector space over $\mathbb{C}$ and let L be an irreducible Lie algebra of linear transformations of V such that $L_2 \neq (0)$. Then either $L = \mathfrak{gl}(V)$, $L = \mathfrak{sl}(V)$, $\dim V = 2r \geq 4$ and $L = \mathfrak{sp}(V)$, or $\dim V = 2r \geq 4$ and $L = \mathfrak{csp}(V)$.*

Theorem 2, which is important in the study of primitive pseudogroups and infinite Lie algebras of Cartan type, has been proved by several authors ([1], [2], [3], [5], [9]). These proofs have involved considerable use of the classification and representation theory of

Received by the editors August 17, 1970.

AMS 1970 subject classifications. Primary 17B20; Secondary 17B65.

Key words and phrases. Cartan prolongation, infinite Lie algebra of Cartan type.

¹ The author has received partial support from a Sloan Foundation grant.

Copyright © 1971, American Mathematical Society

semisimple Lie algebras over $\mathbb{C}$ and hence cannot be generalized to fields of prime characteristic. In the proof presented here we use more elementary techniques which are valid over algebraically closed fields of characteristic $\neq 2, 3, 5$.

We will consider the following three conditions on a Lie algebra L of linear transformations of a vector space V :

CONDITION A. There exists $A \in L$ with $\text{rank } A = 1$ and $A^2 \neq 0$.

CONDITION B. Either $\dim V = 2$ or there exist $A, B \in L$ with $\text{rank } A = \text{rank } B = 1$, $\ker A = \ker B$, and $VA \neq VB$.

CONDITION C. There exists $A \in L$ with $\text{ad } A \neq 0 = (\text{ad } A)^2$.

By Condition $\sim X$ we will mean the negation of Condition X.

Theorem 1 is clearly a consequence of the following two lemmas.

LEMMA 1. *If Φ is an algebraically closed field of arbitrary characteristic, V is a finite dimensional vector space over Φ , and $L \subseteq \text{gl}(V)$ is a Lie algebra of infinite type then L contains a rank one transformation.*

LEMMA 2. *Let V be a finite dimensional vector space over a field Φ of characteristic $\neq 2, 3, 5$. Let L be an irreducible Lie algebra of linear transformations of V . Assume that L contains a rank one transformation. Then:*

- (i) *If Condition A holds $L = \text{gl}(V)$.*
- (ii) *If Conditions $\sim A$, B, and $\sim C$ hold then $L = \text{sl}(V)$.*
- (iii) *If Conditions $\sim A$, $\sim B$, and $\sim C$ hold then $\dim V = 2r \geq 4$ and $L = \text{sp}(V)$ or $L = \text{csp}(V)$.*

It is shown in [2] that if $\Phi = \mathbb{C}$ Lemma 1 is a consequence of Hilbert's Nullstellensatz. The proof given there is in fact independent of the assumption $\Phi = \mathbb{C}$ and could be used to prove our Lemma 1. We will give a somewhat more elementary proof here. Part (i) of Lemma 2 is proved in [6] for the case $\Phi = \mathbb{C}$.

In the proof of Lemma 1 we will need:

LEMMA 3. *Let Φ be an algebraically closed field of arbitrary characteristic. Let V and W be finite dimensional vector spaces over Φ with $2 \leq \dim V \leq \dim W$. Let T be a subspace of $\text{Hom}(V, W)$ such that $\dim T \geq \dim W$. Then there exist $\phi \in T$ and $v \in V$ such that $\phi \neq 0$, $v \neq 0$, and $v\phi = 0$.*

PROOF. Suppose $T \subseteq \text{Hom}(V, W)$ is a counterexample to the lemma such that $\dim W$ is minimal among all counterexamples. As T is a counterexample we have, for any nonzero $v \in V$, $\dim W \geq \dim vT = \dim T \geq \dim W$. Thus for any $w \in W$ there is a unique $\phi \in T$ such that $v\phi = w$. Then if v_1, v_2 are linearly independent elements of V we can find a basis $\{\psi_1, \dots, \psi_n\}$ of T such that

$$(1) \quad \begin{aligned} v_2\psi_i &= v_1\psi_{i+1} \quad \text{for all } 1 \leq i \leq n-1, \quad \text{and} \\ v_2\psi_n &= \sum_{i=1}^n v_1\psi_i a_i \quad \text{where the } a_i \in \Phi. \end{aligned}$$

(For if linearly independent elements $\psi_1, \dots, \psi_j$ satisfying (1) have been found for some $1 \leq j \leq n-1$ then the minimality of $\dim W$ implies that $v_2\psi_j \notin \langle v_1\psi_1, \dots, v_1\psi_j \rangle$. Thus there exists a unique $\psi_{j+1} \in T$ such that $\psi_1, \dots, \psi_{j+1}$ are linearly independent and satisfy (1). Proceeding by induction on j gives the result.) Now there exists $\lambda \in \Phi$ such that $\lambda^n - \sum_{i=0}^{n-1} \lambda^i a_{i+1} = 0$. Setting $b_k = \lambda^{n-k} - \sum_{i=0}^{n-k-1} \lambda^i a_{k+i+1}$ for $1 \leq k \leq n-1$ we see that $\psi = \psi_n + \sum_{i=1}^{n-1} \psi_i b_i \in T$ and $(v_2 - \lambda v_1)\psi = 0$. This contradicts the choice of T and proves the lemma.

PROOF OF LEMMA 1. For $\phi \in L_i$ and $j \leq i$ define

$$\text{im}_j(\phi) = \langle v_j(\dots(v_1\phi)\dots) \mid v_1, \dots, v_j \in V \rangle.$$

Now if $x \in \ker \phi$ and $y \in V$ then $0 = y(x\phi) = x(y\phi)$ so we have $\ker \phi \subseteq \ker(y\phi)$. Thus if $\psi \in \text{im}_j(\phi)$ we have $\ker \psi \supseteq \ker \phi$. Furthermore if $d_i = \min \{ \text{rank } \phi \mid 0 \neq \phi \in L_i \}$ we have $d_0 \leq d_1 \leq \dots \leq \dim V$. Hence there is some integer N such that $d_N = d_{N+i}$ for all $i \geq 0$. Now if $i \geq j \geq 0$ and if $\phi \in L_{N+i}$ satisfies $\text{rank } \phi = d_N$ then for $0 \neq \psi \in \text{im}_j(\phi)$ we have $\ker \psi \supseteq \ker \phi$ and $\text{rank } \psi \geq d_N = \text{rank } \phi$. Thus $\ker \psi = \ker \phi$. Thus $\text{im}_j(\phi) \subseteq \text{Hom}(V/\ker \phi, \text{im}_{j+1}(\phi))$ and $0 \neq \psi \in \text{im}_j(\phi)$ implies $\text{rank } \psi = d_N = \dim(V/\ker \phi)$. Thus if $d_N \geq 2$, Lemma 3 shows that $\dim \text{im}_j(\phi) \leq \dim \text{im}_{j+1}(\phi) - 1$. Hence $\dim L_{N-1} \geq \dim \text{im}_{i+1}(\phi) \geq \dim \text{im}_i(\phi) + i \geq i$ for all $i \geq 0$. Since L_{N-1} is finite dimensional this is impossible. Hence $1 = d_N = d_0$, proving the lemma.

PROOF OF LEMMA 2. Let $n = \dim V$. If $\{x_1, \dots, x_n\}$ and $\{y_1, \dots, y_n\}$ are bases for V we define elements E_{ij} and $F_{ij} \in \text{gl}(V)$ for $1 \leq i, j \leq n$ by $x_k E_{ij} = \delta_{ki} x_j$ and $y_k F_{ij} = \delta_{ki} y_j$. If $2k \leq n$ we define $\text{sp}(x_1, \dots, x_{2k})$ to be the Lie algebra of all $A \in \text{gl}(V)$ such that $VA \subseteq \langle x_1, \dots, x_{2k} \rangle$, $x_r A = 0$ for all $r > 2k$, and A is skew with respect to the skew-symmetric bilinear form defined by $(x_{2i+1}, x_j) = \delta_{j, 2i+2}$, $(x_{2i+2}, x_j) = -\delta_{j, 2i+1}$, and $(x_r, x_j) = 0$ for all $0 \leq i \leq k-1$, $1 \leq j \leq n$, $2k < r \leq n$.

We will presently verify the following statements about an irreducible Lie algebra L of linear transformations of V :

- (a) If $1 \leq k < n$ and $E_{1i} \in L$ for all $1 \leq i \leq k$ then there is a basis $\{y_1, \dots, y_n\}$ of V such that $F_{1i} \in L$ for all $1 \leq i \leq k+1$.
- (b) If $1 \leq k < n$, $E_{1i} \in L$ for all $1 \leq i \leq n$, and $E_{n1} \in L$ for all $1 \leq i \leq k$ then there is a basis $\{y_1, \dots, y_n\}$ of V such that $F_{1i} \in L$ for all $1 \leq i \leq n$ and $F_{n1} \in L$ for all $1 \leq i \leq k+1$.

(c) If $2 \leq k \leq m \leq n$, $E_{1i} \in L$ for all $2 \leq i \leq m$, $E_{i1} \in L$ for all $2 \leq i < k$, and L satisfies Condition $\sim C$ then there is a basis $\{y_1, \dots, y_n\}$ of V such that $F_{1i} \in L$ for all $2 \leq i \leq m$ and $F_{i1} \in L$ for all $2 \leq i \leq k$.

(d) If $3 \leq k < n$ and $E_{1i}, E_{i1} \in L$ for all $2 \leq i \leq k$ then there is a basis $\{y_1, \dots, y_n\}$ of V such that $F_{1i} \in L$ for all $2 \leq i \leq k+1$ and $F_{i1} \in L$ for all $2 \leq i \leq k$.

(e) If $\text{sp}(x_1, \dots, x_{2k}) \subseteq L$ where $\dim V > 2k \geq 2$ and L satisfies Conditions $\sim B$ and $\sim C$ then $\dim V \geq 2k+2$ and there is a basis $\{y_1, \dots, y_n\}$ of V such that $\text{sp}(y_1, \dots, y_{2k+2}) \subseteq L$.

(f) If $\text{sp}(V) \subseteq L$ and L satisfies Condition $\sim B$ then $L = \text{sp}(V)$ or $L = \text{csp}(V)$.

Lemma 2 follows immediately from statements (a)–(f). For if Condition A holds we may choose a basis $\{x_1, \dots, x_n\}$ for V such that $E_{11} \in L$. Then by (a), (b), and induction on k we see that $L = \text{gl}(V)$, proving (i). If Conditions $\sim A$, B, and $\sim C$ hold we may choose a basis $\{x_1, \dots, x_n\}$ of V such that $E_{12} \in L$ and if $n \geq 3$ we may also arrange that $E_{13} \in L$. Then using (c), (d), and induction on k we see that $L = \text{sl}(V)$, proving (ii). Finally if Conditions $\sim A$, $\sim B$, and $\sim C$ hold we may choose a basis $\{x_1, \dots, x_n\}$ of V such that $E_{12} \in L$. By (c) we may assume that $\text{sp}(x_1, x_2) \subseteq L$. Then by (e) and induction on k we have $\dim V = 2r \geq 4$ and $\text{sp}(V) \subseteq L$. Then (f) proves (iii).

We now verify (a)–(f). Throughout we will let $A = \sum E_{ij}a_{ij}$ where the $a_{ij} \in \Phi$.

(a): As $\langle x_1, \dots, x_k \rangle$ is not an invariant subspace there exists $A \in L$ such that $a_{ij} \neq 0$ for some $1 \leq i \leq k < j \leq n$. We may assume that $i = 1$ (replacing A by $A(\text{ad } E_{1i})$ if $i \neq 1$) and that $a_{rj} = 0$ whenever $r > 1$ or $j = 1$ (replacing A by $(A(\text{ad } E_{11}) - A)(\text{ad } E_{11})/2$). Letting $\{y_1, \dots, y_n\}$ be any basis for V satisfying $y_i = x_i$ for $1 \leq i \leq k$, $y_{k+1} = x_1A$, and $y_j \in \langle x_{k+1}, \dots, x_n \rangle$ for $j > k+1$ gives the result.

The proof of (b) is similar to that of (a).

(c): First assume that $\text{ad } E_{1k} = 0$. Then for every $A \in L$ we have $0 = A(\text{ad } E_{1k}) = \sum_i (E_{ik}a_{i1} - E_{1i}a_{ki})$. Hence $a_{ki} = 0$ for all $i \neq k$ so $\langle x_k \rangle$ is an invariant subspace, contradicting the irreducibility of L . Hence $\text{ad } E_{1k} \neq 0$ so by Condition $\sim C$ we have $(\text{ad } E_{1k})^2 \neq 0$. Since $A(\text{ad } E_{1k})^2 = E_{1k}(-2a_{k1})$ we have $E_{1k} \in L(\text{ad } E_{1k})^2$. Since $E_{1k}^2 = 0$ we have $(\text{ad } E_{1k})^3 = 0$. Thus Lemma V.8.2 of [8] shows that there exists $A \in L$ such that $A(\text{ad } E_{1k})^2 = 2E_{1k}$ and $A(\text{ad } E_{1k})(\text{ad } A) = -2A$. Now

$$A(\text{ad } E_{1k})(\text{ad } A) = \sum_{i,j} E_{ij} \left(2a_{kj}a_{i1} - \delta_{i1} \sum_r a_{kr}a_{rj} - \delta_{jk} \sum_r a_{ir}a_{r1} \right).$$

Thus we have $a_{k1} = -1$ and

$$(2) \quad a_{ij} = -a_{i1}a_{kj} + \left(\delta_{i1} \sum_r a_{kr}a_{rj} + \delta_{jk} \sum_r a_{ir}a_{r1} \right) / 2.$$

Setting $i=j=1$ gives $0 = \sum_r a_{kr}a_{r1}$. From this, using (2) to substitute for a_{rj} in $\sum_r a_{kr}a_{rj}$, we conclude that $\sum_r a_{kr}a_{rj} = 0$ for all $1 \leq j \leq n$ and similarly that $\sum_r a_{ir}a_{r1} = 0$ for all $1 \leq i \leq n$. Thus $a_{ij} = -a_{i1}a_{kj}$ and $\sum_r a_{ir}a_{rj} = 0$ for $1 \leq i, j \leq n$. Now set $y_1 = x_k A$, $y_k = x_k$, and $y_j = x_j + a_{j1}x_k$ for $j \neq 1, k$. Then $F_{k1} = A \in L$, $F_{1k} = -E_{1k} \in L$, and $F_{1i} = -E_{1k}a_{i1} - E_{1i} \in L$ for $2 \leq i \leq m$, $i \neq k$. Finally for $2 \leq i < k$ we have $F_{i1} = -E_{i1}(\text{ad } F_{1k})(\text{ad } F_{k1}) - (F_{11} - F_{kk})a_{ki} \in L$ as required.

(d): As in (a) we may find a basis $\{y_1, \dots, y_n\}$ of V such that $F_{1i}, F_{i1} \in L$ for $1 \leq i \leq k$ and $A = \sum F_{ij}b_{ij} \in L$ where $b_{ij} = \delta_{j,k+1}$. Then

$$F_{1,k+1} = A(\text{ad } F_{21})(\text{ad } F_{12})(\text{ad } F_{31})(\text{ad } F_{13}) + (F_{11} - F_{33})b_{22} \in L$$

as required.

(e): It is well known (for example [8, p. 67] where we take $x_{2i+1} = v_{i+1}$ and $x_{2i+2} = v_{r+i+1}$ for $0 \leq i \leq r-1$) that $\text{sp}(x_1, \dots, x_{2r})$ is spanned by the following elements and their transposes: $E_{2i+1,2i+1} - E_{2i+2,2i+2}$, $E_{2i+1,2i+2}$, $E_{2i+1,2j+1} - E_{2j+2,2i+2}$, and $E_{2i+1,2j+2} + E_{2j+1,2i+2}$ for $0 \leq i \neq j \leq r-1$. Thus it is easily checked that $\text{sp}(x_1, \dots, x_{2k+2})$ is generated by $\text{sp}(x_1, \dots, x_{2k})$, $E_{1,2k+1} - E_{2k+2,2}$, and $E_{2k+1,2k+2}$.

We first show that for some basis $\{y_1, \dots, y_n\}$ of V we have $\text{sp}(y_1, \dots, y_{2k}) \subseteq L$ and $A = F_{1,2k+1} + \sum_{i>2k} F_{i2}b_{i2} \in L$ where the $b_{i2} \in \Phi$. As $\langle x_1, \dots, x_{2k} \rangle$ is not an invariant subspace there exists $A \in L$ such that $a_{ij} \neq 0$ for some $1 \leq i \leq 2k < j \leq n$. We may assume that $i=1$ (replacing A by $A(\text{ad } E_{12})$ if $i=2$, by $A(\text{ad}(E_{1,2r+1} - E_{2r+2,2}))$ if $i=2r+1$, and by $A(\text{ad}(E_{1,2r+2} + E_{2r+1,2}))$ if $i=2r+2$), that $a_{rs} = 0$ unless $r=1$, $s>2$ or $r>2$, $s=2$ (replacing A by $A(\text{ad } E_{21})(\text{ad } E_{12}) - (E_{11} - E_{22})(a_{11} - a_{22}) - E_{12}(2a_{12})$), and that $a_{1s} = a_{s2} = 0$ for $s \leq 2k$ (replacing A by

$$A - \sum_{i=1}^{k-1} A((\text{ad } E_{2i+2,2i+1})(\text{ad } E_{2i+1,2i+2}) + (\text{ad } E_{2i+1,2i+2})(\text{ad } E_{2i+2,2i+1})).$$

Then letting $\{y_1, \dots, y_n\}$ be any basis for V satisfying $x_i = y_i$ for $1 \leq i \leq 2k$, $y_{2k+1} = x_1 A$, and $y_j \in \langle x_{2k+1}, \dots, x_n \rangle$ for $j > 2k+1$ gives the result.

Now assume that $\text{sp}(x_1, \dots, x_{2k}) \subseteq L$ and that

$$A = E_{1,2k+1} + \sum_{i>2k} E_{i2}a_{i2} \in L.$$

We will show that for some basis $\{y_1, \dots, y_n\}$ we have $\text{sp}(y_1, \dots, y_{2k}) \subseteq L$ and $F_{1,2k+1} - F_{2k+2,2} \in L$. If $a = a_{2k+1,2} \neq 0$ setting $y_i = x_i$ for $1 \leq i \leq 2k+1$ and $y_j = x_j - a^{-1}a_{j,2}x_{2k+1}$ for $j > 2k+1$ we see that $\text{sp}(y_1, \dots, y_{2k}) \subseteq L$ and $A = F_{1,2k+1} + F_{2k+1,2}a$. Then $F_{21}(\text{ad } A)^3 = (F_{1,2k+1} - F_{2k+1,2}a)3a \in L$ so $F_{1,2k+1} \in L$. This contradicts Condition $\sim B$ so we must have $a = 0$. Also by Condition $\sim B$ some $a_{i2} \neq 0$. Hence we can choose $y_{2k+2} \in \langle x_{2k+2}, \dots, x_n \rangle$ so that $y_{2k+2}A = -x_2$. Then setting $y_i = x_i$ for $1 \leq i \leq 2k+1$ and choosing $y_j \in \langle x_{2k+2}, \dots, x_n \rangle \cap \ker A$ for $j > 2k+2$ so that $\{y_1, \dots, y_n\}$ is a basis for V we have the result.

Now assume $\text{sp}(x_1, \dots, x_{2k}) \subseteq L$ and $E_{1,2k+1} - E_{2k+2,2} \in L$. We will show that for some basis $\{y_1, \dots, y_n\}$ of V we have $\text{sp}(y_1, \dots, y_{2k}) \subseteq L$ and $F_{1,2k+1} - F_{2k+2,2}, F_{2k+1,2k+2} \in L$, thus proving (e). We have $E_{2k+2,2k+1} = E_{21}(\text{ad}(E_{1,2k+1} - E_{2k+2,2}))^2/2 \in L$. Then as in (c) we have $B = \sum E_{ij}b_{ij} \in L$ where the $b_{ij} \in \Phi$ and satisfy $b_{2k+1,2k+2} = -1$, $b_{ij} = -b_{i,2k+2}b_{2k+1,j}$, and $\sum_r b_{ir}b_{rj} = 0$ for $1 \leq i, j \leq n$. Then if $y_{2k+2} = -x_{2k+1}B$, $y_{2k+1} = x_{2k+1}$, and $y_j = x_j + b_{j,2k+2}x_{2k+1}$ for $j \neq 2k+1, 2k+2$ we have $F_{2k+1,2k+2} = -B \in L$, $F_{2k+2,2k+1} = E_{2k+2,2k+1} \in L$, and $F_{1,2k+1} - F_{2k+2,2} = E_{1,2k+1} - E_{2k+2,2} - E_{2k+2,2k+1}(b_{2,2k+2} - b_{2k+1,1}) \in L$. Also, for $1 \leq i, j \leq 2k$, we have

$$F_{ij} = E_{ij} - E_{ij}(\text{ad } F_{2k+1,2k+2})(\text{ad } F_{2k+2,2k+1}) - F_{2k+2,2k+1}b_{ji}.$$

Hence $\text{sp}(y_1, \dots, y_{2k}) \subseteq L$ and $F_{1,2k+1} - F_{2k+2,2}, F_{2k+1,2k+2} \in L$ as required.

(f): If $\dim V = 2k$ and $\text{sp}(V) \subseteq L$ then for $0 \leq i \neq j \leq k-1$ if $A \in L$ then

$$\begin{aligned} E_{2i+1,2j+1}a_{2i+1,2j+1} + E_{2j+2,2i+2}a_{2j+2,2i+2} \\ = A(\text{ad } E_{2i+2,2i+1})(\text{ad } E_{2i+1,2i+2})(\text{ad } E_{2j+1,2j+2})(\text{ad } E_{2j+2,2j+1}) \in L. \end{aligned}$$

Then by Condition $\sim B$ we must have $a_{2i+1,2j+1} + a_{2j+2,2i+2} = 0$. Similarly we see that $a_{2i+1,2j+2} = a_{2j+1,2i+2}$ and $a_{2i+2,2j+1} = a_{2j+2,2i+1}$. Hence $A = D + S$ where $S \in \text{sp}(V)$ and $D = \text{diag}\{d_1, \dots, d_{2k}\} \in L$. Now

$$D(\text{ad}(E_{1,2i+1} - E_{2i+2,2})) = E_{1,2i+1}(d_1 - d_{2i+1}) - E_{2i+2,2}(d_{2i+2} - d_2) \in L.$$

Thus, again by Condition $\sim B$, we have $d_1 - d_{2i+1} = d_{2i+2} - d_2$ for all $0 \leq i \leq k-1$. Thus $2D = I(d_1 + d_2) + E$ where $E \in \text{sp}(V)$. Thus $L = \text{sp}(V)$ or $L = \text{csp}(V)$.

REFERENCES

1. E. Cartan, *Les groupes de transformations continus, infinis, simples*, Ann. Sci. École Norm. Sup. **26** (1909), 93-161; *Oeuvres complètes*, Vol. 2, Partie II, Gauthier-Villars, Paris, 1953, pp. 857-925.

2. V. Guillemin, D. Quillen and S. Sternberg, *The classification of the irreducible complex algebras of infinite type*, J. Analyse Math. **18** (1967), 107–112. MR **36** #221.
3. V. G. Kac, *Simple irreducible graded Lie algebras of finite growth*, Izv. Akad. Nauk SSSR Ser. Mat. **32** (1968), 1323–1367 = Math. USSR Izv. **2** (1968), 1271–1311.
4. S. Kobayashi and T. Nagano, *On filtered Lie algebras and geometric structures*. I, J. Math. Mech. **13** (1964), 875–907. MR **29** #5961.
5. ———, *On filtered Lie algebras and geometric structures*. III, J. Math. Mech. **14** (1965), 679–706. MR **32** #5803.
6. B. Kostant, *A characterization of the classical groups*, Duke Math. J. **25** (1958), 107–123. MR **20** #2381.
7. A. I. Kostrikin, *Simple Lie p -algebras*, Trudy Mat. Inst. Steklov. **64** (1961), 79–89; English transl., Amer. Math. Soc. Transl. (2) **55** (1966), 195–206. MR **24** #A1933.
8. G. B. Seligman, *Modular Lie algebras*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 40, Springer-Verlag, Berlin, 1967. MR **39** #6933.
9. I. M. Singer and S. Sternberg, *The infinite groups of Lie and Cartan*. I. *The transitive groups*, J. Analyse Math. **15** (1965), 1–114. MR **36** #911.

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, NEW YORK, NEW YORK 10012