A CLASS OF COMMUTATIVE BANACH ALGEBRAS WITH UNIQUE COMPLETE NORM TOPOLOGY AND CONTINUOUS DERIVATIONS

JOHN A. LINDBERG, JR.

ABSTRACT. Let A be a semisimple commutative complex algebra with identity and $\alpha(x)$ a monic polynomial over A . Two results are proved. If $B = A[x]/(\alpha(x))$ is a Banach algebra under some norm, then B has a unique complete norm topology. Furthermore, B has nontrivial derivations if and only if B has a nontrivial radical.

Introduction. It is known that if A is a commutative Banach algebra with identity and $\alpha(x)$ is a monic polynomial over A , then $B = A[x]/(\alpha(x))$ can be made into a Banach algebra with a norm extending the norm on A [1]. In [2], D. T. Brown has shown that if A is semisimple and regular (in the sense of Šilov), then B has unique complete norm topology. The assumption of regularity can be dropped as we show in §1. In general, the extension B is not semisimple; indeed, B has a nontrivial radical if and only if the discriminant of $\alpha(x)$ is zero or a zero divisor in A [5].

§2 is motivated by a recent result of R. J. Loy that states, in our context, that all derivations on B are continuous [4]. However, he does not show the existence of derivations on (nonsemisimple) B even though he gives their form. We consider the existence of nontrivial derivations in §2.

1. Uniqueness of complete norm topology on $A[x]/(\alpha(x))$. It is convenient to set first some of our notation and to make several comments. Lemma 1 of [2] states that if A is a semisimple algebra (all algebras in this paper are assumed to be over the complex field $\mathbb{C}$ and to possess an identity element) and if $B = A[x]/(\alpha(x))$ is a Banach algebra, then A is a Banach algebra under some norm. Actually, it is proved that if $K(A)$ denotes the intersection of the kernels of nontrivial complex homomorphisms, then $A/K(A)$ is a Banach algebra. However, it is known that each maximal ideal M in A is contained in a maximal ideal M_0 in B such that $M = M_0 \cap A$ (see p. 259 of [7]). Since B is a Banach algebra, M_0 is the kernel of a com-

Received by the editors June 26, 1970.

AMS 1969 subject classifications. Primary 4655, 4650; Secondary 1360.

Key words and phrases. Uniqueness of complete norm topology, continuous derivations, normed extensions, commutative Banach algebras.

Copyright © 1971, American Mathematical Society

plex homomorphism; hence, so is M . Thus, $K(A) = (0)$ since A is semisimple and Brown's Lemma 1 follows.

For a complex algebra A , let Φ_A denote the space of nontrivial complex homomorphisms on A , together with the Gelfand topology (see Chapter III of [6]). If $p(x) \in A[x]$, let $p(x)$ denote the coset $p(x) + (\alpha(x))$, and if $\phi \in \Phi_A$, let $p_\phi(x)$ denote $\sum \phi(p_i)x^i$, $p(x) = \sum p_i x^i$. The space Φ_B is identifiable with $\{(\phi, \lambda) \in \Phi_A \times C : \alpha_\phi(\lambda) = 0\}$. The action of $(\phi, \lambda) \in \Phi_B$ on $p(x)$ is given by $(\phi, \lambda)(p(x)) = p_\phi(\lambda)$.

Finally, before proceeding to the theorem of this section, we discuss a recent result of B. E. Johnson which will be used in our proof. In the setting of a Banach algebra, he proved in [3] that if d is a derivation of the Banach algebra A into the bounded functions on Φ_A , then the point derivations d_ϕ defined by $\phi \circ d$ are continuous on A for all but at most a finite number of ϕ 's. Now, suppose that at each point $\phi \in \Phi_A$ we have a finite number of point derivations, but no more than n , some positive number. Let them be denoted by $d_\phi^1, \dots, d_\phi^{k_\phi}$, $k_\phi \leq n$, $\phi \in \Phi_A$. If $d_\phi^i(a)$ is bounded for all i , ϕ and $a \in A$, then by at most n applications of Johnson's result, we have that for all i and for all except possibly a finite number of ϕ 's, d_ϕ^i is continuous on A .

THEOREM 1. *Let A be a semisimple complex algebra and $\alpha(x)$ a monic polynomial of degree n over A . If $B = A[x]/(\alpha(x))$ is a Banach algebra with norm $\|\cdot\|$, then A is $\|\cdot\|$ -closed in B . Consequently, B has unique complete norm topology.*

PROOF. By the comments preceding the statement of the theorem, we can conclude that each $\phi \in \Phi_A$ is $\|\cdot\|$ -continuous on A , where $\|\cdot\|$ is the norm on B . Thus, $\Phi_{\bar{A}} = \Phi_A$, $\bar{A}$ the $\|\cdot\|$ -closure of A in B .

Suppose $\{a_n\}$ is a sequence in A converging to $p(x) \in B$ with respect to $\|\cdot\|$. Then $\phi(a_n) = (\phi, \lambda)(a_n) \rightarrow (\phi, \lambda)(p(x)) = \phi(p(\lambda))$ for all $(\phi, \lambda) \in \Phi_B$. This implies that there exists an $a \in A$ such that $\phi(a_n) \rightarrow \phi(a)$ for all $\phi \in \Phi_A$. The existence of such an a is shown in the proof of Theorem 3 in [2]. Hence, $p(x)$ can be uniquely expressed as the sum of an element in A and an element in the radical $R(B)$ of B , namely, $p(x) = a + [p(x) - a]$ and $p(x) - a \in \bar{A} \cap R(B)$.

We next show that there is a finite subset F of Φ_B such that if $p(x) \in \bar{A}$, $\partial p \leq n-1$, then $\phi(p^{(j-1)}(\lambda)) = 0$ for j satisfying $2 \leq j \leq M(\phi, \lambda)$ and for all $(\phi, \lambda) \in \Phi_B \setminus F$, where $M(\phi, \lambda)$ denotes the multiplicity of λ as a root of $\alpha_\phi(x) = 0$. For $j \geq 1$, $p^{(j)}(x)$ denotes the j th derivative of $p(x)$ and $p^{(0)}(x) = p(x)$. Consider the statement: there exists a finite subset $F_i \subset \Phi_B$ such that $2 \leq j \leq \min(i, M(\phi, \lambda))$ implies $\phi(p^{(j-1)}(\lambda)) = 0$ for all $p(x) \in \bar{A}$ with $\partial p \leq n-1$ and for all $(\phi, \lambda) \in \Phi_B \setminus F_i$. Suppose that

the statement is true for i . We proceed to show that it is true for $i+1$ as well. Define $d_{(\phi, \lambda)}$ for $(\phi, \lambda) \in \Phi_B$ as follows:

$$\begin{aligned} d_{(\phi, \lambda)}(p(\mathbf{x})) &= 0 && \text{if } M(\phi, \lambda) \leq i \text{ or } (\phi, \lambda) \in F_i, \\ &= \phi(p^{(i)}(\lambda)) && \text{otherwise,} \end{aligned}$$

where $p(\mathbf{x}) \in \overline{A}$ and $\partial p \leq n-1$. Clearly, for each $p(\mathbf{x}) \in \overline{A}$ with $\partial p \leq n-1$,

$$\sup_{(\phi, \lambda) \in \Phi_B} |d_{(\phi, \lambda)}(p(\mathbf{x}))| < +\infty,$$

and $d_{(\phi, \lambda)}$ is a linear functional on $\overline{A}$. To show that $d_{(\phi, \lambda)}$ is a point derivation at ϕ on $\overline{A}$, let $p(\mathbf{x}), q(\mathbf{x}) \in \overline{A}$, $\partial p \leq n-1$, $\partial q \leq n-1$ and write $p(\mathbf{x})q(\mathbf{x}) = Q(\mathbf{x})\alpha(\mathbf{x}) + r(\mathbf{x})$, where $Q(\mathbf{x}), r(\mathbf{x}) \in A[\mathbf{x}]$ and $\partial r \leq n-1$. If $M(\phi, \lambda) \leq i$ or $(\phi, \lambda) \in F_i$, $d_{(\phi, \lambda)} = 0$ and there is nothing to prove. Suppose, therefore, $M(\phi, \lambda) \geq i+1$ and $(\phi, \lambda) \notin F_i$. Then $d_{(\phi, \lambda)}(p(\mathbf{x})q(\mathbf{x})) = \phi(r^{(i)}(\lambda))$. But

$$r^{(i)}(x) = \sum_{j=0}^i \binom{i}{j} \{p^{(i-j)}(x)q^{(j)}(x) - Q^{(i-j)}(x)\alpha^{(j)}(x)\}.$$

Thus,

$$\begin{aligned} d_{(\phi, \lambda)}(p(\mathbf{x})q(\mathbf{x})) &= \sum_{j=0}^i \binom{i}{j} \{ \phi(p^{(i-j)}(\lambda))\phi(q^{(j)}(\lambda)) - \phi(Q^{(i-j)}(\lambda))\phi(\alpha^{(j)}(\lambda)) \} \\ &= \phi(p^{(i)}(\lambda))\phi(q(\lambda)) + \phi(p(\lambda))\phi(q^{(i)}(\lambda)) \\ &= d_{(\phi, \lambda)}(p(\mathbf{x}))(\phi, \lambda)(q(\mathbf{x})) + (\phi, \lambda)(p(\mathbf{x}))d_{(\phi, \lambda)}(q(\mathbf{x})), \end{aligned}$$

since $\phi(\alpha^{(j)}(\lambda)) = 0$ for $j = 0, 1, 2, \dots, i$ and $p(\mathbf{x}), q(\mathbf{x}) \in \overline{A}$, $\partial p \leq n-1$, $\partial q \leq n-1$, $(\phi, \lambda) \notin F_i$ imply that $\phi(p^{(j-1)}(\lambda)) = \phi(q^{(j-1)}(\lambda)) = 0$ for $j = 2, \dots, \min(i, M(\phi, \lambda)) = i$. Hence, $d_{(\phi, \lambda)}$ is a point derivation. By the extension of Johnson's result mentioned above, we can conclude that there is a finite set F_{i+1} such that $d_{(\phi, \lambda)}$ is continuous whenever $(\phi, \lambda) \in \Phi_B \setminus F_{i+1}$. We may and do assume that $F_{i+1} \supset F_i$. Now, if

$$a_n \xrightarrow{\|\cdot\|} p(\mathbf{x}), \quad \{a_n\} \subset A,$$

then $d_{(\phi, \lambda)}(a_n) = 0$ for all n so that $(\phi, \lambda)(p^{(i)}(\mathbf{x})) = 0$ whenever $M(\phi, \lambda) \geq i+1$ and $(\phi, \lambda) \notin F_{i+1}$. Thus $2 \leq j \leq \min(i+1, M(\phi, \lambda))$ implies that $\phi(p^{(j-1)}(\lambda)) = 0$ whenever $(\phi, \lambda) \notin F_{i+1}$. Since the argument also shows that the statement is true for $i=2$, we have by induction that $\phi(p^{(j-1)}(\lambda)) = 0$ whenever $2 \leq j \leq M(\phi, \lambda)$, except possibly for (ϕ, λ) belonging to a finite subset F of Φ_B . (For the case $i=2$, $d_{(\phi, \lambda)}(p(\mathbf{x}))$ is simply $\phi(p^{(1)}(\lambda))$ when $2 \leq M(\phi, \lambda)$ and zero otherwise.)

Let $F' = (\pi)^{-1}(\pi(F))$, and let $p(x) \in \overline{A} \cap R(B)$, where $\pi(\phi, \lambda) = \phi$ for $(\phi, \lambda) \in \Phi_B$. Then $(\phi, \lambda) \in \Phi_B$ implies that $\phi(p(\lambda)) = 0$. We may assume that the degree ∂p of $p(x)$ is smaller than n . Now, suppose that $(\phi, \lambda) \notin F'$ and that $p_\phi(x)$ is not the zero polynomial. Then λ is a root of $p_\phi(x) = 0$ of at least multiplicity $M(\phi, \lambda)$. But this implies that $p_\phi(x) = 0$ has at least $n = \sum_{(\phi, \lambda) \in (\pi)^{-1}(\phi)} M(\phi, \lambda)$ roots, each repeated according to multiplicity. This is a contradiction since $\partial p \leq n-1$. Hence $(\phi, \lambda) \in F'$ implies that $p_\phi(x)$ is the zero polynomial. Thus, if $p(x) = \sum_{k=0}^{n-1} p_k x^k \in R(B) \cap \overline{A}$, then each $\hat{p}_k$ vanishes on Φ_A , except possibly for a finite subset. If $\hat{p}_k(\phi) \neq 0$, then ϕ is an isolated point of Φ_A .

We now continue as in [2]. If ϕ is an isolated point in Φ_A , then there exists an idempotent u_ϕ in A such that $u_\phi(\theta) = 1$ if and only if $\theta = \phi$ (see [6, p. 168]), since A is a Banach algebra under some norm (see [2, Lemma 1]). Since A is semisimple $u_\phi a = \phi(a)u_\phi$, $a \in A$, and therefore $u_\phi A$ is closed in B . Now, if $p(x) = p_0 + \cdots + p_{n-1}x^{n-1} \in B$, then $u_\phi p(x) = \phi(p_0)u_\phi + \phi(p_1)u_\phi x + \cdots + \phi(p_{n-1})u_\phi x^{n-1}$. Furthermore if

$$a_n \xrightarrow{\|\cdot\|} p(x),$$

then $u_\phi a_n \rightarrow u_\phi p(x)$ and $u_\phi p(x)$ must lie in $u_\phi A$. Hence, $u_\phi p_k = 0$, $1 \leq k \leq n-1$, or equivalently, $\phi(p_k) = 0$ for $1 \leq k \leq n-1$. Thus, $p(x) \in \overline{A}$ implies that $p(x) = p_0 \in A$ and $A = \overline{A}$ since A is semisimple.

The second assertion follows from Theorem 2 of [2].

2. Derivations on $A[x]/(\alpha(x))$. In our context, the work of Loy in [4] shows that if A is a semisimple Banach algebra, then any derivation D on $B = A[x]/(\alpha(x))$ must be of the form $D(p(x)) = p'(x)D(x)$, $p(x) = p_0 + p_1x + \cdots + p_{n-1}x^{n-1}$, $n = \partial\alpha$. This particularly simple form reflects the fact that A has no nontrivial derivations into itself [3]. It is easily established that any linear transformation on B of the form $p(x) \rightarrow p'(x)p_0(x)$, $p_0(x)$ fixed, is a derivation on B if and only if $\alpha'(x)p_0(x) = 0$. Note that $p(x) \rightarrow p'(x)p_0(x)$ is continuous with respect to any Banach algebra norm on B , by virtue of Theorem 1.

THEOREM 2. *Let A be a semisimple commutative Banach algebra and $\alpha(x) \in A[x]$ a monic polynomial. Then B has a nontrivial derivation if and only if the radical of B is nontrivial.*

PROOF. The only if part follows from Theorem 4 of [3].

Suppose, now, that $R(B) \neq (0)$. By Corollary 9.3 of [5], we have that the discriminant d_α of $\alpha(x)$ is either zero or a zero divisor in A , say $d_\alpha c = 0$, $c \neq 0$. From Lemma 9.1 of [5], we have the existence of

polynomials $\gamma(x)$, $\delta(x)$, $R(x) \in A[x]$ and nonzero elements $a, b \in A$ such that

- (i) $a\alpha(x) = \gamma(x)R(x)$,
- (ii) $b\alpha'(x) = \delta(x)R(x)$,
- (iii) $\phi(a) = 0$ if and only if $\phi(b) = 0$, and $\phi(c) = 0$ implies $\phi(a) = 0$, and
- (iv) if $\beta_\phi(x) \mid \alpha_\phi(x)$, $\beta_\phi(x) \mid \alpha'_\phi(x)$, then $\beta_\phi(x) \mid R_\phi(x)$, where $\phi \in \Phi_A$.

Set $p_0(x) = ab\gamma(x)$, and $D(p(x)) = p'(x)p_0(x)$. To show that D is a derivation, we need only verify that $\alpha'(x)p_0(x) = 0$, that is, $\alpha'(x)p_0(x) \in (\alpha(x))$. By conditions (i) and (ii), we have that

$$p_0(x)\alpha'(x) = ab\gamma(x)\alpha'(x) = a\gamma(x)\delta(x)R(x) = a^2\delta(x)\alpha(x).$$

We next show that D is nontrivial.

Suppose that $p_0(x) = q(x)\alpha(x)$. Then $q(x)\alpha'(x)\alpha(x) = a^2\delta(x)\alpha(x)$. Since $\alpha(x)$ is monic and hence not a zero divisor in $A[x]$, $q(x)\alpha'(x) = a^2\delta(x)$. Now, $a^2b\alpha'(x) = a^2\delta(x)R(x) = q(x)\alpha'(x)R(x)$. Hence, $a^2b = q(x)R(x)$. If $\phi \in \Phi_A$ and $\phi(a) \neq 0$, then $q_\phi(x)R_\phi(x) = \phi(a^2b) \neq 0$ by (iii), and consequently, $q_\phi(x)$ and $R_\phi(x)$ are constant polynomials. On the other hand, if $\phi \in \Phi_A$ and $\phi(a) \neq 0$, then $\phi(c) \neq 0$. Therefore, $\phi(d_a) = 0$ so that there exists a complex number λ such that $(x - \lambda) \mid \alpha_\phi(x)$ and $(x - \lambda) \mid \alpha'_\phi(x)$. By (iv), $(x - \lambda) \mid R_\phi(x)$ so that $\partial R_\phi(x) > 0$, a contradiction. It follows that $p_0(x) \notin (\alpha(x))$, and D as defined is not trivial. This completes the proof.

In conclusion, we note that since the results from [5] used above are valid for semisimple complex algebras, Theorem 2 is true for semisimple complex algebras A without nontrivial derivations.

REFERENCES

1. R. Arens and K. Hoffman, *Algebraic extension of normed algebras*, Proc. Amer. Math. Soc. **7** (1956), 203–210. MR **17**, 1113.
2. D. T. Brown, *A class of Banach algebras with a unique norm topology*, Proc. Amer. Math. Soc. **17** (1966) 1429–1434. MR **34** #4938.
3. B. E. Johnson, *Continuity of derivations on commutative algebras*, Amer. J. Math. **91** (1969), 1–10. MR **39** #7433.
4. R. J. Loy, *Uniqueness of the complete norm topology and continuity of derivations on Banach algebras*, Tôhoku Math. J. **22** (1970), 371–378.
5. J. A. Lindberg, *Algebraic extensions of commutative Banach algebras*, Pacific J. Math. **14** (1964), 559–583. MR **30** #3380.
6. C. E. Rickart, *General theory of Banach algebras*, University Series in Higher Math., Van Nostrand, Princeton, N. J., 1960. MR **22** #5903.
7. O. Zariski and P. Samuel, *Commutative algebra*. Vol. I, University Series in Higher Math., Van Nostrand, Princeton, N. J., 1958. MR **19**, 833.

SYRACUSE UNIVERSITY, SYRACUSE, NEW YORK 13210