A WHITEHEAD TYPE THEOREM

H. B. HASLAM¹

ABSTRACT. Let $\mathfrak{F}$ denote the Serre class of finite abelian groups. We consider, for example, conditions under which a map which induces an $\mathfrak{F}$ -epimorphism in homotopy also induces an $\mathfrak{F}$ -epimorphism in homology.

1. Introduction. Let $f: X \rightarrow Y$ be a map, $\mathfrak{C}$ a Serre class of abelian groups. Modulo some technical assumptions on the spaces X and Y or the class $\mathfrak{C}$, the Whitehead theorem states that f induces a $\mathfrak{C}$ -isomorphism in homotopy in each dimension if and only if it induces a $\mathfrak{C}$ -isomorphism in homology in each dimension. We are concerned here with finding conditions under which the word "isomorphism" can be replaced by "epimorphism" or "monomorphism" for the class $\mathfrak{F}$ of finite abelian groups. For example, we show that if a map g from an H -space Y to a 1-connected finite CW-complex X induces an $\mathfrak{F}$ -epimorphism in homotopy, then it induces an $\mathfrak{F}$ -epimorphism in homology and X is an H -space mod $\mathfrak{F}$. As a special case we recover a result of [4]: If X is a 1-connected finite CW-complex and a G -space mod $\mathfrak{F}$ (i.e. the evaluation map $\omega: (X^{\mathfrak{F}}, 1) \rightarrow (X, *)$ induces an $\mathfrak{F}$ -epimorphism in homotopy), then X is an H -space mod $\mathfrak{F}$. (The converse is also true.) Moreover, ω induces an $\mathfrak{F}$ -epimorphism in homology. The proof given here is much simpler than that given in [4].

I wish to thank W. Browder for a conversation which motivated this note.

All spaces are assumed to have the based homotopy type of a CW-complex and all maps and homotopies are to preserve base points. We will frequently not distinguish between a map and its homotopy class. The symbol " $\mathfrak{h}$ " will be used to denote Hurewicz homomorphisms. We assume that the reader is familiar with [1] and [2].

2. The result. Let X be a 1-connected finite CW-complex, Y be a 1-connected space with $H_m(Y)$ finitely generated for all m and let $f: X \rightarrow Y, g: Y \rightarrow X$ be maps.

THEOREM 1. *Suppose that Y is an H -space.*

Received by the editors August 18, 1970.

AMS 1969 subject classifications. Primary 5540.

Key words and phrases. H -space (mod $\mathfrak{F}$), H' -space (mod $\mathfrak{F}$).

¹ Supported by NSF Grant GP 19964.

Copyright © 1971, American Mathematical Society

(i) If $f_*: \pi_m(X) \rightarrow \pi_m(Y)$ is an $\mathfrak{F}$ -monomorphism for all m , then so is $f_*: H_m(X) \rightarrow H_m(Y)$.

(ii) If $g_*: \pi_m(Y) \rightarrow \pi_m(X)$ is an $\mathfrak{F}$ -epimorphism for all m , then so is $g_*: H_m(Y) \rightarrow H_m(X)$. Moreover, X , in each case, is an H -space mod $\mathfrak{F}$.

THEOREM 2. Suppose that Y is an H' -space.

(i) If $f_*: H_m(X) \rightarrow H_m(Y)$ is an $\mathfrak{F}$ -monomorphism for all m , then so is $f_*: \pi_m(X) \rightarrow \pi_m(Y)$.

(ii) If $g_*: H_m(Y) \rightarrow H_m(X)$ is an $\mathfrak{F}$ -epimorphism for all m , then so is $g_*: \pi_m(Y) \rightarrow \pi_m(X)$. Moreover, X , in each case, is an H' -space mod $\mathfrak{F}$.

REMARK. The 1-connectedness assumption on Y is needed only for Theorem 2 (ii) and neither assumption on Y is needed for Theorem 1 (ii).

We will need the following result, the proof of which depends only on the universal coefficient theorem and the representability of cohomology.

LEMMA 1. Let B be a space for which $H_m(B)$ is finitely generated and let $\beta \in \pi_m(B)$. Then $\mathfrak{h}(\beta) \neq 0$ if and only if there is a map $h: B \rightarrow K(\pi, m)$ (=an Eilenberg-Mac Lane space) such that $h\beta$ is not homotopic to a constant. (We may take $\pi = \mathbb{Z}_p$ or $\mathbb{Z}$ depending on whether $\mathfrak{h}(\beta)$ has finite or infinite order.)

LEMMA 2. Let $a: A \rightarrow B$ be a map from an H -space A to a finite CW-complex B . If $a_*: \pi_{2n}(A) \rightarrow \pi_{2n}(B)$ is an $\mathfrak{F}$ -epimorphism, then $\mathfrak{h}(\pi_{2n}(B)) \in \mathfrak{F}$.

PROOF. It suffices to show that $\mathfrak{h}(\beta)$ has finite order for each $\beta \in \pi_{2n}(B)$. In order to obtain a contradiction, assume that there is a $\beta \in \pi_{2n}(B)$ such that $\mathfrak{h}(\beta)$ has infinite order. By Lemma 1, there is a map $h: B \rightarrow K(\mathbb{Z}, 2n)$ such that $h\beta$ is not homotopic to a constant. Since $a_*: \pi_{2n}(A) \rightarrow \pi_{2n}(B)$ is an $\mathfrak{F}$ -epimorphism there is an $\alpha \in \pi_{2n}(A)$ such that $a_*(\alpha) = r\beta$ where r is some nonzero integer. Now, if $\rho: \Omega\Sigma A \rightarrow A$ is a retraction map (A is an H -space), then $h\rho \circ \Omega\Sigma\alpha: \Omega\Sigma S^{2n} \rightarrow K(\mathbb{Z}, 2n)$ is a nontrivial map which factors through a finite complex. This is clearly impossible (consider the ring structure of $H^*(\Omega\Sigma S^{2n})$) and the lemma is proved.

COROLLARY 1. If B is $(2n-1)$ -connected, then $\pi_{2n}(B) \in \mathfrak{F}$.

PROOF OF THEOREM 1. (i) Since the homotopy suspension homomorphism for an H -space is a monomorphism in all dimensions, it follows that the suspension homomorphism $i_*: \pi_m(X) \rightarrow \pi_m(\Omega\Sigma X)$ (i is the inclusion map) is an $\mathfrak{F}$ -monomorphism for all m and therefore that X is an H -space mod $\mathfrak{F}$. Let $h: S \rightarrow X$ be a weak $\mathfrak{F}$ -equivalence,

where S is a finite product of odd dimensional spheres. Since the Hurewicz homomorphism $\mathfrak{h}:\pi_m(Y)\rightarrow H_m(Y)$ is an $\mathfrak{F}$ -monomorphism it follows from Lemma 1 that fh induces an $\mathfrak{F}$ -epimorphism in cohomology and hence an $\mathfrak{F}$ -monomorphism in homology. Thus f induces an $\mathfrak{F}$ -monomorphism in homology.

(ii) We first show, by induction, that $\pi_{2n}(X)\in\mathfrak{F}$ for all n . For $n=1$, this follows from Corollary 1. Assume that $\pi_{2n}(X)\in\mathfrak{F}$ for $2n<N$, N odd. Since $g_*:\pi_m(Y)\rightarrow\pi_m(X)$ is an $\mathfrak{F}$ -epimorphism for all m , we can use the multiplication on Y to obtain a map $h_N:S\rightarrow Y$ such that gh_N induces an $\mathfrak{F}$ -isomorphism in homotopy in dimensions $\leq N$, where S is a finite product of odd dimensional spheres S^{n_i} , $3\leq n_i\leq N$. We can assume that gh_N is an inclusion map. Then $\pi_m(X, S)\in\mathfrak{F}$ for all $m\leq N$; by the Hurewicz theorem, $\mathfrak{h}:\pi_{N+1}(X, S)\rightarrow H_{N+1}(X, S)$ is an $\mathfrak{F}$ -isomorphism. Since $\pi_{N+1}(S)\in\mathfrak{F}$, it follows that $\mathfrak{h}:\pi_{N+1}(X)\rightarrow H_{N+1}(X)$ is an $\mathfrak{F}$ -monomorphism and so, by Lemma 2, $\pi_{N+1}(X)\in\mathfrak{F}$. Thus $\pi_{2n}(X)\in\mathfrak{F}$ for all n . It is now a simple matter to show that for $N\geq\dim X$, gh_N is a weak $\mathfrak{F}$ -equivalence. Therefore X is an H -space mod $\mathfrak{F}$ and $g_*:H_m(Y)\rightarrow H_m(X)$ is an $\mathfrak{F}$ -epimorphism for all m .

PROOF OF THEOREM 2. (i) Since $f_*:H_m(X)\rightarrow H_m(Y)$ is an $\mathfrak{F}$ -monomorphism for all m , $f^*:H^m(Y)\rightarrow H^m(X)$ is an $\mathfrak{F}$ -epimorphism for all m . Let $\{\beta_i\}$ be a basis for the free part of $H^*(X)$ and let $\{\gamma_i\}\subset H^*(Y)$ be chosen so that $f^*(\gamma_i)=t_i\beta_i$ for some nonzero integer t_i . Let $r>\dim X$ be arbitrary, $\gamma'_i=\gamma_i|Y^r$, where Y^r is the r -skeleton of Y .

Since $\gamma_i\cup\gamma_i=0$ (Y is an H' -space), $\gamma'_i\cup\gamma'_i=0$ and [3] there is a map $h_i:Y^r\rightarrow S^{n_i}$, $n_i=\dim \gamma_i$, which maps the fundamental class of S^{n_i} to some nonzero multiple of γ'_i . Making use of the fact that Y is an H' -space we obtain a map $h:Y^r\rightarrow \bigvee S^{n_i}$ (as in [3]) such that hf is a weak $\mathfrak{F}$ -equivalence (by the cellular approximation theorem we can assume $f(X)\subset Y^r$). Therefore X is an H' -space mod $\mathfrak{F}$ and $f_*:\pi_m(X)\rightarrow\pi_m(Y)$ is an $\mathfrak{F}$ -monomorphism for $m<r$. Since r was arbitrary the result follows.

(ii) Since the Hurewicz homomorphism for an H' -space is an $\mathfrak{F}$ -epimorphism in all dimensions, it follows that $\mathfrak{h}:\pi_m(X)\rightarrow H_m(X)$ is an $\mathfrak{F}$ -epimorphism for all m and hence that X is an H' -space mod $\mathfrak{F}$. Moreover, it is clear that there is a map $h:\bigvee S^{n_i}\rightarrow Y$ such that fh is a weak $\mathfrak{F}$ -equivalence and the result follows.

REMARK. In contrast to the Whitehead theorem, the converse of each assertion in Theorems 1 and 2 is false. Counterexamples are given as follows:

1(i). The inclusion map $S^{2n}\rightarrow\Omega\Sigma S^{2n}$.

1(ii). The quotient map $S^n\times S^n\rightarrow S^n\wedge S^n=S^{2n}$, $n=3$ or 7 .

2(i). The Whitehead product map $S^{m+n-1} \rightarrow S^m \vee S^n$, $m+n$ even, $m, n \geq 2$.

2(ii). The inclusion map $S^m \vee S^n \rightarrow S^m \times S^n$.

BIBLIOGRAPHY

1. M. Arkowitz and C. R. Curjel, *The Hurewicz homomorphism and finite homotopy invariants*, Trans. Amer. Math. Soc. 110 (1964), 538–551. MR 28 #1624.
2. ———, *Zum Begriff des H -Raumes mod $\mathfrak{F}$* , Arch. Math. 16 (1965), 186–190. MR 31 #4031.
3. Israel Berstein, *Homotopy mod $\mathfrak{C}$ of spaces of category 2*, Comment. Math. Helv. 35 (1961), 9–14. MR 23 #A652.
4. H. B. Haslam, *G -spaces mod $\mathfrak{F}$ and H -spaces mod $\mathfrak{F}$* , Duke Math. J. (to appear).

FLORIDA STATE UNIVERSITY, TALLAHASSEE, FLORIDA 32306