THE CLASS NUMBER OF $Q(\sqrt{-p})$, FOR $p \equiv 1 \pmod{8}$ A PRIME

EZRA BROWN

ABSTRACT. Let $h(-p)$ be the class number of the quadratic field $Q(\sqrt{-p})$, where $p \equiv 1 \pmod{8}$ is a prime. Write $p = a^2 + b^2 = 2e^2 - d^2$, where $a \equiv e \equiv d \equiv b + 1 \equiv 1 \pmod{2}$ and $e > 0$. We prove that $h(-p) \equiv 0$ or $4 \pmod{8}$ according as $(e|p) = 1$ or -1 ; using this, we prove that $h(-p) \equiv (p-1)/2 + b \pmod{8}$. The proofs are elementary, relying on the theory of composition of binary quadratic forms.

1. Introduction. Let $h(-d)$ denote the class number of the algebraic number field $Q(\sqrt{-d})$. Recently, Barrucand and Cohn (see [1]) and Hasse (see [2] and [3]) have studied congruence conditions on $h(-p)$ and $h(-2p)$ modulo 8, where p is an odd prime. In this paper we present some new results related to this problem; in particular, we prove the following theorem.

THEOREM 1. *Let $p \equiv 1 \pmod{8}$ be a prime, and write $p = a^2 + b^2$, where a is odd and b is even. Then $h(-p) \equiv (p-1)/2 + b \pmod{8}$.*

2. Preliminary definitions and results. Our study uses the Gaussian theory of composition of binary quadratic forms; we begin by reviewing some standard definitions and results. For details and proofs, see [4].

DEFINITIONS. A binary quadratic form $[a, b, c] = ax^2 + bxy + cy^2$ is called *primitive* if $(a, b, c) = 1$ and *ambiguous* if $a|b$. An *ambiguous class* of forms is one which contains an ambiguous form. The *principal form* of discriminant d is $f_1 = [1, \varepsilon, (\varepsilon-d)/4]$, where $\varepsilon = 0$ or $1 \equiv d \pmod{4}$; the class (genus) containing f_1 is called the *principal class* (genus). The primitive classes of discriminant d form a group under Gaussian composition, called the *class group* (if $d < 0$, only the positive-definite classes are considered). The symbol $(p|q)_4 = 1$ or -1 according as p is or is not a biquadratic residue $\pmod{q}$, and $(p|q)$ is the Legendre quadratic residue symbol.

PROPOSITION. *If A and B are primitive classes of discriminant d , there exist forms $[a_1, b, a_2c]$ in A and $[a_2, b, a_1c]$ in B . The product (composite)*

Received by the editors March 29, 1971.

AMS 1970 subject classifications. Primary 12A25, 12A50; Secondary 10A15, 10C05.

Key words and phrases. Class number, quadratic residues, quadratic fields, algebraic number fields, composition of binary quadratic forms.

© American Mathematical Society 1972

class AB of A and B is the class containing $[a_1 a_2, b, c]$. The ambiguous classes are precisely those classes A such that $A^2=I$, the principal class; every class in the principal genus is a square, and each genus contains the same number of classes. If $p \equiv 1 \pmod{8}$ is a prime, $h(-p)$ is the number of positive-definite classes of discriminant $-4p$ (=determinant p).

3. The main results. Proof of Theorem 1 uses the following.

THEOREM 2. Let $p \equiv 1 \pmod{8}$ be a prime. Write $p = a^2 + b^2 = 2e^2 - d^2$, where $a \equiv d \equiv e \equiv b + 1 \equiv 1 \pmod{2}$ and $e > 0$. Then $h(-p) \equiv 0$ or $4 \pmod{8}$ according as $(e|p) = 1$ or -1 .

PROOF. If $p = 2e^2 - d^2$, then the form $f = [e, 2d, 2e]$ has determinant p . Denote by A the class of f , by B the class of $g = [2, 2, (p+1)/2]$, and by I the class of $f_1 = [1, 0, p]$. The forms of determinant p have generic characters $(m|p)$ and $(-1|m)$, and are divided into two genera: the two ambiguous classes are I and B . Since $(2|p) = (-1|(p+1)/2) = 1$, B is in the principal genus.

If $(e|p) = 1$, then A is in the principal genus; by the proposition, $A = C^2$ for some class C . Now $A^2 = B$, since $[e, 2d, 2e]^2 = [e^2, 2d, 2]$, which is equivalent to g . Since A^3 contains $[2, 2d, e^2] \cdot [e, 2d, 2e] = [2e, 2d, e]$, $A^3 = A^{-1}$. Hence, A is an element of order 4 in the class group G . Since C^2 has order 4, C has order 8 (this is easy to check). Hence G has an element of order 8, so we have that $h(-p) = \text{ord } G \equiv 0 \pmod{8}$.

If $(e|p) = -1$, then f is in the nonprincipal genus, and so A is not the square of any class (for, if $A = C^2$, the generic characters of A would be the squares of those of C , and A would be in the principal genus). By the above, A has order 4. Let D be a class whose order is 2^n . If $n=0$, $D=I$; if $n=1$, $D=B$; if $n=2$, $D^2=B$, so $(DA)^2 = B^2 = I$, and hence $D=A$ or A^{-1} . If $n \geq 3$, then either $A = (D^{2^{n-1}})^2$ or $A = ((D^{-1})^{2^{n-1}})^2$, contrary to our assumption that A is in the nonprincipal genus, and thus not a square. Hence, the only elements of G of order 2^n are A , A^{-1} , B and I . Let H be the subgroup of G generated by A ; we show that G/H has odd order. If there is a class K whose square is in H , then $K^2 = A$, A^2 , A^3 or I . In these cases, we see that either A is in the principal genus, $K=A$ or A^{-1} , A is in the principal genus, or $A=B$. None of these are possible by previous reasoning; hence G/H has odd order, and we have $h(-p) = \text{ord } G = \text{ord } H \cdot \text{ord } (G/H) = 4 \text{ ord } (G/H) \equiv 4 \pmod{8}$. Q.E.D.

PROOF OF THEOREM 1. Let $p = a^2 + b^2 = 2e^2 - d^2$, where $a \equiv d \equiv e \equiv b + 1 \equiv 1 \pmod{2}$. From the congruences $2e^2 \equiv d^2 \pmod{p}$ and $2e^2 \equiv p \pmod{d}$, we find that $(e|p) = (2|p)_4 (d|p) = (2|p)_4 (p|d) = (2|p)_4 (2|d)$. Since e is odd, $2e^2 \equiv 2 \pmod{16}$; hence $p \equiv 2 - d^2 \pmod{16}$. Thus, $p \equiv 1$ or $9 \pmod{16}$ according as $(2|d) = 1$ or -1 . Then $(e|p) = (2|p)_4$ or $-(2|p)_4$ according as $p \equiv 1$ or $9 \pmod{16}$. It is well known (see [5]) that if $p \equiv 1$

$(\text{mod } 8)$ is a prime represented by $a^2 + b^2$ with b even, then $(2|p)_4 = (-1)^{b/4}$. Now, if $p \equiv 1 \pmod{16}$, then $(e|p) = 1$ if and only if $b \equiv 0 \pmod{8}$; if $p \equiv 9 \pmod{16}$, then $(e|p) = 1$ if and only if $b \equiv 4 \pmod{8}$. In each case, $(e|p) = 1$ or -1 according as $(p-1)/2 + b \equiv 0$ or $4 \pmod{8}$. Hence, by Theorem 2, we have that $h(-p) \equiv (p-1)/2 + b \pmod{8}$. Q.E.D.

REFERENCES

1. P. Barrucand and H. Cohn, *Note on primes of type $x^2 + 32y^2$, class number, and residuacity*, J. Reine Angew. Math. **238** (1969), 67–70. MR **40** #2641.
2. H. Hasse, *Über die Klassenzahl des Körpers $P(\sqrt{-p})$ mit einer Primzahl $p \equiv 1 \pmod{2^3}$* , Aequationes Math. **3** (1969), 165–169. MR **40** #2642.
3. ———, *Über die Klassenzahl des Körpers $P(\sqrt{-2p})$ mit einer Primzahl $p \neq 2$* , J. Number Theory **1** (1969), 231–234. MR **39** #1424.
4. B. W. Jones, *The arithmetic theory of quadratic forms*, Carus Monograph Series, no. 10, Math. Assoc. of America, Buffalo, New York, 1950. MR **12**, 244.
5. A. L. Whiteman, *The sixteenth power residue character of 2*, Canad. J. Math. **6** (1954), 364–373. MR **16**, 14.

DEPARTMENT OF MATHEMATICS, VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY, BLACKSBURG, VIRGINIA 24061