A FLAG TRANSITIVE PLANE OF ORDER 49

M. L. NARAYANA RAO

ABSTRACT. A non-Desarguesian translation plane of order 49, whose collineation group is transitive on the ideal points, is constructed.

1. Introduction. Among the known non-Desarguesian translation (projective) planes, there are only four planes (besides Luneburg planes), whose collineation groups are transitive on the ideal points. They are: (i) the translation plane coordinatized by the near field of order 9, (ii) the Hering's plane of order 27 [3, p. 236], (iii) the two flag transitive planes of Foulser [4] of order 25. The aim of this paper is to construct a translation plane of order 49 and show that its collineation group is transitive on the ideal points.

2. Let F be the set of all ordered pairs (a, b) over $\text{GF}(7)$. Let C be a set of 2×2 matrices over $\text{GF}(7)$ satisfying

- (i) C contains $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ and $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.
(2.1) (ii) C contains 49 matrices.
(iii) If M, N are from C and $|M - N| = 0$, then $M = N$, where $|X|$ denotes the determinant of the matrix X .

The conditions (2.1) imply that corresponding to each ordered pair (a, b) in F , there is exactly one matrix of the form $\begin{bmatrix} a & b \\ p & q \end{bmatrix}$. The matrix $\begin{bmatrix} a & b \\ p & q \end{bmatrix}$ corresponding to the ordered pair (a, b) is denoted by $M(a, b)$. We now define addition "+" and multiplication "·" on F as follows:

$$\begin{aligned} (a, b) + (c, d) &= (a + c, b + d), \\ 2.2) \quad (a, b) \cdot (c, d) &= (ca + dp, cb + dq) = (c, d)M(a, b) \\ &\quad \text{where } \begin{bmatrix} a & b \\ p & q \end{bmatrix} \text{ is the matrix } M(a, b) \text{ in } C. \end{aligned}$$

THEOREM 2.1. *The set F with operations $+$ and $\cdot$ defined by (2.2) is a left Veblen-Wedderburn system.*

PROOF. See [3, §5] or [2, §5].

Received by the editors October 5, 1970.

AMS 1969 subject classifications. Primary 5070; Secondary 5060.

Key words and phrases. Veblen-Wedderburn systems, projective planes, affine planes, flag transitivity.

© American Mathematical Society 1972

TABLE 3.1

i	L_i	$A+L_iP$	$L_iT=$ L_{i+1}	i	L_i	$A+L_iP$	$L_iT=$ L_{i+1}	i	L_i	$A+L_iP$	$L_iT=$ L_{i+1}
1	0 0 0 0	1 0 1 4	1 0 0 1	17	1 1 3 6	6 5 2 2	2 4 2 5	34	4 4 0 4	0 6 1 0	1 4 6 0
2	1 0 0 1	6 6 1 3	5 5 1 2	18	2 4 2 5	4 1 4 4	2 2 0 6	35	1 4 6 0	6 2 3 5	1 2 1 1
3	5 5 1 2	5 4 6 1	2 6 3 3	19	2 2 0 6	4 3 1 5	3 3 1 3	36	1 2 1 1	6 4 6 2	6 6 0 5
4	2 6 3 3	4 6 2 5	6 4 2 2	20	3 3 1 3	2 1 6 0	6 3 3 1	37	6 6 0 5	3 2 1 6	2 3 1 6
5	6 4 2 2	3 4 4 0	2 1 4 1	21	6 3 3 1	3 5 2 0	4 2 2 3	38	2 3 1 6	4 2 6 4	4 5 3 4
6	2 1 4 1	4 4 0 6	3 4 6 3	22	4 2 2 3	0 1 4 6	4 1 1 0	39	4 5 3 4	0 5 2 4	5 2 3 0
7	3 4 6 3	2 0 3 2	4 0 5 2	23	4 1 1 0	0 2 6 3	4 2 4 0	40	5 2 3 0	5 0 2 1	3 0 2 4
8	4 0 5 2	0 3 5 4	6 5 5 0	24	4 3 4 0	0 0 0 0	—	41	3 0 2 4	2 4 4 5	1 5 5 1
9	6 5 5 0	3 3 5 6	1 3 4 4	25	2 0 5 3	4 5 5 3	5 1 6 2	42	1 5 5 1	6 1 5 5	1 6 2 6
10	1 3 4 4	6 3 0 3	0 4 5 6	26	5 1 6 2	5 1 3 3	6 2 6 4	43	1 6 2 6	6 0 4 3	6 0 4 6
11	0 4 5 6	1 3 5 0	3 5 4 3	27	6 2 6 4	3 6 3 1	5 3 0 2	44	6 0 4 6	3 1 0 1	0 1 1 4
12	3 5 4 3	2 6 0 4	5 4 2 1	28	5 3 0 2	5 6 1 2	6 1 1 5	45	0 1 1 4	1 6 6 6	0 5 6 5
13	5 4 2 1	5 5 4 1	4 6 6 1	29	6 1 1 5	3 0 6 5	5 0 4 5	46	0 5 6 5	1 2 3 0	5 6 5 4
14	4 6 6 1	0 4 3 4	0 6 2 0	30	5 0 4 5	5 2 0 2	0 2 4 2	47	5 6 5 4	5 3 5 2	3 1 0 3
15	0 6 2 0	1 1 4 2	3 2 5 5	31	0 2 4 2	1 5 0 5	0 3 3 5	48	3 1 0 3	2 3 1 1	2 5 6 6
16	3 2 5 5	2 2 5 1	1 1 3 6	32	0 3 3 5	1 4 2 3	3 6 3 2	49	2 5 6 6	4 0 3 6	2 0 5 3
				33	3 6 3 2	2 5 2 6	4 4 0 4				

3. We now construct a set C of matrices over $\text{GF}(7)$ and show that C satisfies the conditions (2.1). Let $A = \begin{bmatrix} 1 & 0 \\ 1 & 4 \end{bmatrix}$, $P = \begin{bmatrix} 5 & 6 \\ 0 & 6 \end{bmatrix}$ be 2×2 matrices over $\text{GF}(7)$. Let D be the set of matrices containing $\begin{bmatrix} 3 & 4 \\ 3 & 4 \end{bmatrix}$ and all 2×2 matrices L over $\text{GF}(7)$ such that $|A + LP| \neq 0$. Let T be a mapping on D defined by

$$(3.1) \quad LT = (A + LP)^{-1}A, \quad \text{for } L \in D.$$

Let $L_1 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ and $L_{25} = \begin{bmatrix} 2 & 0 \\ 5 & 3 \end{bmatrix}$ be two elements from D . Let $L_{i+1} = L_i T$ for $1 \leq i \leq 23$, $25 \leq i < 49$. A reference to Table 3.1 shows that the matrices L_i for $1 \leq i \leq 49$ are all defined. In the table any entry of the form $\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}$ stands for the matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$. The square brackets $[\]$ are dropped for convenience.

LEMMA 3.1. *The set $C = \{L_i | 1 \leq i \leq 49\}$ satisfies the conditions (2.1).*

PROOF. The conditions (i) and (ii) of (2.1) are obviously satisfied. An examination of Table 3.1 shows that

$$(3.2) \quad \begin{aligned} (a) \quad & |L_i| \neq 0, \quad 1 < i \leq 49, \\ (b) \quad & |L_{25} - L_i| \neq 0, \quad 1 \leq i \leq 24, 26 \leq i \leq 49. \end{aligned}$$

Let X and Y be two matrices from D such that $XT = U$ and $YT = V$ with $|U| \neq 0 \neq |V|$. Then it follows from the relation $(X - Y)P = AU^{-1}(V - U)V^{-1}$ that

$$(3.3) \quad |V - U| = 0 \quad \text{if and only if} \quad |X - Y| = 0.$$

To prove the condition (iii) of (2.1) we use induction. From (3.2) we have $|L_1 - L_i| \neq 0$ for $1 < i < 24$ since L_1 is the zero matrix. Suppose $|L_k - L_i| \neq 0$ for $k < i \leq 24$. Choosing $X = L_k$ and $Y = L_i$ in (3.3) and using (3.2) we obtain $|L_{k+1} - L_i| \neq 0$ for $k + 1 < i \leq 24$. Thus we have $|L_i - L_j| \neq 0$ for $i < j$, $1 \leq i < 24$, $2 \leq j \leq 24$. Since we are dealing with 2×2 matrices we have $|L_i - L_j| = |L_j - L_i| \neq 0$. It now follows that

$$(3.4) \quad |L_i - L_j| \neq 0, \quad i \neq j, 1 \leq i \leq 24, 1 \leq j \leq 24.$$

Similarly by choosing X and Y of (3.3) appropriately and using induction, we obtain

$$(3.5) \quad |L_i - L_j| \neq 0, \quad i \neq j, 25 \leq i \leq 49, 25 \leq j \leq 49,$$

$$(3.6) \quad |L_i - L_j| \neq 0, \quad 1 \leq i \leq 24, 25 \leq j \leq 49.$$

From the inequalities (3.4), (3.5) and (3.6) we obtain that if X and Y are from C and $X \neq Y$, then $|X - Y| \neq 0$. Thus the condition (iii) of (2.1) is proved. Hence the lemma.

4. Let Π be the projective plane coordinatized by $F(+, \cdot)$ [5, p. 353], [6, p. 45], [7, p. 31], i.e. Π has points (c) , (a, d) , (∞) , and lines $[k]$,

$[m, b], [\infty] = L_\infty$ for $a, b, c, d, k, m \in F$ and $\infty \notin F$. Incidence in Π is defined by $(x, y) \in [m, b]$, if $y = m \cdot x + b$ and $(x, y) \in [k]$, if $x = k$. The plane Π may also be considered as a four dimensional right vector space V over $\text{GF}(7)$, the points of Π being ordered quadruples over $\text{GF}(7)$ and the lines being two dimensional subspaces of V . The line corresponding to the equation $y = m \cdot x$, $m \in F$, is given by the subspace

$$(4.1) \quad V_m = \{(a, b, c, d) \mid (a, b) \in F, (c, d) = (p, q)M(m), \\ \text{where } M(m) \text{ is the matrix from } C \text{ corresponding to } m\}.$$

The line $x=0$ corresponds to the subspace

$$(4.2) \quad V_\infty = \{(0, 0, c, d) \mid (c, d) \in F\}.$$

The lines $y = m \cdot x + b$ correspond to the appropriate translates of V_m for $m \in F$ or $m = \infty$. The group G_0 of all collineations fixing $(0, 0)$ of Π consists of all linear transformations of V which permute the subspaces V_m for $m \in F$ or $m = \infty$ among themselves [1, Satz 19, p. 178], [7, p. 208].

Let R be a nonsingular linear transformation partitioned as $\begin{bmatrix} A & B \\ P & Q \end{bmatrix}$ where A, B, P, Q are 2×2 matrices over $\text{GF}(7)$. Let V_m be the two dimensional subspace corresponding to the line $y = m \cdot x$. The subspace V_m is transformed by R on to a subspace U_m of V given by

$$(4.3) \quad U_m = \{(a, b, c, d) \mid (a, b) = (p, q)A + (p, q)MP; \\ (c, d) = (p, q)B + (p, q)MQ; (p, q) \in F; M = M(m)\}$$

and V_∞ is transformed on to the subspace U_∞ given by

$$(4.4) \quad U_\infty = \{(a, b, c, d) \mid (a, b) = (p, q)P, (c, d) = (p, q)Q, (p, q) \in F\}.$$

The following lemma gives a set of sufficient conditions so that a nonsingular linear transformation R on V is a collineation of Π belonging to G_0 .

LEMMA 4.1. *A nonsingular transformation $R = \begin{bmatrix} A & B \\ P & Q \end{bmatrix}$ on V where A, B, P, Q are 2×2 matrices is a collineation of Π belonging to G_0 , if the following two conditions are satisfied:*

- (i) *either $P = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ and $|Q| \neq 0$ or $|P| \neq 0$ and $P^{-1}Q \in C$,*
- (ii) *for each $M \in C$, either $A + MP = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ and $|B + MQ| \neq 0$ or $|A + MP| \neq 0$ and there exists a unique matrix N in C such that $(A + MP)^{-1}(B + MQ) = N$.*

PROOF. At the outset it may be observed that if $P = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, then $A \neq \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, for, otherwise, R becomes singular. Further there exists at most one matrix $M \in C$ such that $A + MP = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$. Suppose $H \in C$ and $A + HP = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$. Then we have $M - H = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$. The theorem is proved, if we show that R leaves the set $\{V_m \mid m = \infty, \text{ or } m \in F\}$ of subspaces of V invariant. Here two cases arise.

Case (i). Let $m = \infty$. If $P = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, then

$$\begin{aligned} U_m &= \{(0, 0, c, d) \mid (c, d) = (a, b)Q; (a, b) \in F\} \\ &= \{(0, 0, c, d) \mid (c, d) \in F\} = V_\infty, \quad \text{since } |Q| \neq 0. \end{aligned}$$

If $|P| \neq 0$, then

$$\begin{aligned} U_\infty &= \{(a, b, c, d) \mid (a, b) \in F, (c, d) = (a, b)P^{-1}Q\} = V_r, \\ &\quad \text{where } P^{-1}Q = M(r). \end{aligned}$$

Case (ii). Let $m \in F$. If $A + MP = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ where $M = M(m)$, then

$$\begin{aligned} U_m &= \{(0, 0, a, b) \mid (a, b) = (c, d)(B + MQ); (c, d) \in F; |B + MP| \neq 0\} \\ &= \{(0, 0, a, b) \mid (a, b) \in F\} = V_\infty. \end{aligned}$$

If $|A + MP| \neq 0$, then

$$\begin{aligned} U_m &= \{(a, b, c, d) \mid (a, b) \in F, (c, d) = (a, b)(A + MP)^{-1}(B + MQ)\} \\ &\quad \text{where } M(r) = (A + MP)^{-1}(B + MQ). \end{aligned}$$

Thus the members of the set $\{V_m \mid m = \infty, m \in F\}$ are permuted among themselves by the mapping R . Incidentally it follows that the distinguished point (r) is mapped on to the distinguished point (m) where $V_m R = V_r$. This completes the proof of the lemma.

5. Let

$$R = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 1 & 4 & 1 & 4 \\ 5 & 6 & 0 & 0 \\ 0 & 6 & 0 & 0 \end{bmatrix} \quad \text{and} \quad S = \begin{bmatrix} 0 & 3 & 3 & 5 \\ 5 & 0 & 0 & 5 \\ 4 & 5 & 0 & 6 \\ 2 & 5 & 2 & 1 \end{bmatrix}$$

be two nonsingular linear transformations on V . The matrix R may be partitioned as $R = \begin{bmatrix} A & B \\ P & Q \end{bmatrix}$ where $A = \begin{bmatrix} 1 & 0 \\ 1 & 4 \end{bmatrix} = B$, $P = \begin{bmatrix} 5 & 6 \\ 0 & 0 \end{bmatrix}$ and $Q = \begin{bmatrix} 1 & 0 \\ 1 & 4 \end{bmatrix}$. Similarly S may be partitioned as $S = \begin{bmatrix} A & B \\ P & Q \end{bmatrix}$ where $A = \begin{bmatrix} 0 & 3 \\ 4 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 3 & 5 \\ 0 & 5 \end{bmatrix}$, $P = \begin{bmatrix} 5 & 0 \\ 2 & 5 \end{bmatrix}$ and $Q = \begin{bmatrix} 0 & 3 \\ 4 & 5 \end{bmatrix}$. In order to show that S is a collineation belonging to G_0 we need Table 5.1. In Table 5.1, A, B, C, D are the matrices obtained by partitioning S and $MW = (A + MP)^{-1}(B + MQ)$. Any entry of the form $\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}$ stands for the matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$.

It may be noticed that the linear transformations R and S satisfy the conditions of Lemma 4.1 by referring to Tables 3.1 and 5.1. Consequently R and S are collineations of Π belonging to G_0 and the restrictions of R and S to the points of L_∞ induce two permutations f and g respectively

TABLE 5.1

$M \in C$	$A+MP$	$B+MQ$	MW	$M \in C$	$A+MP$	$B+MQ$	MW	$M \in C$	$A+MP$	$B+MQ$	MW
0 0	0 3	3 5	0 1	0 3	6 4	2 1	4 0	0 5	3 0	6 3	2 1
0 0	5 0	0 5	1 4	3 5	6 5	0 3	5 2	6 5	4 6	4 3	4 1
0 1	2 1	5 6	4 3	4 0	2 2	6 5	6 0	2 1	3 4	3 6	5 0
1 4	3 4	0 2	4 0	5 2	1 0	6 0	4 6	4 1	2 4	5 6	4 5
4 3	1 3	5 1	0 2	6 0	3 5	4 5	1 1	5 0	6 0	5 5	2 2
4 0	0 6	3 5	4 2	4 6	5 1	1 4	3 6	4 5	3 3	6 3	0 6
0 2	4 6	0 0	0 0	1 1	6 6	4 6	0 3	2 2	5 2	5 0	0 5
4 2	4 2	0 0	0 0	3 6	1 3	2 4	3 5	0 6	3 2	5 4	6 5
1 0	4 1	2 5	6 2	0 4	1 2	4 2	5 2	0 6	5 5	1 4	1 6
0 1	0 5	2 6	6 4	5 6	2 6	0 4	3 0	2 0	6 3	5 5	2 6
6 2	0 1	1 0	4 1	5 2	3 3	2 0	4 6	1 6	2 3	0 4	1 3
6 4	2 1	2 2	1 0	3 0	3 1	4 5	6 1	2 6	4 5	3 4	4 4
4 1	4 0	1 6	2 5	4 6	0 4	4 4	1 2	1 3	3 2	1 1	3 6
1 0	2 5	6 5	6 6	6 1	3 0	3 6	1 1	4 4	1 5	4 2	3 2
2 5	4 3	4 3	1 0	1 2	1 4	6 0	0 4	3 6	3 6	5 4	0 6
6 6	6 4	6 4	0 1	1 1	4 3	1 6	5 6	3 2	0 4	1 0	2 0
1 4	5 0	3 2	2 6	1 5	0 5	5 3	5 5	2 0	1 6	1 5	6 5
6 0	1 2	1 5	3 3	5 1	6 2	4 6	1 2	5 3	3 5	1 1	5 0
2 6	6 1	6 4	3 0	5 5	2 4	1 3	4 4	6 5	6 2	0 3	5 1
3 3	2 2	3 1	2 4	1 2	6 1	3 0	0 4	5 0	4 4	2 5	6 2
3 0	5 4	0 5	6 3	4 4	3 1	0 2	4 2	5 1	1 5	0 6	3 2
2 4	0 2	6 2	3 1	0 4	6 6	1 2	2 3	6 2	5 5	5 0	5 5
6 3	2 6	3 1	1 4	4 2	6 5	3 0	1 5	3 2	2 0	4 0	2 0
3 1	5 6	6 6	6 0	2 3	5 4	4 1	5 1	5 5	0 1	5 3	5 3
2 4	2 5	2 2	4 5	3 1	0 2	2 6	3 3	3 5	1 1	3 3	5 6
2 5	2 0	1 3	3 4	0 3	4 1	6 1	1 3	4 3	6 0	2 1	5 4
4 5	5 6	2 3	3 4	3 3	4 5	6 1	5 3	5 6	4 2	3 4	5 4
3 4	4 0	5 2	6 3	1 3	1 6	5 1	0 2	5 4	5 3	3 2	2 1
3 4	6 3	1 2	6 6	5 3	5 1	4 1	6 4	5 4	0 6	6 2	6 1
6 3	0 3	0 1	0 5	0 2	2 3	4 0	2 2	2 1	1 1	0 6	1 5
6 6	1 0	2 4	2 4	6 4	4 4	5 2	3 1	6 1	5 3	6 6	3 5
0 5	1 4	3 3	2 5	2 2	3 6	2 0	0 3	1 5	5 2	2 3	4 3
				2 3	0 0	0 1	—				
				1 6	0 0	4 4					

on the points of L_∞ given by

$$\begin{aligned} f: & (\infty, 1, 2, \dots, 24)(25, 26, \dots, 49), \\ g: & (2, 27, 23, 25)(45, 24, 31, 1)(32, 8, 44, 17)(11, 40, 14, 12)(46, 6, 30, 19) \\ & (15, 43, 10, 13)(35, 4, 41, 21)(42, 3, 34, 22)(25, 9, 26, 16)(18, 39, 7, 37) \\ & (48, 30, 28, 5)(12, 47, 13, 29)(\infty, 38), \end{aligned}$$

where ∞ is the distinguished point (∞) and i is the distinguished point corresponding to the slope (a_i, b_i) the first row of L_i , $1 \leq i \leq 49$.

It now follows that $\langle f, g \rangle$ is transitive on the points of L_∞ and consequently the collineation group of Π is flag transitive on Π^* , the affine plane obtained from Π by deleting the distinguished line. Since the set C is not closed under the matrix addition, $F(+, \cdot)$ is not a field [3, p. 220] and consequently Π is not Desarguesian.

The full collineation group of this plane and Foulser's flag transitive planes will be discussed elsewhere.

REFERENCES

1. J. André, *Über nicht-Desarguessche Ebenen mit transitiver Translationsgruppe*, Math. Z. **60** (1954), 156–186. MR **16**, 64.
2. R. H. Bruck and R. C. Bose, *The construction of translation planes from projective spaces*, J. Algebra **1** (1964), 85–102. MR **28** #4414.
3. P. Dembowski, *Finite geometries*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 44, Springer-Verlag, New York, 1968. MR **38** #1597.
4. D. A. Foulser, *Solvable flag transitive affine groups*, Math. Z. **86** (1964), 191–204. MR **30** #1190.
5. M. Hall, Jr., *The theory of groups*, Macmillan, New York, 1959. MR **21** #1996.
6. D. R. Hughes, *Review of some results in collineation groups*, Proc. Sympos. Pure Math., vol. 1, Amer. Math. Soc., Providence, R.I., 1959, pp. 42–55. MR **22** #7053.
7. G. Pickert, *Projective Ebenen*, Springer-Verlag, Berlin, 1955, p. 208.

DEPARTMENT OF MATHEMATICS, OSMANIA UNIVERSITY, HYDERABAD-7 A.P., INDIA