A NOTE ON HOMOTOPY EQUIVALENCES

R. M. VOGT

ABSTRACT. Given a homotopy equivalence $f: X \rightarrow Y$, a homotopy inverse g of f , and a homotopy $H: X \times I \rightarrow X$ from $g \circ f$ to 1_X . We show that there is a homotopy $K: Y \times I \rightarrow Y$ from $f \circ g$ to 1_Y such that $f \circ H \simeq K \circ (f \times 1_I)$ rel $X \times \partial I$ and $H \circ (g \times 1_I) \simeq g \circ K$ rel $Y \times \partial I$.

In [1], R. Lashof introduced the notion of a strong homotopy equivalence in context with the study of reductions of topological microbundles to piecewise linear or differentiable microbundles.

DEFINITION. A strong homotopy equivalence between two spaces X and Y is a quadruple (f, g, H, K) where $f: X \rightarrow Y$ and $g: Y \rightarrow X$ are maps and $H: X \times I \rightarrow X$ and $K: Y \times I \rightarrow Y$ are homotopies, $H: g \circ f \simeq 1_X$, $K: f \circ g \simeq 1_Y$, such that $f \circ H \simeq K \circ (f \times 1_I)$ rel endpoints and $H \circ (g \times 1_I) \simeq g \circ K$ rel endpoints (here I denotes the unit interval).

Now the question arises, whether any homotopy equivalence can be made into a strong one. The following proposition gives an affirmative answer. The result is also quite useful for many questions in homotopy theory. Although it is implicitly contained in various papers, I have never found it stated explicitly. The proof is very elementary, only using the basic facts about track addition of homotopies.

PROPOSITION. Let $f: X \rightarrow Y$ be a homotopy equivalence with homotopy inverse g . Let $H: X \times I \rightarrow X$ be a homotopy from $g \circ f$ to 1_X . Then there exists a homotopy $K: Y \times I \rightarrow Y$ from $f \circ g$ to 1_Y such that $f \circ H \simeq K \circ (f \times 1_I)$ rel endpoints and $H \circ (g \times 1_I) \simeq g \circ K$ rel endpoints.

Let $[X, Z]_p^q$ denote the set of equivalence classes of homotopies $F: X \times I \rightarrow Z$ such that $F|_{X \times 0} = p: X \rightarrow Z$ and $F|_{X \times 1} = q: X \rightarrow Z$. Two such homotopies are defined to be equivalent if they are homotopic rel endpoints.

Maps $h: X \rightarrow Y$ and $k: Z \rightarrow W$ induce maps $h^*: [Y, Z]_p^q \rightarrow [X, Z]_{p \circ h}^{q \circ h}$, given by $h^*[M] = [M \circ (h \times 1)]$, where $[M]$ denotes the equivalence class of M , and $k_*: [X, Z]_u^v \rightarrow [X, W]_{k \circ u}^{k \circ v}$, given by $k_*[N] = [k \circ N]$.

LEMMA 1. Let $[R] \in [U, V]_k^l$ and $[Q] \in [V, W]_u^v$. Then

$$[u \circ R + Q \circ (l \times 1)] = [Q \circ (k \times 1) + v \circ R]$$

Received by the editors August 19, 1970.
AMS 1970 subject classifications. Primary 55D10.

© American Mathematical Society 1972

in $[U, W]_{u \circ k}^{v \circ l}$, where “+” denotes track addition, the left homotopy is applied first.

PROOF. The map $G: U \times I \times I \rightarrow W$, given by $G(x, t_1, t_2) = Q(R(x, t_1), t_2)$ is the required homotopy rel endpoints between $u \circ R + Q \circ (I \times 1)$ and $Q \circ (k \times 1) + v \circ R$, because $G|_{U \times 0 \times I} = Q \circ (k \times 1)$, $G|_{U \times 1 \times I} = Q \circ (l \times 1)$, $G|_{U \times I \times 0} = u \circ R$, $G|_{U \times I \times 1} = v \circ R$.

LEMMA 2. With the notation of the proposition, $f^*: [Y, Z]_p^q \rightarrow [X, Z]_{p \circ f}^{q \circ f}$ and $g^*: [X, Z]_u^v \rightarrow [Y, Z]_{u \circ g}^{v \circ g}$ are bijections.

PROOF. Consider the composite

$$(*) \quad r: [Y, Z]_p^q \xrightarrow{f^*} [X, Z]_{p \circ f}^{q \circ f} \xrightarrow{g^*} [Y, Z]_{p \circ f \circ g}^{q \circ f \circ g} \xrightarrow{s} [Y, Z]_p^q$$

where s is given by $s[M] = [-p \circ F + M + q \circ F]$, and $F: f \circ g \simeq 1_Y$ is an arbitrary fixed homotopy. For $[Q] \in [Y, Z]_p^q$, we have

$$r[Q] = [-p \circ F + Q \circ (f \circ g \times 1) + q \circ F].$$

Substituting R by F in the previous lemma, we find that $r[Q] = [Q]$. Since s is a bijection (with inverse $[N] \mapsto [p \circ F + N - q \circ F]$), the composite $g^* \circ f^*$ is a bijection, and hence f^* injective and g^* surjective. Analogously one shows that in the sequence

$$(**) \quad [X, Z]_u^v \xrightarrow{g^*} [Y, Z]_{u \circ g}^{v \circ g} \xrightarrow{f^*} [X, Z]_{u \circ g \circ f}^{v \circ g \circ f}$$

g^* is injective and f^* is surjective. Hence, in (*), g^* and therefore f^* are bijective. Similarly in (**), f^* and hence g^* are bijective.

PROOF OF THE PROPOSITION. We are given $H: g \circ f \simeq 1_X$. Choose K to be a representative of $f^{*-1}[f \circ H] \in [Y, Y]_{f \circ g}^1$, and let H' be a representative of $g^{*-1}[g \circ K] \in [X, X]_{g \circ f}^1$. Then $K \circ (f \times 1) \simeq f \circ H$ rel endpoints, and $H' \circ (g \times 1) \simeq g \circ K$ rel endpoints. Consider the composite

$$r: [Y, X]_{g \circ f \circ g}^g \xrightarrow{f^*} [Y, Y]_{f \circ g \circ f \circ g}^{f \circ g} \xrightarrow{s} [Y, Y]_{f \circ g}^1 \xrightarrow{g^*} [Y, X]_{g \circ f \circ g}^{g \circ f \circ g}$$

where s is defined by $s[M] = [-K \circ (f \circ g \times 1) + M + K]$. Let $[R] \in [Y, X]_{g \circ f \circ g}^g$, then

$$\begin{aligned} r[R] &= [-g \circ K \circ (f \circ g \times 1) + g \circ f \circ R + g \circ K] \\ &= [-H' \circ (g \circ f \circ g \times 1) + g \circ f \circ R + H' \circ (g \times 1)] = [R]. \end{aligned}$$

The last equality follows from Lemma 1 by substituting Q by H' .

Now $[H \circ (g \times 1)] \in [Y, X]_{g \circ f \circ g}^g$, and therefore

$$\begin{aligned} [H \circ (g \times 1)] &= r[H \circ (g \times 1)] \\ &= [-g \circ K \circ (f \circ g \times 1) + g \circ f \circ H \circ (g \times 1) + g \circ K] \\ &= [-g \circ K \circ (f \circ g \times 1) + g \circ K \circ (f \circ g \times 1) + g \circ K] \\ &= [g \circ K]. \end{aligned}$$

So $H \circ (g \times 1) \simeq g \circ K$ rel endpoints.

REFERENCES

1. R. Lashof, *The immersion approach to triangulation and smoothing*, Proc. Advanced Study Institute on Algebraic Topology, Matematisk Institut, Aarhus Universitet, Aarhus, 1970.

MATEMATISK INSTITUT, AARHUS UNIVERSITET, AARHUS, DENMARK

Current address: Mathematisches Institut, Universität des Saarlandes, Saarbrücken, Federal Republic of Germany