PL INVOLUTIONS OF SOME 3-MANIFOLDS¹

MYUNG MI MYUNG

ABSTRACT. Let h_1 and h_2 be PL involutions of connected, oriented, closed, irreducible 3-manifolds M_1 and M_2 , respectively. Let $a_i, i=1, 2$, be a fixed point of h_i such that near a_i the fixed point sets of h_i are of the same dimension. Then we obtain a PL involution $h_1 \# h_2$ on $M_1 \# M_2$ induced by h_i by taking the connected sum of M_1 and M_2 along neighborhoods of a_i . In this paper, we study the possibility for a PL involution h on $M_1 \# M_2$ having a 2-dimensional fixed point set F_0 to be of the form $h_1 \# h_2$, where M_i are lens spaces. It is shown that: (1) if F_0 is orientable, then $M_1 = -M_2$ and h is the obvious involution, (2) if the fixed point set F contains a projective plane, then $M_1 = M_2 =$ a projective 3-space, and in this case, F is the disjoint union of two projective planes and h is unique up to PL equivalences, (3) if F contains a Klein bottle K , then F is the disjoint union of a Klein bottle and two points.

1. Introduction. Let M be a closed, oriented 3-manifold which is the connected sum $M_1 \# M_2$ of two irreducible 3-manifolds M_1 and M_2 , and let h be a PL involution of M with a fixed point set F containing a non-orientable surface F_0 . Since F_0 is one-sided, it would seem that h cannot interchange M_1 -part and M_2 -part and h must be obtained from involutions h_i of $M_i, i=1, 2$, by attaching two involutions along an invariant neighborhood of fixed points a_i of h_i , where near a_i the fixed point sets are of the same dimension.

Kwun [4] proved that no lens spaces except the real projective space P_3 admits orientation reversing PL involutions, and in case of P_3 there exists a unique PL involution up to PL equivalences and F is the disjoint union of a projective plane and a point. Hence it leads us to consider the possibility of the above question when M_i are isomorphic to lens spaces.

2. PL involutions of lens spaces. Let h_1 and h_2 be PL involutions of M_1 and M_2 , respectively, where $M_i, i=1, 2$, is isomorphic to a lens space (not necessarily having the natural orientation). The connected sum $M=$

Received by the editors March 1, 1971 and, in revised form, August 2, 1971.

AMS 1970 subject classifications. Primary 57A65.

Key words and phrases. PL involution, lens space, connected sum.

¹ This paper is part of the author's doctoral thesis prepared at Michigan State University under the direction of Professor Kyung W. Kwun, and the last stage of this research was supported in part by NSF Grant GP 19462.

$M_1 \# M_2$ is obtained by removing the interior of a nice 3-cell from each, and then matching the resulting boundaries using an orientation reversing homeomorphism.

If Z_2 acts on M , $H^*(M/Z_2; Q) \approx H^*(M; Q)^{Z_2}$, and since the involution reverses orientation, we obtain $H^3(M; Q)^{Z_2} = 0$, and hence $\chi(M/Z_2) = 1$. Therefore $2\chi(M/Z_2) = \chi(F) + \chi(M)$ implies $\chi(F) = 2$. Let $b_i(F)$ denote the i th mod 2 betti number of F . Then $\sum b_i(F) \leq \sum b_i(M) \leq 1 + 2 + 2 + 1 = 6$ (see [2, p. 42]). This, together with $\chi(F) = 2$ and $\dim F$ is even, implies that the possible fixed point sets are S^2 , the disjoint union of two projective planes, the disjoint union of a projective plane and a point, the disjoint union of a Klein bottle and two points, the disjoint union of a Klein bottle and S^2 , the disjoint union of $S^1 \times S^1$ and S^2 , and the disjoint union of $S^1 \times S^1$ and two points.

Now we first consider the case that F contains an orientable surface.

THEOREM 1. *Let h be a PL involution of $M = M_1 \# M_2$. If the fixed point set F contains an orientable surface, then F is a 2-sphere and $M_2 = -M_1$, h being the obvious involution.*

PROOF. Let S be an orientable surface in F . By the Alexander duality theorem [9], over the rationals Q , F separates M into two parts U and V . Hence $M = 2\bar{U}$, and we have $U \xrightarrow{i} M \xrightarrow{r} U$, such that ri is the identity, where i is the inclusion and r is a retraction. Hence we obtain the exact sequence

$$H_i(\bar{U}; Q) \xrightarrow{i_*} H_i(M; Q) \xrightarrow{r_*} H_i(\bar{U}; Q)$$

where r_*i_* is the identity. Hence $H_i(U; Q) = 0$ for $i = 1, 2$, and therefore F must be a 2-sphere.

By Milnor [8], we may say that U is isomorphic to M_1 -part and V is isomorphic to M_2 -part, and since U and V must be interchanged by h , $M_2 = -M_1$ and h is the obvious involution. This proves the theorem.

Hence we have eliminated the case that F is the disjoint union of $S^1 \times S^1$ and S^2 , the disjoint union of $S^1 \times S^1$ and two points, or the disjoint union of a Klein bottle and S^2 .

Since the case where a 2-dimensional component of F is orientable has been taken care of, we have to consider only the case where each 2-dimensional component of F is nonorientable.

THEOREM 2. *Let M be $M_1 \# M_2$, where M_i is isomorphic to a lens space, and h a PL involution on M . If a real projective plane P_2 is contained in the fixed point set F of h , then $M = P_3 \# P_3$.*

PROOF. Suppose that h fixes a real projective plane P and assume that M has been triangulated so that h is simplicial and the simplicial neighborhood U of P is an invariant regular neighborhood of P . Moreover,

assume that $h|_{U-P}$ is fixed point free. Let $p: N_1 \rightarrow M$ be the double covering obtained from M by cutting along P . Since U is orientable, but P is not, (U, P) is homeomorphic to (N, P) , where N is the mapping cylinder of a double covering $S^2 \rightarrow P$. Let U' and $(M-U)'$ be the connected manifolds obtained from U and $(M-U)$ by attaching a 3-cell to each. Then by Milnor [8], U' is isomorphic to S^3 , M_1 , or M_2 . But since $\pi_1(U')$ is Z_2 , U' cannot be isomorphic to S^3 , and we may say U' is isomorphic to M_1 . Now $h|_U$ can be extended to an orientation reversing PL involution h' of U' defined by the cone over $h|_{\text{Bd}(U)}$. Then by Kwun [4], $U' \approx M_1 \approx P_3$. Similarly $M_2 \approx P_3$. This completes the proof.

THEOREM 3. *Let $M = M_1 \# M_2$ and h a PL involution of M . If a projective plane is contained in the fixed point set F of h , then F is the disjoint union of two projective planes, and h being of the form $h_1 \# h_2$ is unique.*

PROOF. Suppose that a projective plane P is contained in F and assume that h is simplicial, the simplicial neighborhood U of P is invariant regular, and $h|_{U-P}$ is fixed point free. We have seen in Theorem 2 that $h|_{M-U}$ can be extended to an orientation reversing PL involution h' of P_3 . Let F' be the fixed point set of h' . Then by Kwun [3], F' is the disjoint union of a projective plane and a point p . By the way we extended h to h' , p is the cone vertex. Therefore $h|_{M-U}$ has a projective plane as the fixed point set F' . Hence F is the disjoint union of two projective planes.

Let P and P' be the two projective planes whose union is F . Take U and U' to be disjoint invariant regular neighborhoods of P and P' , respectively. Then by Milnor [8], $M - (U \cup U')$ is isomorphic to $S^2 \times [0, 1]$, and by Livesay [7], there is a unique involution on $S^2 \times [0, 1]$ up to PL equivalences. Hence it suffices to analyse $h|_U$ and $h|_{U'}$. But by Kwun [3] these are unique, and $h = h_1 \# h_2$, where h_1 and h_2 are the extensions of $h|_U$ and $h|_{U'}$ to P_3 . This proves the theorem.

From now on we consider the case that F contains a Klein bottle K . The only possible F is the disjoint union of K and two points. The union of K and two points can be the fixed point of an involution h . For, let the disjoint union of a projective plane A_i and a point p_i , $i=1, 2$, be the fixed point set of h_i of P_3 . Taking the connected sum $P_3 \# P_3$ along an invariant neighborhood of a_i , $a_i \in A_i$, we obtain a PL involution $h_1 \# h_2$ whose fixed point set is the disjoint union of a Klein bottle and two points. Hence we have

THEOREM 4. *Let $M = M_1 \# M_2$ and h a PL involution of M with a Klein bottle in the fixed point set F . Then F is the disjoint union of a Klein bottle and two points.*

The uniqueness question for h in case F is the disjoint union of a Klein bottle and two points is not settled, but the following theorem gives some idea how the Klein bottle is located in $P_3 \# P_3$.

THEOREM 5. *Let M be the connected sum $P_3 \# P_3$ and let K be the Klein bottle in the fixed point set F of an involution h of M . Then $\pi_1(M-U)$ is the integers Z and $M-U$ is homeomorphic to $D^2 \times S^1$.*

For the proof of Theorem 5, we first prove the following lemma.

LEMMA 6. *Let h be a PL involution of $M = P_3 \# P_3$. Then there exists a PL involution $h': S^1 \times S^2 \rightarrow S^1 \times S^2$ such that $p'h' = hp'$, where $p': S^1 \times S^2 \rightarrow P_3 \# P_3$ is a 4-to-1 covering projection.*

PROOF. Consider the usual 2-to-1 covering map $p: S^1 \times S^2 \rightarrow M$. Let $H = p_{\#}\pi_1(S^1 \times S^2)$ and $G = \pi_1(M)$. Suppose $h_{\#}H \neq H$. Since $[G:H] = [G:h_{\#}H] = 2$, there is no inclusion relation between H and $h_{\#}H$. Let $L = H \cap h_{\#}H$. Then L is a normal subgroup of G , and

$$H/L = H \cdot h_{\#}H / h_{\#}H = G / h_{\#}H = Z_2$$

implies $[H:L] = 2$. Hence $[G:L] = 4$. Furthermore $h_{\#}L = L$. Hence by the lifting theorem there is a PL involution h' on $S^1 \times S^2$ such that $p'h' = hp'$, where p' is 4-to-1. If $H = h_{\#}H$, then the construction is similar and easier. This proves the lemma.

PROOF OF THEOREM 5. Consider the double covering $p: N \rightarrow M$ obtained from M cutting along K . Then $p^{-1}(K)$ is homeomorphic to $S^1 \times S^1$. Since each component of $M_1 - p^{-1}(K)$ maps homeomorphically onto $M - K$, $p^{-1}(U)$ is a collar of $p^{-1}(K)$. So $\text{Bd}(U)$ is homeomorphic to $S^1 \times S^1$. By the Mayer-Vietoris sequence

$$H_2(M) \rightarrow H_1(S^1 \times S^1) \rightarrow H_1(\bar{U}) + H_1(M - U) \rightarrow H_1(M) \rightarrow 0$$

we obtain that $H_1(M - U)$ is a group of rank 1.

Now we show that $\pi_1(M - U)$ is the integers Z . Let $p: S^1 \times S^2 \rightarrow P_3 \# P_3$ be a 4-to-1 covering such that $h_{\#}p_{\#}\pi_1(S^1 \times S^2) = p_{\#}\pi_1(S^1 \times S^2)$. Then $h': S^1 \times S^2 \rightarrow S^1 \times S^2$ exists such that the following diagram commutes:

$$\begin{array}{ccc} S^1 \times S^2, \tilde{x}_0 & \xrightarrow{h'} & S^1 \times S^2, \tilde{x}_0 \\ \downarrow p & & \downarrow p \\ P_3 \# P_3, x_0 & \xrightarrow{h} & P_3 \# P_3, x_0 \quad (x_0 \in K). \end{array}$$

Consider the component A of $\tilde{x}_0$ in $p^{-1}(K)$. Then A is either $S^1 \times S^1$ or a Klein bottle K .

Case (1). If $A = S^1 \times S^1$, h' fixes $S^1 \times S^1$ and $S^1 \times S^2 - A = R^2 \times S^1 \cup R^2 \times S^1$ [5]. The covering $A \rightarrow K$ is at least 2-to-1. If it is actually 2-to-1, then $p^{-1}(K) = A \cup B$, where B is another $S^1 \times S^1$. Then B is contained in one of the two $R^2 \times S^1$. But h' interchanges two $R^2 \times S^1$, while $h'(p^{-1}(K)) = p^{-1}(K)$ and therefore $h'(B) = B$ which is impossible. Hence $p^{-1}(K) = A$. Hence each of the two $R^2 \times S^1$ double cover $P_3 \# P_3 - K$. Hence we have an exact sequence

$$0 \longrightarrow Z \xrightarrow{f} \pi_1(M - U) \xrightarrow{g} Z_2 \longrightarrow 0.$$

Hence if we choose $x \in \pi_1(M - U)$ such that $g(x) \neq 0$ and a generator y in Z , then $\pi_1(M - U)$ is generated by x and y . Suppose $xyx^{-1} = y^{-1}$. By abelianizing we have $y^2 = 1$. And since x is of finite order too, $H_1(M - U)$ is finite which contradicts the fact that $H_1(M - U)$ is of rank 1. Hence $\pi_1(M - U)$ is abelian, and hence $\pi_1(M - U) = Z + \text{torsion part}$. But since we have a universal covering $R^2 \times R^1 \rightarrow M - U$ and no nontrivial finite group can act freely on a finite-dimensional, contractible space, $\pi_1(M - U) = Z$.

Case (2). If $A = K$, then h' fixes only A and $S^1 \times S^2 - A = R^2 \times S^1$. If $p^{-1}(K)$ had any other component B , then $B \approx K$ and $B \subseteq R^2 \times S^1 \subseteq R^3$. But K cannot be embedded in R^3 . Hence $p^{-1}(K) = A$. Then $R^3 - A \approx R^2 \times S^1$ covers $P_3 \# P_3$ 4-to-1. Hence

$$0 \longrightarrow Z \longrightarrow \pi_1(M - U) \xrightarrow{\zeta} Z_2 + Z_2 \longrightarrow 0$$

is exact, or

$$0 \longrightarrow Z \longrightarrow \pi_1(M - U) \xrightarrow{\zeta} Z_4 \longrightarrow 0$$

is exact. Let N be a subgroup of $Z_2 + Z_2$ (or Z_4) such that $N \approx Z_2$. Let $p': X \rightarrow M - U$ be the double covering corresponding to $\zeta^{-1}(N)$. Then there exists a double covering $p'': R^2 \times S^1 \rightarrow X$. Then the argument in Case (1) shows that $\pi_1 X \approx Z$. Now consider the double covering $p': X \rightarrow M - U$. Repeating the same argument in Case (1), we get $\pi_1(M - U) = Z$.

Since $M - U$ is an irreducible, orientable, compact 3-manifold with $\pi_1(M - U) = Z$ and $\text{Bd}(M - U)$ is homeomorphic to $S^1 \times S^1$, $M - U$ is homeomorphic to $D^2 \times S^1$. This proves the theorem.

The author wishes to thank the referee for his helpful suggestions.

REFERENCES

1. J. Alexander, *On the subdivision of 3-space by a polyhedron*, Proc. Nat. Acad. Sci. U.S.A. **10** (1924), 6-8.
2. A. Borel, *Seminar on transformation groups*, Ann. of Math. Studies, no. 46, Princeton Univ. Press, Princeton, N.J., 1960. MR **22** #7129.
3. K. W. Kwun, *Scarcity of orientation reversing PL involutions of lens spaces* (to appear).

4. K. W. Kwun, *Nonexistence of orientation reversing involutions on some manifolds*, Proc. Amer. Math. Soc. **23** (1969), 725–726. MR **40** #893.
5. ———, *Piecewise linear involutions of $S^1 \times S^2$* , Michigan Math. J. **16** (1969), 93–96. MR **39** #3495.
6. G. R. Livesay, *Involutions with two fixed points on the three sphere*, Ann. of Math. (2) **78** (1963), 582–593. MR **27** #5257.
7. ———, *Fixed-point-free involutions on the 3-sphere*, Ann. of Math. (2) **72** (1960), 603–611. MR **22** #7131.
8. J. W. Milnor, *A unique decomposition theorem for 3-manifolds*, Amer. J. Math. **84** (1962), 1–7. MR **25** #5518.
9. E. H. Spanier, *Algebraic topology*, McGraw-Hill, New York, 1966. MR **35** #1007.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING,
MICHIGAN 48823

Current address: Department of Mathematics, University of Northern Iowa, Cedar
Falls, Iowa 50613