ON MODULAR COMPLEMENTED AND ANNIHILATOR ALGEBRAS

T. HUSAIN¹ AND PAK-KEN WONG²

ABSTRACT. Let A be a modular complemented algebra. We give some necessary and sufficient conditions for some special classes of algebras A to be annihilator or dual algebras.

1. Introduction. Modular complemented algebras were introduced by Yood in [7], where he developed a structure and ideal theory for these algebras. In this paper our aim is to investigate the relationship among modular complemented, annihilator and dual algebras.

Let A be a modular complemented algebra. We show that if $I + I^\perp = A$ for all minimal (left, right) ideals I of A , then A is an annihilator algebra if and only if every nonzero left ideal of A contains an element with right adjoint. If $J + J^\perp = A$ for all (left, right) ideals J of A , and A is a left or right annihilator algebra, then it is a dual algebra.

2. Notation and preliminaries. Let A be an algebra over the complex field C which is a pre-Hilbert space in the inner product (x, y) . A is called a right-modular complemented algebra if it satisfies the following conditions:

- (a) the product xy is continuous in each variable separately,
- (b) every right or left ideal I for which $I^\perp = (0)$ is dense in A ,
- (c) the intersection of all closed modular maximal right ideals is (0) and $M^\perp$ is a right ideal for each closed modular maximal right ideal M .

REMARK. From (c) it follows that every right-modular complemented algebra is semisimple.

Similarly we define a left-modular complemented algebra. It has been shown [7] that an algebra A is right-modular complemented if and only

Received by the editors July 19, 1971.

AMS 1969 subject classifications. Primary 4650; Secondary 4655.

Key words and phrases. Modular complemented algebra, annihilator algebra, dual algebra, Hilbert algebra, proper H^* -algebra.

¹ This work was supported by a N.R.C. grant.

² The second author was supported by a postdoctoral fellowship at McMaster University.

© American Mathematical Society 1972

if it is left-modular complemented. In this case, $K^\perp$ is a right (left) ideal for any right (left) ideal K of A (see [7, p. 262, Theorem 2.1]).

Let A be an algebra which is a pre-Hilbert space. As in [7], we say that u^l is a left adjoint for u if $(ux, y) = (x, u^l y)$ for all $x, y \in A$. If A is semi-simple, then the left adjoint is unique if it exists. Let $u, v \in A$. If u^l, v^l exists then $(uv)^l$ exists and $(uv)^l = v^l u^l$. If $u = u^l$, then u is said to be left selfadjoint. Similarly u^r is called the right adjoint if $(xu, y) = (x, yu^r)$ for all $x, y \in A$.

For any subset E in an algebra A , let $L(E)$ ($R(E)$) denote the left (right) annihilator of E in A . Suppose A is a topological algebra then we call A an annihilator algebra provided that, for every closed left ideal M and for every closed right ideal N , we have $R(M) = (0)$ if and only if $M = A$ and $L(N) = (0)$ if and only if $N = A$. If $M = L(R(M))$ and $N = R(L(N))$, then A is called a dual algebra.

In this paper, all algebras and linear spaces are over the complex field $\mathbb{C}$. The results of this paper heavily depend on [6] and [7].

3. Characterizations of annihilator and dual algebras. It is known that a modular complemented algebra need not be an annihilator algebra (see [7, p. 263]). In our first result, we give a characterization for annihilator modular complemented algebras.

Let A be a topological algebra. Then A is called a left annihilator algebra if for every closed right ideal M of A , $L(M) \neq (0)$ and $L(A) = (0)$. Similarly we define a right annihilator algebra.

THEOREM 3.1. *Let A be a modular complemented algebra such that $I \oplus I^\perp = A$ for all minimal (left, right) ideals I of A . Then the following statements are equivalent:*

- (i) A is an annihilator algebra.
- (ii) A is a right annihilator algebra.
- (iii) Every element of the socle S of A has a right adjoint.
- (iv) Every nonzero left ideal of A contains an element with right adjoint.

PROOF. (i) $\Rightarrow$ (ii). This is clear.

(ii) $\Rightarrow$ (iii). Suppose A is a right annihilator algebra. Let I be a minimal left ideal of A . Since $A = I \oplus I^\perp$, it follows easily that $I^\perp$ is a maximal closed left ideal. Thus by [1, p. 568, Theorem 3.4], $I^\perp$ is a modular maximal left ideal. Hence by the proof of [6, p. 44, Theorem 4.5] each element of I has a right adjoint and so does S .

(iii) $\Rightarrow$ (iv). Let M be a nonzero left ideal of A . Since S is dense in A (see the proof of [7, p. 262, Theorem 2.1]), we have $S \cap M \neq (0)$. Therefore (iv) follows from (iii).

(iv) $\Rightarrow$ (ii). Assume (iv). Let T be the set given in the proof of [7, p. 262, Theorem 2.1]. Then T is a dense right ideal of A and each element in T has a left adjoint. Let J be a proper closed left ideal of A . Then $J^\perp \neq (0)$. Let x be an element in $J^\perp$ which has a right adjoint x^r . Let $z \in J$ and $y \in T$. Then $(yx, z) = (x, y^l z) = 0$. Since T is dense in A , we have $(Ax, z) = (0)$ and so $(A, zx^r) = (0)$ which gives $zx^r = 0$. Hence $Jx^r = (0)$ and so $R(J) \neq (0)$. Therefore A is a right annihilator algebra.

(ii) $\Rightarrow$ (i). Suppose (ii) holds. Let I be a minimal left ideal of A . By the argument in (ii) $\Rightarrow$ (iii), $I^\perp$ is a modular maximal left ideal. Hence by the proof of [7, p. 263, Theorem 2.3], each element of I has a left adjoint and so does S . Hence by a similar argument in the proof (iv) $\Rightarrow$ (ii), we can show that A is a left annihilator algebra. This completes the proof of the theorem.

REMARK. A similar form of Theorem 3.1 holds for left annihilator algebras.

THEOREM 3.2. *Let A be a modular complemented algebra such that $J \oplus J^\perp = A$ for all closed (left, right) ideals J of A . Then the following statements are equivalent:*

- (i) A is a dual algebra.
- (ii) A is a left or right annihilator algebra.

PROOF. (i) $\Rightarrow$ (ii). This is clear. (ii) $\Rightarrow$ (i). Assume (ii). Then by Theorem 3.1 and its remark, A is an annihilator algebra. Let I be a proper closed right ideal of A . Since $I^\perp \neq (0)$ and since the socle of A is dense in A , it follows that $I^\perp$ contains a minimal right ideal J . By the proof of Theorem 3.1, $J^\perp$ is a closed modular maximal right ideal and clearly $J^\perp \supset I$. Let $\{M_\lambda : \lambda \in \Lambda\}$ be the family of all closed modular maximal right ideals of A containing I . Then $\{M_\lambda : \lambda \in \Lambda\}$ is not empty. Let $M = \bigcap \{M_\lambda : \lambda \in \Lambda\}$. We claim that $M = I$. In fact, suppose this is not so. Then there exists an element $x \in M$ such that $x \notin I$. Since $I \oplus I^\perp = A$, we can write $x = x_1 + x_2$ with $x_1 \in I$ and $x_2 \in I^\perp$. Hence $x_2 = x - x_1 \in I^\perp \cap M$ and so $I^\perp \cap M$ contains a minimal right ideal J whose orthogonal complement $J^\perp$ is a closed modular maximal right ideal by the above argument. Therefore by the proof of [6, p. 44, Theorem 4.5], J , and hence $I^\perp \cap M$, contains a left selfadjoint minimal idempotent e . Since $eA \subset I^\perp$, we have $(1-e)A = (eA)^\perp \supset I$. Therefore $(1-e)A \supset M$. But $e \in M$, a contradiction. Hence $M = I$. Since each $M_\lambda = R(L(M_\lambda))$, we have $I = R(L(I))$. Similarly we can show that $N = L(R(N))$ for all closed left ideals N of A . Therefore A is a dual algebra. This completes the proof.

A modular complemented algebra is said to be complete if it is complete in its inner product norm.

COROLLARY 3.3. *Let A be a complete modular complemented algebra. Then the conditions (i), (ii), (iii) and (iv) in Theorem 3.1 are equivalent to the statement that A is a dual algebra.*

PROOF. This follows from Theorems 3.1 and 3.2.

REMARK. It is well known that a proper H^* -algebra A is dual. This result also follows from Corollary 3.3, since every element of A has left and right adjoints.

If a modular complemented algebra is a normed algebra under its inner product norm, then we say that it is a normed modular complemented algebra.

THEOREM 3.4. *Let A be a complete normed modular complemented algebra. Then*

- (i) *A is a dense left ideal of a dual B^* -algebra B .*
- (ii) *If A is also a right ideal of B , then A is a dual algebra.*

PROOF. (i) For each $x \in A$, let U_x be the operation of left multiplication by x and $|U_x|$ the operator bound of U_x . Let B be the closure of $\{U_x : x \in A\}$ in the uniform topology. Since A is semisimple, the mapping $x \rightarrow U_x$ is one-one. Therefore we can identify A as a subalgebra of B . Let T be the algebraic sum of the $M^\perp$ as M ranges over the set of all closed modular maximal right ideals of A . By the proof of [6, p. 44, Theorem 4.5], T is dense in A , the left adjoint a^l exists for all $a \in T$ and $a^l \in T$. Clearly $U_a^l = (U_a)^*$ and so $T^* = T$ in B . Since T is dense in B , it follows that $B = B^*$. Therefore B is a B^* -algebra. Since the socle of B contains T , it follows from [4, p. 222, Theorem 2.1] that B is a dual algebra. For all $x, y \in A$, we have $\|xy\| \leq \|x\| \|y\|$. This shows that A is a left ideal of B and completes the proof of (i).

(ii) Now it follows from (i) that A is a dense two-sided ideal of B . For example, a proper H^* -algebra has such a property. Let M be a non-zero right ideal of A . Then M contains a minimal idempotent e of A . It is clear that $eB = eA$ and so eA is a minimal right ideal of B . Hence by [5, p. 261, Lemma (4.10.1)], eA contains a hermitian minimal idempotent $f \in B$. It is easy to see that $f \in A$ and f is a left selfadjoint element in A . Since $f \in M$, it follows from Corollary 3.3 that A is a dual algebra.

4. Hilbert algebras. A complex algebra A is called a Hilbert algebra (see [3]) if it is a pre-Hilbert space with an involution $x \rightarrow x^*$ having the following properties:

- (a) $(xy, z) = (y, x^*z)$ for all $x, y, z \in A$,
- (b) $(x, y) = (y^*, x^*)$ for all $x, y \in A$,

(c) the operation of left multiplication $y \rightarrow xy$ is continuous on A for each $x \in A$,

(d) the set of elements of the form xy is dense in A .

It follows easily from (a) and (b) that $(yx, z) = (y, zx^*)$ for all $x, y, z \in A$. It is easy to see that the right multiplication is also continuous. Since $\|x\| = \|x^*\|$, the involution is continuous.

A net $\{e_\alpha\}$ in a normed algebra A is called an approximate identity if it is bounded and $x = \lim_\alpha x e_\alpha = \lim_\alpha e_\alpha x$ for all $x \in A$.

The following result was proved by Yood for B^* -algebras (see [7, p. 267, Theorem 3.1]).

THEOREM 4.1. *Let A be an infinite-dimensional Banach $*$ -algebra with an approximate identity $\{e_\alpha\}$. Then there does not exist an inner product for A under which A becomes a pre- H^* -algebra with the same involution.*

PROOF. Suppose otherwise. Let $|x| = (x, x)^{1/2}$ be the inner product norm and let H be the completion of A in the norm $|x|$. By the proof of [7, p. 267, Theorem 3.1], H is a proper H^* -algebra and, for all $x \in A$, $|x| \leq k\|x\|$, where k is a constant. Now it is easy to see that $\{e_\alpha\}$ is also an approximate identity in H . Since H is reflexive, by [2, p. 855, Lemma 3.8], H has an identity. Therefore by [7, p. 266, Corollary 2.7], H is finite dimensional and so is A , a contradiction. This completes the proof.

THEOREM 4.2. *Let A be a Hilbert algebra. Then A is a dual algebra if and only if $I = I^{\perp\perp}$ for all closed right ideals I of A .*

PROOF. By the proof of [7, p. 268, Lemma 3.2], $x \in xA \cap Ax$. Therefore $L(A) = R(A) = (0)$. Suppose A is a dual algebra. Let I be a closed right ideal of A . Then by the proof of [7, p. 268, Lemma 3.2], we have

$$I = R(L(I)) = ((I^\perp)^{**})^\perp = I^{\perp\perp}.$$

Conversely suppose $I = I^{\perp\perp}$ for all closed right ideals I of A . It is easy to see that $I^{\perp\perp} = R(L(I)) = I$. Let J be a closed right ideal of A . Hence $J^* = (J^*)^{\perp\perp}$ and so $J = J^{\perp\perp}$. It is now easy to see that $J = L(R(J))$. Therefore A is a dual algebra.

COROLLARY 4.3. *Let A be a Hilbert algebra. If $I \oplus I^\perp = A$ for all closed right ideals I of A , then A is a dual algebra.*

REFERENCES

1. B. A. Barnes, *Modular annihilator algebras*, Canad. J. Math. **18** (1966), 566–578. MR **33** #2681.
2. P. Civin and B. Yood, *The second conjugate space of a Banach algebra as an algebra*, Pacific J. Math. **11** (1961), 847–870. MR **26** #622.

3. J. Dixmier, *Les algèbres d'opérateurs dans l'espace Hilbertien (Algèbres de von Neumann)*, Gauthier-Villars, Paris, 1969.
4. I. Kaplansky, *The structure of certain operator algebras*, Trans. Amer. Math. Soc. **70** (1951), 219–255. MR **13**, 48.
5. C. E. Rickart, *General theory of Banach algebras*, University Series in Higher Math., Van Nostrand, Princeton, N.J. 1960. MR **22** #5903.
6. B. Yood, *Ideals in topological rings*, Canad. J. Math. **16** (1964), 28–45. MR **28** #1505.
7. ———, *On algebras which are pre-Hilbert spaces*, Duke Math. J. **36** (1969), 261–272. MR **41** #4242.

DEPARTMENT OF MATHEMATICS, MCMASTER UNIVERSITY, HAMILTON, ONTARIO,
CANADA