A CONDITION FOR ANALYTIC STRUCTURE¹

RICHARD F. BASENER

ABSTRACT. Let X be a compact Hausdorff space, A a uniform algebra on X , M the maximal ideal space of A . Let $f \in A$ and let W be a component of $C \setminus f(X)$. Suppose that, for all $\lambda \in W$, $f^{-1}(\lambda) = \{x \in M \mid f(x) = \lambda\}$ is at most countable. Then there is an open dense subset U of $f^{-1}(W)$ which can be given the structure of a one-dimensional complex analytic manifold so that for all $g \in A$, g is analytic on U .

Let X be a compact Hausdorff space, let A be a uniform algebra on X , and let M be the maximal ideal space of A .

THEOREM. *Let $f \in A$ and let W be a component of $C \setminus f(X)$. Suppose that for all $\lambda \in W$, $f^{-1}(\lambda) = \{x \in M \mid f(x) = \lambda\}$ is at most countable. Then there is an open dense subset of $f^{-1}(W)$ which can be given the structure of a one-dimensional complex analytic manifold so that the functions in A become analytic there.*

This theorem partially generalizes results for the case when $f^{-1}(\lambda)$ is finite for all λ in a sufficiently large subset of W , which are essentially contained in a paper of E. Bishop ([1], see Theorem 11.2 in [2]). A key result in the proof of this earlier result is also important in the proof of our theorem (see Lemma 13 in [1] or Theorem 10.7 in [2]).

DEFINITION. Let $p \in M$, and let Φ be a continuous one-to-one map from $\{|z| < 1\}$ into M , with $\Phi(0) = p$. The set $\{\Phi(z) \mid |z| < 1\}$ is called an analytic disk through p if for all $h \in A$, $h \circ \Phi$ is analytic on $\{|z| < 1\}$.

LEMMA 1. *Let $f \in A$ and suppose that:*

(a) $|f| = 1$ on X ;

(b) $\exists p \in M$ with $f(p) = 0$;

(c) $\exists$ a closed subset Γ_0 of $\{|z| = 1\}$ having positive linear measure such that for each $\lambda \in \Gamma_0$ there is a unique point $q \in X$ with $f(q) = \lambda$.

Then $f^{-1}(\{|z| < 1\})$ is an analytic disk through p .

Presented to the Society, October 13, 1971; received by the editors December 30, 1971.

AMS 1970 subject classifications. Primary 46J10, 46J15.

Key words and phrases. Analyticity, maximal ideal space.

¹ This research was supported in part by the National Science Foundation under grant GP-28574.

© American Mathematical Society 1972

We also need the following elementary result (Lemma 11.1 in [2]).

LEMMA 2. *Let $f \in A$ and let W be a component of $C \setminus f(X)$. Fix $\lambda \in W$. If f takes on the value λ on M , then f takes on every value in W on M .*

One further result is needed.

LEMMA 3. *Let $f \in A$ and let W be a component of $C \setminus f(X)$. Let $z \in W$, and suppose that $f^{-1}(z)$ is at most countable. Given a neighborhood $\mathcal{O}$ of a point $x \in f^{-1}(z)$, there is a compact neighborhood N of x with the properties that:*

- (a) N is A -convex, i.e., the maximal ideal space of $\bar{A}|_N$ is N ;
- (b) $z \notin f(\partial N)$;
- (c) $N \subseteq \mathcal{O} \cap f^{-1}(W)$.

Furthermore, $f(N)$ is a neighborhood of z in C .

PROOF OF LEMMA 3. Choose $\varepsilon > 0$ and functions $g_1, g_2, \dots, g_n \in A$ so that

$$N = \{y \in M \mid |g_j(y) - g_j(x)| \leq \varepsilon, 1 \leq j \leq n\} \subseteq f^{-1}(W) \cap \mathcal{O}.$$

Since $f^{-1}(z)$ is at most countable, we may adjust ε so that, for all $y \in f^{-1}(z)$, $|g_j(y) - g_j(x)| \neq \varepsilon$, $j = 1, 2, \dots, n$. Now if $y \in \partial N$, then $|g_j(y) - g_j(x)| = \varepsilon$ for some j , so $f^{-1}(z) \cap \partial N = \emptyset$ and it is clear that (a), (b) and (c) are satisfied.

By the local maximum modulus principle, the Shilov boundary of $\bar{A}|_N$ is contained in ∂N (note that N does not meet the Shilov boundary of A). Hence $\bar{A}|_N$ may be regarded as a uniform algebra on ∂N . Also, $z \in f(N) \setminus f(\partial N)$. By Lemma 2, $f(N)$ contains the component of z in $C \setminus f(\partial N)$, so $f(N)$ is a neighborhood of z as claimed.

PROOF OF THEOREM. We will prove the following.

Assertion. If $f^{-1}(W)$ is nonempty, $\exists p \in f^{-1}(W)$ with a neighborhood in M which is an analytic disk through p .

Assume for the moment that the assertion has been established. Given any $x \in f^{-1}(W)$ and any neighborhood $\mathcal{O}$ of x , we may apply Lemma 3 to find a compact neighborhood N of x with properties (a), (b) and (c). Then the assertion may be applied to $f|_N \in \bar{A}|_N$, to yield a point $p \in \mathcal{O}$ with a neighborhood which is an analytic disk through p . Thus the set of such points is dense in $f^{-1}(W)$, and it is obviously open, so the Theorem follows.

PROOF OF ASSERTION. Suppose, on the contrary, that no point of $f^{-1}(W)$ has a neighborhood which is an analytic disk. We will use this assumption to find a point $z \in W$ such that $f^{-1}(z)$ is uncountable, contradicting the hypothesis of the Theorem. The uncountable set will consist of limit points obtained from sequences $\{x_{i_1 i_2 \dots i_n}\}_{n=1}^\infty$ in M , which we will now define inductively.

Step 1. Since $f^{-1}(W)$ is nonempty, by Lemma 2 we have $f(f^{-1}(W)) = W$, so by Lemma 1 and the assumption that no point of $f^{-1}(W)$ has a neighborhood which is an analytic disk, there are distinct points $x_0, x_1 \in M$ and a corresponding $z_1 \in W$ such that $f(x_0) = f(x_1) = z_1$. By Lemma 3 there are disjoint compact A -convex neighborhoods N_{i_1} of x_{i_1} with the properties $N_{i_1} \subseteq f^{-1}(W)$, $z_1 \notin f(\partial N_{i_1})$, and $f(N_{i_1})$ is a neighborhood of z_1 , $i_1 = 0, 1$. Choose ε_1 with $0 < \varepsilon_1 < 1$ so that

$$(A_1) \quad \{|z - z_1| \leq \varepsilon_1\} \subseteq f(N_{i_1}) \setminus f(\partial N_{i_1}), \quad i_1 = 0, 1.$$

Fix i_1 and look at

$$M_{i_1} = \{y \in N_{i_1} \mid |f(y) - z_1| \leq \varepsilon_1\} = N_{i_1} \cap f^{-1}(\{|z - z_1| \leq \varepsilon_1\}).$$

M_{i_1} is a nonempty compact A -convex set, so the maximal ideal space of $\bar{A}|_{M_{i_1}}$ is M_{i_1} . Let $X_{i_1} = \{y \in N_{i_1} \mid |f(y) - z_1| = \varepsilon_1\}$.

Observe that $\partial M_{i_1} \subseteq (\partial N_{i_1} \cup X_{i_1}) \cap M_{i_1} = X_{i_1}$ since $\partial N_{i_1} \cap M_{i_1} = \emptyset$ by (A_1) . Also observe that M_{i_1} does not meet the Shilov boundary of A . By the local maximum modulus principle, the Shilov boundary of $\bar{A}|_{M_{i_1}}$ is contained in X_{i_1} . Thus we may apply Lemma 1 with $X = X_{i_1}$, $A = \bar{A}|_{X_{i_1}}$, $M = M_{i_1}$, $f = (f - z_1)/\varepsilon_1$, to conclude that (since there is no analytic disk through any point of $f^{-1}(W)$):

$$(B_1) \quad \begin{aligned} &\text{linear measure } \{z \in C \mid |z - z_1| = \varepsilon_1 \\ &\quad \text{and } f^{-1}(z) \text{ is unique in } N_{i_1}\} = 0. \end{aligned}$$

(Note that $f^{-1}(z) \cap N_{i_1} = f^{-1}(z) \cap M_{i_1}$ if $|z - z_1| = \varepsilon_1$. The set described in statement (B_1) is measurable since Lemma 3 implies that it is closed. In fact, suppose that $\zeta_1, \zeta_2, \dots \in C$, $\zeta_j \rightarrow \zeta \in C$, $|\zeta_j - z_1| = \varepsilon_1$ and there is precisely one $\alpha_j \in N_{i_1}$ such that $f(\alpha_j) = \zeta_j$, $j = 1, 2, \dots$. Let $\alpha \in f^{-1}(\zeta) \cap N_{i_1}$ and let V be any neighborhood of α in M . By Lemma 3, $f(V \cap N_{i_1})$ is a neighborhood of ζ in C , hence ζ_j is in $f(V \cap N_{i_1})$ for all $j \geq j_0$ for some j_0 . But $f^{-1}(\zeta_j) \cap N_{i_1} = \{\alpha_j\}$, so α_j is in V for $j \geq j_0$. Thus α_j converges to α , and α is unique.)

This completes the first step.

Inductive hypothesis. Suppose that $z_n \in W$, $x_a \in M$, $N_a \subseteq M$ (where $a = i_1 \cdots i_n$) and $\varepsilon_n > 0$ have been chosen for $i_j = 0, 1$, $1 \leq j \leq n$. Assume that they satisfy the conditions: the 2^n points x_a are all distinct; $f(x_a) = z_n$; N_a is a compact A -convex neighborhood of x_a ; $N_a \cap N_b = \emptyset$ (where $b = j_1 \cdots j_n$) unless $i_1 = j_1, \dots, i_n = j_n$; $N_a \subseteq \text{interior } N_{i_1 \cdots i_{n-1}}$, if $n > 1$; $\varepsilon_n < 1/n^2$;

$$(A_n) \quad \{|z - z_n| \leq \varepsilon_n\} \subseteq f(N_a) \setminus f(\partial N_a);$$

$$(B_n) \quad \begin{aligned} &\text{linear measure } \{z \in C \mid |z - z_n| = \varepsilon_n \\ &\quad \text{and } f^{-1}(z) \text{ is unique in } N_a\} = 0. \end{aligned}$$

Step $n+1$. By (B_n) , $\exists z_{n+1} \in \{|z - z_n| = \varepsilon_n\}$ such that for all $i_1 \cdots i_n$, there are two distinct points $x_c \in N_a$ (where $c = i_1 \cdots i_{n+1}$, $i_{n+1} = 0, 1$) with $f(x_c) = z_{n+1}$. By (A_n) , we have $x_c \in N_a \setminus f^{-1}(f(\partial N_a))$. Fix $a = i_1 \cdots i_n$ and choose disjoint open sets $\mathcal{O}_0, \mathcal{O}_1$ with

$$x_c \in \mathcal{O}_{i_{n+1}} \subseteq N_a \setminus f^{-1}(f(\partial N_a)), \quad i_{n+1} = 0, 1.$$

Fix $i_{n+1} = 0$ or 1 and apply Lemma 3 with

$$A = \bar{A}|_{\partial N_a}, \quad X = \partial N_a, \quad M = N_a, \quad f = f|_{N_a},$$

$$W = \text{component of } z_{n+1} \text{ in } C \setminus f(\partial N_a), \quad z = z_{n+1}, \quad \mathcal{O} = \mathcal{O}_{i_{n+1}}, \quad x = x_c.$$

We conclude that there exist disjoint compact A -convex neighborhoods N_c of x_c , $i_{n+1} = 0, 1$, with the properties $z_{n+1} \notin f(\partial N_c)$,

$$N_c \subseteq N_a \setminus f^{-1}(f(\partial N_a)),$$

$f(N_c)$ is a neighborhood of z_{n+1} . Do this for all indices $c = i_1 \cdots i_{n+1}$, $i_j = 0, 1$, $1 \leq j \leq n+1$.

From the definition of the N_c it follows that we can choose an ε_{n+1} with $0 < \varepsilon_{n+1} < 1/(n+1)^2$ so that for all $i_1 \cdots i_{n+1}$ we have

$$(A_{n+1}) \quad \{|z - z_{n+1}| \leq \varepsilon_{n+1}\} \subseteq f(N_c) \setminus f(\partial N_c).$$

Fix $c = i_1 \cdots i_{n+1}$ and define

$$M_c = \{y \in N_c \mid |f(y) - z_{n+1}| \leq \varepsilon_{n+1}\},$$

$$X_c = \{y \in N_c \mid |f(y) - z_{n+1}| = \varepsilon_{n+1}\}.$$

As in Step 1, the maximal ideal space $\bar{A}|_{M_c}$ is M_c and the Shilov boundary of $\bar{A}|_{M_c}$ is contained in X_c . ((A_{n+1}) guarantees that $\partial M_c \subseteq X_c$.) We may therefore apply Lemma 1 with

$$X = X_c, \quad A = \bar{A}|_{X_c}, \quad M = M_c, \quad f = (f - z_{n+1})/\varepsilon_{n+1}.$$

We conclude that

$$(B_{n+1}) \quad \begin{aligned} &\text{linear measure } \{z \in C \mid |z - z_{n+1}| = \varepsilon_{n+1} \\ &\text{and } f^{-1}(z) \text{ is unique in } N_c\} = 0. \end{aligned}$$

This equation holds for all indices $c = i_1 \cdots i_{n+1}$, $i_j = 0, 1$, $1 \leq j \leq n+1$, and the induction is complete.

By the above construction $\{z_n\}$ is Cauchy, so $z_n \rightarrow z$ for some $z \in C$; $z \in W$ since, for each n , $z_n \in f(N_a) \subseteq f(N_0) \cup f(N_1)$, a compact subset of W .

Let $I = (i_1, i_2, \cdots, i_n, \cdots)$ be an infinite sequence of 0's and 1's. Some subnet of the sequence $x_{i_1}, x_{i_1 i_2}, x_{i_1 i_2 i_3}, \cdots$ converges to a point $x_I \in M$, and by continuity $f(x_I) = z$. In this way we associate an x_I with each I .

Since there are uncountably many distinct I 's, the proof will be complete if we show that $x_I \neq x_J$ whenever $I \neq J$. Suppose therefore that $i_1 = j_1$, $i_2 = j_2, \dots, i_m = j_m, i_{m+1} \neq j_{m+1}$. We know that $x_{i_1} \in N_{i_1}, x_{i_1 i_2} \in N_{i_1 i_2}, \dots$ and $N_{i_1} \supseteq N_{i_1 i_2} \supseteq N_{i_1 i_2 i_3} \dots$ by the inductive hypotheses. Thus $x_I \in \bigcap_{n=1}^{\infty} N_a$. Similarly $x_J \in \bigcap_{n=1}^{\infty} N_b$ where $b = j_1 \dots j_n$. But $N_{i_1 \dots i_{m+1}} \cap N_{j_1 \dots j_{m+1}} = \emptyset$ since $i_{m+1} \neq j_{m+1}$, so $x_I \neq x_J$.

B. Cole has pointed out that if one merely assumes that $f^{-1}(\lambda)$ is countable for λ in a subset of W of positive plane measure, the same proof shows that there is at least one point $p \in f^{-1}(W)$ with a neighborhood in M which is an analytic disk through p .

I would like to thank J. Wermer, who pointed out to me the question answered by the Theorem, for many helpful discussions and suggestions.

REFERENCES

1. E. Bishop, *Holomorphic completions, analytic continuation, and the interpolation of semi-norms*, Ann. of Math. (2) 78 (1963), 468–500. MR 27 #4958.
2. J. Wermer, *Banach algebras and several complex variables*, Markham, Chicago, Ill., 1971.

DEPARTMENT OF MATHEMATICS, BROWN UNIVERSITY, PROVIDENCE, RHODE ISLAND 02912

Current address: Department of Mathematics, Yale University, New Haven, Connecticut 06520