CLASSICAL QUOTIENT RINGS OF PWD'S

ROBERT GORDON

ABSTRACT. Piecewise domains which are right orders in semi-primary rings are characterized. An example is given showing the result obtained is "best possible". A further example is obtained of a prime right Goldie ring possessing a regular element which becomes a left zero divisor in some prime overring. This example leads to the construction of a PWD R not satisfying the regularity condition, but for which $R/N(R)$ is right Goldie.

Introduction. The principle result of this note is a characterization of PWD's (piecewise domains) which are right orders in semiprimary rings. The main device employed here is L. W. Small's concept of exhaustive set. Probably the most important consequence of the characterization is that a right noetherian PWD is a right order in a right artinian PWD. In the more general setting our result is less satisfactory because an assumption is made about each of the factor rings $R/T_i(R)$ (see the theorem in §1). However, an example given at the end of §1 shows that this is unavoidable: There we exhibit a right Goldie PWD R having $R/N(R)$ right Goldie which is not a right order in a semiprimary ring.

In §2, as a problem closely related to the existence of classical quotient rings, we study the regularity condition in PWD's. Necessary and sufficient conditions are obtained for a PWD R to satisfy the regularity condition in terms of the torsion-freeness of R as an $R/N(R)$ -module. It is shown that any PWD R with $R/N(R)$ right Goldie satisfies a certain one-sided regularity condition. That the regularity condition then need not be implied is demonstrated by an example. Since this example is surely of more general interest than its application, we single it out here:

A right Ore domain D is constructed so that for some elements $a, b, c, d \in D$, the equations $ax + by = 0, cx + dy = 0$ have a nontrivial solution in a domain E extending D , but no nontrivial solution in D . (This means that the prime right Goldie ring D_2 contains a regular element which becomes a left zero divisor in the prime overring E_2 .) It is amusing to note that there

Received by the editors September 7, 1971.

AMS 1970 subject classifications. Primary 16A08; Secondary 16A02, 15A06.

Key words and phrases. Piecewise domain, essentiality condition, exhaustive set, order, classical quotient ring, Goldie ring, principal right ideals projective, torsion-free, left regularity condition, Ore domain, linear equations.

© American Mathematical Society 1972

is no such example where the domain D is two-sided Ore. Moreover, the domain E extending D in which the equations have a nontrivial solution cannot imbed in a division ring.

Throughout the paper, the prime radical of a ring R is denoted by $N(R)$ and $T_i(R)$ stands for the intersection of $N(R)$ with the left annihilator, $\ell(N(R)^i)$, of $N(R)^i$. The essentiality condition (see [3]) holds in R if aR is an essential right ideal whenever a is an element of R with zero right annihilator, $\iota(a)$. As defined in [4], R is a PWD with respect to the complete set $e_1, \dots, e_n$ of orthogonal idempotents if $xy=0$ for $x \in e_i Re_k$ and $y \in e_k Re_j$ implies $x=0$ or $y=0$. The reader is referred to [9] for the notion of exhaustive set. I would like to thank Professors A. V. Jategaonkar and L. W. Small for conversations about the material in this paper which were both stimulating and helpful.

1. Classical quotient rings. We would like to show first that a semi-primary classical right quotient ring of a PWD is a PWD. This is easily accomplished by means of the following theorem.

THEOREM A. *Suppose e and f are nonzero idempotents in a ring R which is a right order in a semiprimary ring Q . Then $eQf = eRffQf$, and the semiprimary ring fQf is the classical right quotient ring of fRf .*

PROOF. The proof follows from the technique used in proving a theorem of Small [9, Theorem 3], of which this is a mild generalization.

PROPOSITION. *If the PWD R is a right order in a semiprimary ring Q , then Q is also a PWD.*

PROOF. Let $\{e_i\}$ be a complete set of orthogonal idempotents with respect to which R is a PWD. Suppose $xy=0$ for $x \in e_i Qe_k$ and $y \in e_k Qe_j$. By Theorem A, $x=ab^{-1}$ and $y=cd^{-1}$ for elements $a \in e_i Re_k$, $b \in e_k Re_k$, $c \in e_k Re_j$, and $d \in e_j Re_j$. Since $b^{-1}c \in e_k Qe_k e_k Re_j \subseteq e_k Qe_j$, $b^{-1}c=uv^{-1}$ where $u \in e_k Re_j$ and $v \in e_j Re_j$. We have $auv^{-1}d^{-1}=0$ implying $au=0$. Since R is a PWD with respect to $\{e_i\}$, $a=0$ or $u=0$, so $a=0$ or $c=0$. Thus $x=0$ or $y=0$, and Q is a PWD.

The lemma which follows is critical to the characterization mentioned in the introduction.

LEMMA. *Let S be a semiprime right Goldie PWD with respect to the complete set of orthogonal idempotents $\{e_i\}_{i=1}^n$. Then*

$$M = \left\{ \sum_{i=1}^n s_i \mid 0 \neq s_i \in e_i S e_i, 1 \leq i \leq n \right\}$$

is an exhaustive set of S .

PROOF. That M is a multiplicatively closed set of regular elements of S is inherent in the definition of PWD. We must show that if $q \in Q$, where Q is the classical right quotient ring of S (which exists by Goldie's theorem), then $q = sm^{-1}$ for some $s \in S$ and $m \in M$. To this end, write $q = \sum_{i,j} q_{ij}$, $q_{ij} \in e_i Q e_j$. By Theorem A we have $q_{ij} = a_{ij} b_{ij}^{-1}$ for some $a_{ij} \in e_i S e_j$ and $b_{ij} \in e_j S e_j$. Using a well-known property of quotient rings, there exist an element $d_j \in e_j S e_j$ and elements $c_{ij} \in e_j S e_j$ such that $b_{ij}^{-1} = c_{ij} d_j^{-1}$ for all i ($d_j^{-1} \in e_j Q e_j$). So q_{ij} may be written $q_{ij} = x_{ij} y_j^{-1}$ for all i, j , where $x_{ij} \in e_i S e_j$ and $y_j \in e_j S e_j$. Thus $q = \sum_{i,j} x_{ij} y_j^{-1} = (\sum_{i,j} x_{ij}) (\sum_j y_j^{-1}) = (\sum_{i,j} x_{ij}) (\sum_j y_j)^{-1}$, and we see q has the required form.

We proceed to state and prove our main result.

THEOREM. *If R is a PWD, then R is a right order in a semiprimary ring if and only if $R/T_i(R)$ satisfies the essentiality condition for all i .*

PROOF. It is well known that the "only if" part of the theorem holds for any ring R . To prove the "if" part, let R be a PWD with respect to $\{e_i\}$ and assume the rings $R/T_i(R)$ all have the essentiality condition. Denote the canonical image of $r \in R$ in $R/N(R)$ by $\bar{r}$. By the main theorem of [4], $\bar{R}$ is a PWD with respect to $\{\bar{e}_i\}$. Also, since $N(R)$ is known to be nilpotent [4], $N(R) = T_i(R)$ for some i . In particular $\bar{R}$, as a semiprime PWD with the essentiality condition, is a right Goldie ring [3, Proposition 1].

Let $M = \{\sum r_i | 0 \neq r_i \in e_i R e_i\}$. Then, since $\bar{M} = \{\sum s_i | 0 \neq s_i \in \bar{e}_i \bar{R} \bar{e}_i\}$ and $\bar{R}$ is a semiprime right Goldie PWD with respect to $\{\bar{e}_i\}$, $\bar{M}$ is an exhaustive set of $\bar{R}$ by the lemma above. But M is clearly a multiplicatively closed set of regular elements of R and thus an exhaustive set of regular elements. The theorem now follows from the following general criterion of Small.

THEOREM B. *Let R be a ring satisfying*

- (1) $N(R)$ is nilpotent;
- (2) $R/T_i(R)$ has the essentiality condition for all i ;
- (3) $R/N(R)$ is right Goldie; and
- (4) R has an exhaustive set of regular elements.

Then R is a right order in a semiprimary ring.

PROOF. The proof is virtually that of [9, Theorem 1 and Corollary 1]; we omit the details.

COROLLARY 1. *A necessary and sufficient condition for a PWD R to be a right order in a right artinian ring is for $R/T_i(R)$ to have finite right Goldie dimension for all i .*

COROLLARY 2. *A right noetherian PWD is a right order in a right artinian ring.*

As an application, consider a right noetherian ring R which is a direct sum of uniform right ideals and has zero right singular ideal (so R is a PWD). Then R is a finite direct product of indecomposable rings R_i each of which satisfies the same hypothesis that R does. Each R_i is a right order in an indecomposable (by Theorem A) right artinian ring Q_i which also is a direct sum of uniform right ideals and has zero right singular ideal. A typical Q_i is characterized in [2, Theorem 4.1] as a certain blocked triangular matrix ring with entries in subgroups of the additive group of a division ring. This can be shown to lead to a similar, albeit considerably more involved, characterization of the R_i 's.

In [4], it was shown that any PWD R is a split extension of $N(R)$ by a unitary subring of R uniquely determined up to an inner automorphism of R . Consequently R can be considered as an $R/N(R)$ -bimodule in an essentially unique way. The following is by way of generalization of a remark made in [4, §4] about rings with principal right ideals projective.

COROLLARY 3. *If the PWD R is a finitely generated projective right $R/N(R)$ -module (e.g., if $N(R)_R$ is finitely generated projective [4]) and $R/N(R)$ is right Goldie, then R is a right order in a right artinian ring.*

PROOF. It follows from [4, §2, Corollary 1] that, for each i , R is a split extension by $R/T_i(R)$ and $R/T_i(R)$ is a split extension by $R/N(R)$. Clearly, then, each ring $R/T_i(R)$ is finitely generated projective as a right $R/N(R)$ -module. But then each has finite right Goldie dimension by [4, §2, Proposition 1].

EXAMPLE. Let R be the ring

$$\begin{bmatrix} F[x] & F[x] & F(x) \\ 0 & F & F(x) \\ 0 & 0 & F(x) \end{bmatrix}$$

where F is any field. Then R has the following properties.

- (1) R is a left but not a right order in a semiprimary ring.
- (2) R is a right (but not left) Goldie PWD.
- (3) $R/N(R)$ is an hereditary noetherian ring.
- (4) R satisfies the regularity condition.

PROOF. First, R is right Goldie because the right socle of R is essential and artinian. Also, R is a left order in the semiprimary ring

$$\begin{bmatrix} F(x) & F(x) & F(x) \\ 0 & F & F(x) \\ 0 & 0 & F(x) \end{bmatrix}$$

(which is not one-sided artinian). But R is not a right order in a semi-primary ring, since

$$R/T_1(R) \simeq \begin{bmatrix} F[x] & F[x] \\ 0 & F \end{bmatrix} \times F(x)$$

does not satisfy the essentiality condition (by an example of Herstein and Small [5, §3]). The rest is obvious (see the corollary in §2).

According to [4, §4, Theorem], the ring R in the preceding example cannot be right semihereditary. In fact, it is not very hard to see that R does not have principal right ideals projective. Is there an example of a ring R with principal right ideals projective having R and $R/N(R)$ right Goldie, which is not a right order in a semiprimary ring? Such an example (if it exists!) would be interesting in the light of the rather weak conditions which suffice for a semihereditary ring to be an order in a semiprimary ring (see [3] and [4]). We have one trivial comment on this problem: The right Goldie ring

$$R = \begin{bmatrix} K & K & K \\ 0 & k & K \\ 0 & 0 & K \end{bmatrix},$$

where K is an infinite field extension of the field k , has $R/N(R)$ right Goldie and principal right ideals projective; it is a right order in a semi-primary ring but not in a one-sided artinian ring (cf. [4, §4, Theorem]).

2. The regularity condition. It turns out to be quite natural to study a one-sided regularity condition in PWD's. Accordingly, we say that a PWD R satisfies the *left regularity condition* if $a + N(R)$ regular in $R/N(R)$ implies a left regular in R (i.e., $\ell(a) = 0$). (Then, of course, R satisfies the regularity condition if and only if R satisfies both the left and right regularity conditions.) We use the term "torsion-free" in the sense of Levy [7].

PROPOSITION 1. *If R is a PWD, the following statements are equivalent.*

- (1) *R satisfies the left regularity condition.*
- (2) *R is a torsion-free right $R/N(R)$ -module.*
- (3) *$f_i R f_j$ is a torsion-free right $f_j R f_i$ -module for every complete set $f_1, \dots, f_r$ of orthogonal idempotents in R which are centrally primitive modulo $N(R)$.*

PROOF. We prove (2) $\Rightarrow$ (1), the remaining implications being routine. The proof is by induction on the index of nilpotence of $N(R)$. Noting the case $N(R) = 0$ is trivial, we assume $N(R) \neq 0$. As follows from [4, §2, Corollary 1], there exist (unitary) subrings S and S_1 of R with $S \subseteq S_1 \subset R$ such that R is a split extension of $N(R)$ by S and a split extension of $T_1(R)$

by S_1 . Now the right S -module S_1 , as a submodule of the torsion-free (since (2) holds) right S -module R , is torsion-free. Moreover, the index of nilpotence of $N(S_1)$ is one less than that of $N(R)$ and S_1 is a PWD [4, §1, Corollary 2]. So our induction hypothesis implies $R/T_1(R) \simeq S_1$ satisfies the left regularity condition.

Let $a \in R$ be regular mod $N(R)$. Since $N(R/T_1(R)) = N(R)/T_1(R)$, the canonical image of a in $R/T_1(R)$ is regular mod $N(R/T_1(R))$. Thus $\ell(a) \subseteq T_1(R) \subseteq \ell(N(R))$ by the left regularity condition in $R/T_1(R)$. But a can be written $a = s + n$ where $n \in N(R)$ and s is a regular element of S . So if $x \in \ell(a)$, then $xs = xa = 0$ which has the consequence $x = 0$.

COROLLARY. *A PWD R for which $R/N(R)$ is a finite direct product of domains (cf. [4, §1, Corollary 3]) satisfies the regularity condition.*

PROPOSITION 2. *If R is a PWD such that $R/N(R)$ is right Goldie, then R satisfies the left regularity condition.*

PROOF. The same method used in the proof of Proposition 1 works here: Assume $N(R) \neq 0$, the proposition being trivial if $N(R) = 0$. Clearly $R/T_1(R)$ satisfies the same hypothesis that R does. So we may assume by induction that the left regularity condition holds in $R/T_1(R)$. Hence, assuming $a + N(R)$ regular in $R/N(R)$, we obtain $\ell(a) \subseteq \ell(N(R))$. But $aR + N(R)$ is an essential right ideal of R and, if $x \in \ell(a)$, $x(aR + N(R)) = xaR + xN(R) = 0$. So $x = 0$, since R has zero right singular ideal according to [3, Lemma 1].

We proceed to give an example of a ring R satisfying the hypothesis of Proposition 2, but which does not satisfy the regularity condition: Let F be a field and K an extension field of F possessing a nonsurjective (nonzero) endomorphism σ which fixes F elementwise. (This situation can be realized given any field F .) Let D be the twisted polynomial ring $K[z, \sigma]$ in the indeterminate z where multiplication is defined by $kz = z\sigma(k)$ ($k \in K$) and its consequences.

The domain D is well known to be right but not left Ore. In particular, there exist nonzero elements $u, v \in D$ such that $Du \cap Dv = 0$. The choice of u, v implies the directness of the sum $Du + Duv + Duv^2 + Duv^3$. So, since F is contained in the center of D , the F -subalgebra of D generated by u, uv, uv^2 , and uv^3 is in fact freely generated by u, uv, uv^2 , and uv^3 , making use of a result of Jategaonkar [6]. Furthermore, notice that the relation $uvu^{-1} = uv^2(uv^3)^{-1}$ in the right quotient division ring $Q(D)$ of D would lead to the absurdity $v^2 = 1$. Thus setting $a = u$, $b = uv^3$, $c = uv$, and $d = uv^2$, we have $ca^{-1} \neq db^{-1}$ in $Q(D)$.

Let $A = F\langle a, b, c, d, \zeta, \eta \rangle$ be the free associative F -algebra on the letters a, b, c, d, ζ , and η and let I be the ideal of A generated by $a\zeta + b\eta$

and $c\zeta + d\eta$. By making repeated use of the substitutions $a\zeta \rightarrow -b\eta$, $c\zeta \rightarrow -d\eta$, an element $f \in A$ can be brought to a unique "normal form" f' ; that is, no monomial in the expression for f' contains an occurrence of $a\zeta$ or $c\zeta$. Since $f \in I \Leftrightarrow f' = 0$, we obtain $F\langle a, b, c, d \rangle \cap I = 0$, $\zeta \notin I$, and $\eta \notin I$. Thus the F -algebra $L = A/I$ contains (a canonical copy of) $F\langle a, b, c, d \rangle$ and the equations

$$(*) \quad ax + by = 0, \quad cx + dy = 0$$

have a nontrivial solution in L . Further, using straightforward free algebra techniques (see, for example, Shepherdson [8, §4]) one may easily verify that L is a domain.

By a theorem of P. M. Cohn [1, Theorem 2.5], the free product (= coproduct here) $E = D *_F L$ of the F -algebras D and L is a domain. The domain E contains, of course, both D and L as subdomains. In particular, the equations (*) have a nontrivial solution in E . We claim they have none in D . Indeed, the contrary assumption leads via a short computation to $ca^{-1} = db^{-1}$ holding in $Q(D)$, against our construction.

Finally, we take R to be the ring

$$\begin{bmatrix} D_2 & 0 \\ E_2 & D_2 \end{bmatrix}.$$

R is a PWD and $R/N(R)$ is right Goldie (because D is right Ore). Let

$$\alpha = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \text{and} \quad \beta = \begin{bmatrix} x_0 & 0 \\ y_0 & 0 \end{bmatrix}$$

where $x = x_0$, $y = y_0$ is a nontrivial solution of (*) in E . Since the equations (*) have no nontrivial solution in D , $\iota(\alpha) = 0$ in D_2 implying the regularity of α in D_2 . But, in R , $\begin{bmatrix} 1 & 0 \\ 0 & \alpha \end{bmatrix} \begin{bmatrix} 0 & 0 \\ \beta & 0 \end{bmatrix} = 0$, $\begin{bmatrix} 1 & 0 \\ 0 & \alpha \end{bmatrix}$ is regular modulo $N(R)$, and $0 \neq \begin{bmatrix} 0 & 0 \\ \beta & 0 \end{bmatrix} \in N(R)$. Therefore the regularity condition fails to hold in R .

REFERENCES

1. P. M. Cohn, *On the free product of associative rings. II. The case of (skew) fields*, Math. Z. 73 (1960), 433–456. MR 22 #4747.
2. R. Gordon, *Rings faithfully represented on their left socle*, J. Algebra 7 (1967), 303–342. MR 36 #1483.
3. ———, *Semi-prime right Goldie rings which are direct sums of uniform right ideals*, Bull. London Math. Soc. 3 (1971), 277–282.
4. R. Gordon and L. W. Small, *Piecewise domains*, J. Algebra (to appear).
5. I. N. Herstein and L. W. Small, *Regular elements in P.I.-rings*, Pacific J. Math. 36 (1971), 327–330.

6. A. V. Jategaonkar, *Ore domains and free algebras*, Bull. London Math. Soc. **1** (1969), 45–46. MR **39** #241.
7. L. S. Levy, *Torsion-free and divisible modules over non-integral domains*, Canad. J. Math. **15** (1963), 132–151. MR **26** #155.
8. J. C. Shepherdson, *Inverses and zero divisors in matrix rings*, Proc. London Math. Soc. (3) **1** (1951), 71–85. MR **13**, 7.
9. L. W. Small, *Orders in Artinian rings. II*, J. Algebra **9** (1968), 266–273. MR **37** #6315.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UTAH 8411