SELF-UNIVERSAL CRUMPLED CUBES AND A DOGBONE SPACE

E. H. ANDERSON¹

ABSTRACT. The question of whether each self-universal crumpled cube is universal is answered negatively by presenting an example of a dogbone space which is not topologically E^3 but which can be expressed as a sewing of two crumpled cubes, one of which is self-universal.

C. D. Bass and R. J. Daverman [2] presented a brief paper indicating that the solid Alexander horned sphere is an example of a crumpled cube which is self-universal but not universal, thus answering negatively the question asked by C. E. Burgess and J. W. Cannon in [4] of whether each self-universal crumpled cube is universal. The validity of the example presented by Bass and Daverman depends on a claim that a certain upper semicontinuous decomposition of S^3 into points and tame arcs, described in [2], is not topologically S^3 , which in turn depends on the validity of four lemmas which are stated in [2, §2]. The proofs of these four lemmas appear to entail nontrivial arguments which are not included in [2].

In this note, we present an example of a dogbone space, an upper semicontinuous decomposition of S^3 into points and tame arcs whose nondegenerate elements can be expressed as the intersection of a tower of solid double tori, which is not topologically S^3 . The dogbone space can be described as the result of a sewing of two crumpled cubes, one of which is self-universal. Thus, the question asked by Burgess and Cannon in [4] is answered negatively. The argument will be based essentially upon work by Casler [5] and the author [1].

Some recent work by Eaton [6] includes a different proof that the solid Alexander horned sphere H , used in the example presented by Bass and Daverman, is not universal. This was done by sewing H to the crumpled cube T described by Stallings [7] so that the wild points of $\text{Bd } H$ are sewn to the Cantor set of nonpiercing points of T . Other methods developed by Eaton, in papers cited in [6], should offer alternative ways

Presented to the Society, October 30, 1971; received by the editors August 18, 1971.
AMS 1970 subject classifications. Primary 57A10.

Key words and phrases. Crumpled cube, upper semicontinuous decomposition, E^3 .

¹ Supported by Mississippi State University Biological and Physical Sciences Research Institute.

© American Mathematical Society 1972

to show that neither the decomposition space we describe here nor the one described by Bass and Daverman [2] is topologically S^3 .

Definitions relating to crumpled cubes will be those of [2].

1. **The example.** The description of the dogbone space of this paper is modeled after that by Bing in [3]. As in Figure 1, let H_1 and H_2 denote

FIGURE 1

the upper and lower half-spaces of S^3 , and P the xy -plane. Let A_0 denote a solid double torus which intersects P in a disk. Then, embed solid double tori $A_1, \dots, A_4$, linked as indicated, in A_0 ; although each of $A_1, \dots, A_4$ is shown as a finite graph, it is topologically equivalent to A_0 . Then, for each $i=1, \dots, 4$, solid double tori $A_{i,1}, \dots, A_{i,4}$ are embedded in A_i such that there is a homeomorphism of S^3 onto itself which is the identity on the complement of some open set containing A_0 and takes A_0 onto A_i . Succeeding steps of the construction are to be described inductively.

Let $M = A_0 \cap \sum A_i \cap \sum A_{i,j} \cap \sum A_{i,j,k} \dots$. Let G be the set whose elements are components of M and one-point subsets of $S^3 - M$. Then, G is an upper semicontinuous decomposition of S^3 into points and tame

arcs. Let S^3/G denote the associated decomposition space, the dogbone space of this note.

From [1], we have that S^3/G is not topologically S^3 .

The methods of Casler [5] can be easily modified to show that H_2^* , the natural projection of H_2 in S^3/G , is a self-universal crumpled cube.

Since H_1^* , the natural projection of H_1 in S^3/G , is a crumpled cube and S^3/G is the result of a sewing of H_1^* and H_2^* , we have that H_2^* is not a universal crumpled cube.

BIBLIOGRAPHY

1. E. H. Anderson, *Chains of simple closed curves and a dogbone space*, Fund. Math. **72** (1971), 131–144.
2. C. D. Bass and R. J. Daverman, *A self-universal crumpled cube which is not universal*, Bull. Amer. Math. Soc. **76** (1970), 740–742. MR **41** #7655.
3. R. H. Bing, *A decomposition of E^3 into points and tame arcs such that the decomposition space is topologically different from E^3* , Ann. of Math. (2) **65** (1957), 484–500. MR **19**, 1187.
4. C. E. Burgess and J. W. Cannon, *Embeddings of surfaces in E^3* , Rocky Mt. J. Math. **1** (1971), 259–344. MR **43** #4008.
5. B. G. Casler, *On the sum of two solid Alexander horned spheres*, Trans. Amer. Math. Soc. **116** (1965), 135–150. MR **32** #3049.
6. W. T. Eaton, *The sum of solid spheres*, Michigan Math. J. (to appear).
7. J. R. Stallings, *Uncountably many wild disks*, Ann. of Math. (2) **71** (1960), 185–186. MR **22** #1871.

DEPARTMENT OF MATHEMATICS, MISSISSIPPI STATE UNIVERSITY, STATE COLLEGE,
MISSISSIPPI 39762