THE COMPLETION OF A RING WITH A VALUATION

HELEN E. ADAMS

ABSTRACT. This paper proves three main results: the completion of a commutative ring with respect to a Manis valuation is an integral domain; a necessary and sufficient condition is given that the completion be a field; and the completion is a field when the valuation is Harrison and the value group is archimedean ordered.

1. Introduction. In [1] we defined a generalized pseudovaluation on a ring R and found explicitly the completion of R with respect to the topology induced on R by such a pseudovaluation. Using this inverse limit characterization of the completion of R we show, in §2, that when the codomain of a valuation on R is totally ordered, the completion of R with respect to the valuation has no (nonzero) divisors of zero and the valuation on R can be extended to a valuation on the completion. In §3, R is assumed to have an identity and so, from [1, §2], the completion has an identity: a necessary and sufficient condition is given such that each element of the completion of R has a right inverse.

§§2 and 3 are immediately applicable to a Manis valuation φ on a commutative ring R with identity [3], where the valuation φ is surjective and the set $\varphi(R)$ is a totally ordered group. A Harrison valuation on R is a Manis valuation φ on R such that the set $\{x: x \in R, \varphi(x) > \varphi(1)\}$ is a finite Harrison prime [2]. In §4 we show that the completion of a ring with respect to a Harrison valuation is a field if the value group is archimedean ordered.

Let R be a ring (not necessarily either commutative or with identity). Let S be a totally ordered quasi-residuated [1, §1] semigroup and let S_0 be the disjoint union of S and a zero element 0_S with the properties: $0_S 0_S = 0_S$; and, for any $s \in S$, $0_S > s$ and $s 0_S = 0_S = 0_S s$. Let $\varphi: R \rightarrow S_0$ be a valuation on R . That is, for all $a, b \in R$,

- (1.1) $\varphi(ab) = \varphi(a)\varphi(b)$;
- (1.2) $\varphi(a-b) \geq \min\{\varphi(a), \varphi(b)\}$;
- (1.3) $\varphi(0) = 0_S$;
- (1.4) the set $\varphi(R) \setminus \{0_S\}$ is nonempty.

Received by the editors October 19, 1971.

AMS 1970 subject classifications. Primary 12J20, 13J10; Secondary 16A80.

Key words and phrases. Valuation on a ring, valuated ring completion, Manis valuation, Harrison prime.

© American Mathematical Society 1972

Note that, if $\varphi(a) \neq \varphi(b)$, then

$$\varphi(a - b) = \min\{\varphi(a), \varphi(b)\}.$$

Now if S has a greatest element π , it is clear from [1] that the completion of R with respect to φ is isomorphic to the ring $R/\{x: x \in R, \varphi(x) \geq \pi\}$. Hence let S have no greatest element.

Following [1], let $\{P_s\}_{s \in S}$ be the generalized filtration defined on R as follows: For each $s \in S$, $P_s = \{x: x \in R, \varphi(x) \geq s\}$. Note that, in this case, $\bigcup_{s \in S} P_s = R$. From [1, §2], the completion of R with respect to φ is the topological ring $R = \text{proj lim } R/P_s$, with zero $0 = \{P_s\}_{s \in S}$, and addition, multiplication and topology defined in [1, §2]. For reference, $R = \{\{\xi_s\}_{s \in S} \in \prod_{s \in S} R/P_s: \text{for all } s, t \in S \text{ such that } s \geq t, \xi_s \subseteq \xi_t\}$. When there is no risk of ambiguity, $\{\xi_s\}_{s \in S}$ will be written as $\{\xi_s\}$.

2. R has no divisors of zero. First we need

LEMMA 1. Let $\{\xi_s\}_{s \in S} \in R \setminus \{0\}$. Then there exist $u, v \in S$ with $u < v$ such that $\xi_u = P_u$, $\xi_v \neq P_v$, and, for all $x, y \in \xi_v$, $u \leq \varphi(x) = \varphi(y) < v$.

PROOF. (i) Let $s \in S$, $x \in \xi_s$. Since $\bigcup_{s \in S} P_s = R$, there exists $t \in S$ such that $x \in P_t$. If $t \geq s$, then $x \in P_s$ from [1, (1.2)], and so $\xi_s = P_s$. Otherwise $t < s$, $\xi_s \subseteq \xi_t$, and so $\xi_t = x + P_t = P_t$. Hence there exists $u \in S$ such that $\xi_u = P_u$.

(ii) Since $\{\xi_s\}_{s \in S} \neq 0$, there exists $v \in S$ such that $\xi_v \neq P_v$. Now if $v \leq u$, then $P_u = \xi_u \subseteq \xi_v$, and so $\xi_v = P_v$. Therefore $v > u$.

(iii) Now $\varphi(x) < v$ for all $x \in \xi_v$, since otherwise $\xi_v = P_v$. Since $v > u$, $\xi_v \subseteq \xi_u = P_u$ and so $\varphi(x) \geq u$ for all $x \in \xi_v$. Suppose there exist $y, z \in \xi_v$ such that $\varphi(y) \neq \varphi(z)$. Then $\min\{\varphi(y), \varphi(z)\} = \varphi(y - z) \geq v$ since $y - z \in P_v$, which contradicts $\varphi(y), \varphi(z) < v$. Hence $\varphi(x) = \varphi(y)$ for all $x, y \in \xi_v$. This completes the proof.

THEOREM 1. R has no divisors of zero.

PROOF. Let $\{\xi_s\}, \{\eta_s\} \in R$ such that $\{\xi_s\}\{\eta_s\} = 0$. Suppose that $\{\xi_s\}, \{\eta_s\} \neq 0$. Then, by Lemma 1, there exist $a, b, t, u \in S$ such that, for all $x \in \xi_t$, $\varphi(x) = a$; and, for all $y \in \eta_u$, $\varphi(y) = b$.

Since S has no maximal element, there exists $c \in S$ such that $c > ab$; and, by the definition of multiplication in R [1, §2], there exists $v \in S$ such that, for all $x \in \xi_v$ and for all $y \in \eta_v$, $xy \in P_c$.

Let $w \geq t, u, v$. Then, for all $x \in \xi_w \subseteq \xi_t$ and for all $y \in \eta_w \subseteq \eta_u$, $\varphi(xy) = ab < c$, which contradicts $xy \in P_c$.

Hence $\{\xi_s\} = 0$ or $\{\eta_s\} = 0$ which completes the proof.

COROLLARY. *The completion of a commutative ring with respect to a nontrivial Manis valuation is an integral domain.*

Lemma 1 also enables us to define a valuation on R . Let $\{\xi_s\} \in R \setminus \{0\}$. Then there exists $t \in S$ such that $\xi_t \neq P_t$. Define $\varphi(\{\xi_s\}) = \varphi(x)$ where $x \in \xi_t$. Define $\varphi(0) = 0_S$. Then it is a straightforward task to show that $\varphi: R \rightarrow S_0$ is a well defined valuation on R . Moreover, the valuation φ on R is an "extension" of the valuation φ on R in the sense that φ restricted to the image $i(R)$ of R in R , defined in [1, Theorem 2.1], satisfies $\varphi(i(x)) = \varphi(x)$ for each $x \in R$. And it is easy to show that φ defines the topology $\mathcal{T}$ [1, §2] on R .

3. When elements of the completion have inverses. Let R have identity 1. Then, from [1, Proposition 2.1], R has identity $\{1 + P_s\}_{s \in S}$.

PROPOSITION 1. *Each element of R has a right inverse if and only if*
(1) *given $a \in R$ such that $\varphi(a) \neq 0_S$, and given $s \in S$, there exists $b \in R$ such that $\varphi(ab - 1) \geq s$.*

PROOF. (i) Let each element of R have a right inverse. Let $a \in R$ be such that $\varphi(a) \neq 0_S$, and let $w \in S$. Clearly, $\{a + P_s\} \neq 0$ since S has no maximal element. Hence there exists $\{\eta_s\} \in R$ such that $\{a + P_s\}\{\eta_s\} = \{1 + P_s\}$. By the definition of multiplication in R , there exists $t \in S$ such that, for all $x \in \eta_t$, $ax \in 1 + P_w$. Hence (1) holds.

(ii) Conversely, let (1) hold. Denote $\varphi(1)$ by 1_S . Let $\{\xi_s\} \in R \setminus \{0\}$. By Lemma 1 there exist $\rho, \sigma \in S$ such that $\xi_\rho \neq P_\rho$ and, for all $x \in \xi_\rho$, $\varphi(x) = \sigma$; σ is a constant for the chosen $\{\xi_s\}$.

For each $s \in S$, let $x_s \in \xi_s$ and let $y_s \in R$ be such that $\varphi(x_s y_s - 1) > \sigma s$ (S has no maximal element); let $m_s \in S$ be such that $m_s > \sigma s$ and let $g(s) \in S$ be such that $g(s) \geq m_s \sigma, s$. Let $t \in S$ be such that $\sigma t > 1_S$ and $t \geq \rho$. Then, for each $s \in S$ such that $s \geq t$, $\sigma \varphi(y_s) = \varphi(x_s y_s) = 1_S$. Define, for each $s \in S$,

$$\begin{aligned}\eta_s &= y_{g(s)} + P_s \quad \text{if } s \geq t, \\ &= y_{g(t)} + P_s \quad \text{if } s < t.\end{aligned}$$

We shall show that $\{\xi_s\}\{\eta_s\} = \{1 + P_s\}$.

(a) We must show that, for each $s \in S$, η_s is well defined. Accordingly, for each $s \in S$, let $w_s \in \xi_s$, $z_s \in R$ and $\tau, u, n_s, h(s) \in S$ be such that $\xi_\tau \neq P_\tau$; $\varphi(w_s z_s - 1) > \sigma s$; $n_s > \sigma s$ and $h(s) \geq n_s \sigma, s$; $\sigma u > 1_S$ and $u \geq \tau$; and

$$\begin{aligned}\zeta_s &= z_{h(s)} + P_s \quad \text{if } s \geq u, \\ &= z_{h(u)} + P_s \quad \text{if } s < u.\end{aligned}$$

We must show that, for each $s \in S$, $\zeta_s = \eta_s$.

CLAIM. Let $a, b, c \in S$ be such that $a, b \geq c \geq t$. Then $z_{h(a)} - y_{g(b)} \in P_c$.

SUBPROOF. We shall refer to $h(a)$ and $g(b)$ as h and g respectively. Now

$$w_h(z_h - y_g) = (w_h z_h - x_g y_g) + (x_g - w_h) y_g.$$

Clearly, $\varphi\{w_h z_h - 1 - (x_g y_g - 1)\} > \sigma c$. Also $\varphi(x_g - w_h) \geq r\sigma$ where $r = \min\{m_b, n_a\} > \sigma c$. Hence $\varphi\{(x_g - w_h) y_g\} > \sigma c$ since $\sigma\varphi(y_g) = 1_S$, and so $\varphi\{w_h(z_h - y_g)\} > \sigma c$. Thus $\varphi(z_h - y_g) \geq c$ since $\varphi(w_h) = \sigma$. This proves the claim.

Let $s \in S$. Without loss of generality, let $u \geq t$. Then, by applying the claim to each of the cases $s \geq u$, $u \geq s \geq t$, $t > s$, we see that $\zeta_s = \eta_s$. Thus η_s is well defined for each $s \in S$.

(b) By further application of the claim, it is easy to show that $\{\eta_s\}_{s \in S} \in R$.

(c) Finally we must show that $\{\xi_s\}\{\eta_s\} = \{1 + P_s\}$. Let $\{\xi_s\}\{\eta_s\} = \{\Omega_s\}$. Let $s \in S$. Then there exists $v \in S$ such that $xy \in \Omega_s$ for all $x \in \xi_v$, $y \in \eta_v$. Let $d \in S$ be such that $\sigma d \geq s$ and $d \geq v$, t . Then $x_{g(d)} \in \xi_v$, $y_{g(d)} \in \eta_v$ and $\varphi(x_{g(d)} y_{g(d)} - 1) \geq s$. Hence $\Omega_s = x_{g(d)} y_{g(d)} + P_s = 1 + P_s$ which completes the proof of the proposition.

COROLLARY. Let $\varphi: R \rightarrow S_0$ be a nontrivial Manis valuation on a commutative ring R with identity 1. Then the completion of R with respect to φ is a field if and only if condition (1) holds.

4. Harrison valuations. In [2, pp. 11 and 12 and Proposition 2.2, p. 14], Harrison shows that, given a finite prime P of a commutative ring R with identity, one can define a Manis valuation $\varphi_P: R \rightarrow \Gamma_P$ on R such that the set $\{x: x \in R, \varphi_P(x) > \varphi_P(1)\} = P$. Using the corollary to Proposition 1, we have:

THEOREM 2. Let $\varphi: R \rightarrow S_0$ be a nontrivial Harrison valuation on a commutative ring R with identity 1_R , such that S be archimedean ordered. Then the completion of R with respect to φ is a field.

PROOF. Let $a \in R$ be such that $\varphi(a) \neq 0_S$ and let $s \in S$. Now $\{x: x \in R, \varphi(x) > 1_S\} = P$, a finite Harrison prime of R . Suppose $aR \subseteq P$. Then, for all $x \in R$, $\varphi(a)\varphi(x) > 1_S$, which contradicts $\varphi(a) \neq 0_S$ since φ is surjective and S is a totally ordered group. Hence $aR + P$ properly contains P . But $aR + P$ is closed under addition and multiplication. Thus $-1_R \in aR + P$, since P is a Harrison prime: that is, there exist $c \in R$, $d \in P$ such that $ac - 1_R = d$.

Since $\varphi(d) > 1_S$ and S is archimedean ordered, there exists a natural number m such that $\varphi(d^m) = \{\varphi(d)\}^m \geq s$. Let n be an odd integer such that $n \geq m$. Then $\varphi(d^n) \geq s$. Now $d^n = (ac - 1_R)^n = ab - 1_R$ where $b \in R$. The theorem then follows from Proposition 1.

REFERENCES

1. Helen E. Adams, *Filtrations and valuations on rings*, Pacific J. Math. **41** (1972).
2. D. K. Harrison, *Finite and infinite primes for rings and fields*, Mem. Amer. Math. Soc. No. 68 (1966), 62 pp. MR **34** #7550.
3. Merle E. Manis, *Valuations on a commutative ring*, Proc. Amer. Math. Soc. **20** (1969), 193–198. MR **38** #2134.

DEPARTMENT OF MATHEMATICS, MOUNT HOLYOKE COLLEGE, SOUTH HADLEY, MASSACHUSETTS 01075