PERTURBATIONS OF LINEAR m -ACCRETIVE OPERATORS

NOBORU OKAZAWA¹

ABSTRACT. A sufficient condition is given for the sum $A+B$ of two linear m -accretive operators A and B in a Hilbert space to be m -accretive. This condition is expressed in terms of $\operatorname{Re}(Au, Bu)$ for u in D , where D is a certain linear manifold contained in $D(A+B)$.

1. Introduction. A linear operator A (with domain $D(A)$ and range $R(A)$) in a (complex) Banach space X is said to be accretive² if

$$(A) \quad \|(A + \xi)u\| \geq \xi \|u\| \quad \text{for every } u \in D(A) \text{ and } \xi > 0.$$

It is known that $R(A + \xi) = X$ either for every $\xi > 0$ or for no $\xi > 0$; in the former case we say that A is m -accretive. Let X^* be the adjoint space of X . The pairing between $w \in X$ and $f \in X^*$ will be denoted by (w, f) . Let F be the duality map from X to X^* : $F(w) = \{f \in X^*; (w, f) = \|w\|^2 = \|f\|^2\}$ for every $w \in X$. Then (A) is equivalent to the following condition:

$$(A') \quad \text{For every } u \in D(A) \text{ there is } f \in F(u) \text{ such that } \operatorname{Re}(Au, f) \geq 0;$$

see [6] (in which the term "monotonic" was used instead of "accretive"). Note that the inequality is not required to hold for every $f \in F(u)$. But if A is m -accretive and densely defined, then the inequality holds for every $f \in F(u)$ (cf. [8, Remark 1 to Theorem 3.1]).

Now let X be reflexive. Then linear m -accretive operators in X are necessarily densely defined (see [12, p. 218]). So if A and B are linear m -accretive operators in X , then $A+B$ defined on $D(A+B) = D(A) \cap D(B)$ is always accretive according to condition (A').

The main purpose of this note is to give a sufficient condition for the sum of two linear m -accretive operators in a Hilbert space to be m -accretive (§3). In particular we shall improve a perturbation theorem of Glimm

Received by the editors May 23, 1972.

AMS (MOS) subject classifications (1970). Primary 47A55, 47B44, 47B25; Secondary 47D05.

Key words and phrases. Singular perturbations, accretive (dissipative) operators, Yosida approximations, cores, selfadjoint operators, regular (relatively bounded) perturbations.

¹ The writer would like to thank the referee for his kind suggestions.

² In this case $-A$ is said to be dissipative.

and Jaffe [3]. §2 contains a remark for (reflexive) Banach space case. We finally consider the relatively bounded perturbation of linear m -accretive operators in a Hilbert space (§4).

The writer wishes to thank Professor R. Iino for constant encouragement and Professor I. Miyadera for kind advice.

2. Singular perturbation in a reflexive Banach space. Let A and B be linear m -accretive operators in a Banach space X and A_n be the Yosida approximation of A : $A_n = A(1 + n^{-1}A)^{-1} = n(1 - (1 + n^{-1}A)^{-1})$, $n = 1, 2, \dots$; A_n is a bounded linear operator on X . Since A_n is also accretive (see [6, Lemma 2.3]), it is easily seen that $A_n + B$ defined on $D(B)$ is m -accretive. Consequently, for every $v \in X$ there exists a unique $u_n \in D(B)$ such that

$$(2.1) \quad A_n u_n + B u_n + u_n = v, \quad n = 1, 2, \dots$$

LEMMA 2.1. *Let X be reflexive and A be m -accretive in X .*

(a) *If $\{u_n\}$ is a sequence in $D(A)$ such that $u_n \rightharpoonup u \in X$ (we denote by $\rightharpoonup$ weak convergence) and if the $\|Au_n\|$ are bounded, then $u \in D(A)$ and $Au_n \rightharpoonup Au$.*

(b) *If $\{w_n\}$ is a sequence in X such that $w_n \rightharpoonup u \in X$ and if the $\|A_n w_n\|$ are bounded, then $u \in D(A)$ and $A_n w_n \rightharpoonup Au$.*

PROOF. (a) Since X is reflexive and the $\|Au_n\|$ are bounded, there is a subsequence $\{u_m\}$ of $\{u_n\}$ such that $Au_m \rightharpoonup v \in X$. Since A is closed, its graph is weakly closed so that $u \in D(A)$ and $v = Au$.

Since we could have started with any subsequence of $\{u_n\}$ instead of $\{u_n\}$ itself, the result obtained shows that $Au_n \rightharpoonup Au$.

(b) Set $u_n = (1 + n^{-1}A)^{-1}w_n \in D(A)$. Then $Au_n = A_n w_n$ and the $\|Au_n\|$ are bounded. Also $w_n - u_n = (1 - (1 + n^{-1}A)^{-1})w_n = n^{-1}A_n w_n \rightarrow 0$ (we denote by $\rightarrow$ strong convergence) so that $u_n \rightarrow u$. Thus the result of (a) is applicable: $u \in D(A)$ and $A_n w_n = Au_n \rightharpoonup Au$. Q.E.D.

PROPOSITION 2.2. *Let X be reflexive. Let A and B be linear m -accretive operators in X . If $\|A_n(A_n + B + 1)^{-1}\|$ is bounded as n tends to infinity, then $A + B$ is m -accretive.*

PROOF. Since $A + B$ is accretive, it suffices to show that $R(A + B + 1) = X$. To see this, let u_n be the solution of equation (2.1). Since $\|u_n\| \leq \|v\|$, there is a subsequence $\{u_m\}$ of $\{u_n\}$ such that $u_m \rightharpoonup u \in X$. Noting that $\|A_m u_m\|$ and $\|B u_m\|$ are bounded in m by assumption, we see from Lemma 2.1 that $u \in D(A) \cap D(B)$, $B u_m \rightharpoonup B u$ and $A_m u_m \rightharpoonup A u$. Thus we obtain $v = (A + B + 1)u$ which implies $R(A + B + 1) = X$. Q.E.D.

REMARK 2.3. When X^* is uniformly convex, Proposition 2.2 holds even if A and B are nonlinear (see [7, Lemma 1]).

3. Singular perturbation in a Hilbert space. The following lemma is a linear version of a recent result of Brezis, Crandall and Pazy (see [1, Theorem 2.1]):

LEMMA 3.1. *Let A and B be linear m -accretive operators in a Hilbert space H . Then $R(A+B+1)=H$ if and only if $\|A_n(A_n+B+1)^{-1}\|$ is bounded as n tends to infinity.*

Note that the 'if' part of Lemma 3.1 is a particular case of Proposition 2.2. We shall use Lemma 3.1 in the following form:

LEMMA 3.2. *Let A and B be as in Lemma 3.1. Then $R(A+B+1)=H$ if and only if there are nonnegative constants a and $b < 1$ such that*

$$(3.1) \quad 0 \leq 2 \operatorname{Re}(A_n u, Bu) + a \|u\|^2 + b(\|A_n u\|^2 + \|Bu\|^2), \quad u \in D(B).$$

PROOF. It is easily seen that (3.1) is equivalent to

$$(3.2) \quad \|A_n u\|^2 + \|Bu\|^2 \leq c(\|(A_n + B)u\|^2 + \|u\|^2), \quad u \in D(B), \quad c \geq 1.$$

Obviously, (3.2) implies that $\|A_n(A_n+B+1)^{-1}\|$ is bounded in n . Thus it suffices by Lemma 3.1 to show that if $\|A_n(A_n+B+1)^{-1}\|$ is bounded in n , then (3.2) holds. But since $\|B(A_n+B+1)^{-1}\|$ is also bounded in n , there exists a constant $c_1 > 0$ such that

$$(3.3) \quad \|A_n(A_n+B+1)^{-1}v\|^2 + \|B(A_n+B+1)^{-1}v\|^2 \leq c_1 \|v\|^2, \quad v \in H.$$

Now let $u \in D(B)$. Setting $v = (A_n+B+1)u$ in (3.3), we can obtain (3.2).
Q.E.D.

To simplify the statement, let us introduce the following

DEFINITION 3.3. A linear manifold D contained in the domain of a closed linear operator A is called a core of A if the closure of the restriction of A to D is again A .

Then our result is given by

THEOREM 3.4. *Let A and B be linear m -accretive operators in H . Let D be a linear manifold contained in $D(A)$ such that D is mapped into itself by $(1+n^{-1}A)^{-1}$ for $n=1, 2, \dots$. Assume that D is contained in $D(B)$ and there exist nonnegative constants a and $b < 1$ such that for all $u \in D$,*

$$(3.4) \quad 0 \leq 2 \operatorname{Re}(Au, Bu) + a \|u\|^2 + b \|Au\|^2.$$

If D is a core of B , then $A+B$ is also m -accretive.

PROOF. Since $A+B$ is accretive, it suffices to show that $R(A+B+1)=H$. To see this, set $D_0 = (1+A)^{-1}D$. Then D_0 is contained in D and

$\operatorname{Re}(Au, BAu) \geq 0$ for $u \in D_0$. So, we see from (3.4) that for all $u \in D_0$,

$$(3.5) \quad 0 \leq 2 \operatorname{Re}(Au, B(1 + n^{-1}A)u) + a \|u\|^2 + b \|Au\|^2, \quad n = 1, 2, \dots$$

Now let $v \in D$. Then $(1 + n^{-1}A)^{-1}v \in D_0$ (use the resolvent equation). Setting $u = (1 + n^{-1}A)^{-1}v$ in (3.5), we have that for all $v \in D$,

$$0 \leq 2 \operatorname{Re}(A_n v, Bv) + a \|v\|^2 + b \|A_n v\|^2.$$

Since D is a core of B , it follows that for all $u \in D(B)$, $0 \leq 2 \operatorname{Re}(A_n u, Bu) + a \|u\|^2 + b \|A_n u\|^2$. Thus, we see by Lemma 3.2 that $R(A + B + 1) = H$.

Q.E.D.

Now set $D(A^\infty) = \bigcap_n D(A^n)$. Then $D(A^\infty)$ can be taken as D in Theorem 3.4 and we have

COROLLARY 3.5. *Let A and B be selfadjoint operators in H satisfying the inequality (3.4) with $b < 1$ and $u \in D(A^\infty)$. If $D(A^\infty)$ is a core of B , then $A + B$ is also selfadjoint.*

PROOF. The selfadjointness of A and B implies that $\pm iA$ and $\pm iB$ are m -accretive. Since $(\pm iAu, \pm iBu) = (Au, Bu)$ for $u \in D(A^\infty)$, it follows from Theorem 3.4 that $\pm i(A + B)$ are m -accretive. But, $A + B$ is symmetric (note that $D(A^\infty)$ is dense in H) and so $A + B$ is selfadjoint. Q.E.D.

REMARK 3.6. Let A be a positive selfadjoint operator. Let B be a selfadjoint operator such that $D(A^\infty)$ is a core of B and $BD(A^\infty)$ is again contained in $D(A)$. Assume that there are nonnegative constants ε , α and β such that for $u \in D(A^\infty)$, $0 \leq \alpha(Au, u) + (Bu, u) + \beta \|u\|^2$ and

$$0 \leq \varepsilon \|Au\|^2 + (Bu, u) + ([A^{1/2}, [A^{1/2}, B]]u, u) + \beta \|u\|^3, \quad 2\alpha + \varepsilon < 1,$$

where $[T, S] = TS - ST$ and $A^{1/2}$ is the square root of A . Then, since $[A^{1/2}, [A^{1/2}, B]] = [A^{1/2}, [A^{1/2}, B + \beta]]$, we have

$$0 \leq 2 \operatorname{Re}((A + \tfrac{1}{2})u, (B + \beta)u) + (2\alpha + \varepsilon) \|(A + \tfrac{1}{2})u\|^2, \quad u \in D(A^\infty);$$

note that $\|Au\|^2 \leq \|Au\|^2 + (Au, u) + \|u\|^2/4 = \|(A + \tfrac{1}{2})u\|^2$. Consequently, it follows from Corollary 3.5 that $A + B$ is selfadjoint. This fact is proved by Glimm and Jaffe under additional assumptions (see [3, Theorem 8]).

4. Regular perturbation in a Hilbert space. For the regular (relatively bounded) perturbation of linear m -accretive operators in a general Banach space see e.g. [2], [4], [5] and [10]. The following theorem is a variant of the result noted in [9]:

THEOREM 4.1. *Let A be a linear m -accretive operator in a Hilbert space H . Let B be a linear accretive operator in H with $D(B) \supset D(A)$. Assume that*

there are nonnegative constants a and $b \leq 1$ such that

$$(4.1) \quad 0 \leq \operatorname{Re}(Au, Bu) + a \|u\|^2 + b \|Au\|^2, \quad u \in D(A).$$

If $b < 1$, then $A+B$ is also m -accretive. If $b=1$, then the closure of $A+B$ is m -accretive.

Note that (4.1) is similar to the inequality (3.4). First we prove

LEMMA 4.2. If $b < 1$ in Theorem 4.1, then $A+B$ is closed.

PROOF. Since $D(A)$ is dense in H (as noted in §1), so is $D(B)$ and therefore B is closable (see [5, Theorem V-3.4]). Consequently, there exists a constant $c_1 > 0$ such that

$$(4.2) \quad \|Bu\| \leq c_1(\|u\| + \|Au\|), \quad u \in D(A)$$

(see [5, Remark IV-1.5]). Also, we obtain from (4.1)

$$\begin{aligned} 0 &\leq \operatorname{Re}(Au, Bu) + b \|(A+c)u\|^2 \\ &\leq \operatorname{Re}((A+c)u, (A+B+c)u) + (b-1) \|(A+c)u\|^2, \quad c = (a/b)^{1/2}. \end{aligned}$$

Hence it follows that

$$(4.3) \quad (1-b) \|(A+c)u\| \leq \|(A+B+c)u\|, \quad u \in D(A).$$

We see from (4.2) and (4.3) that $A+B$ is closed. Q.E.D.

PROOF OF THEOREM 4.1. It suffices by Lemma 4.2 to show that if $b \leq 1$, then the closure of $A+B$ is m -accretive. Since $A+B$ is densely defined, it suffices to show that the adjoint $(A+B)^*$ of $A+B$ is accretive (cf. [12, p. 251]).

It follows from (4.1) that $0 \leq \operatorname{Re}(Au, (A+B)u) + a\|u\|^2$ for $u \in D(A)$. Hence we obtain

$$(4.4) \quad 0 \leq \operatorname{Re}((1+n^{-1}A)u, (A+B)u) + n^{-1}a\|u\|^2, \quad n = 1, 2, \dots$$

Now let $v \in H$. Then $(1+n^{-1}A)^{-1}v \in D(A)$. Setting $u = (1+n^{-1}A)^{-1}v$ in (4.4), we have that for all $v \in H$,

$$(4.5) \quad 0 \leq \operatorname{Re}(v, (A+B)(1+n^{-1}A)^{-1}v) + n^{-1}a\|(1+n^{-1}A)^{-1}v\|^2.$$

Let $w \in D((A+B)^*)$. Then it follows from (4.5) that

$$0 \leq \operatorname{Re}((A+B)^*w, (1+n^{-1}A)^{-1}w) + n^{-1}a\|w\|^2, \quad n = 1, 2, \dots$$

Going to the limit $n \rightarrow \infty$, we see that $(A+B)^*$ is accretive. Q.E.D.

REMARK 4.3. Theorem 4.1 is essentially proved by Yoshikawa though the closedness of B is assumed in his statement (see [11, Theorem 3.10 and Remark 3.12]).

REMARK 4.4. Let A and B be as in Theorem 4.1. Suppose that there are nonnegative constants a_1 and b_1 such that $\|Bu\|^2 \leq a_1\|u\|^2 + b_1\|Au\|^2$ for $u \in D(A)$. If $b_1 \leq 1$, then (4.1) holds with $b \leq 1$. In fact, setting $a = a_1/2$ and $b = (b_1 + 1)/2$, we have that for $u \in D(A)$,

$$\begin{aligned} \|Bu\|^2 &\leq 2a\|u\|^2 + (2b - 1)\|Au\|^2 \\ &\leq \|(A + B)u\|^2 - \|Au\|^2 + 2a\|u\|^2 + 2b\|Au\|^2 \\ &= \|Bu\|^2 + 2(\operatorname{Re}(Au, Bu) + a\|u\|^2 + b\|Au\|^2). \end{aligned}$$

Thus, Corollary 1 to Theorem 1 and Theorem 2 of [9] are covered by Theorem 4.1.

REFERENCES

1. H. Brezis, M. G. Crandall and A. Pazy, *Perturbations of nonlinear maximal monotone sets in Banach space*, Comm. Pure Appl. Math. **23** (1970), 123–144. MR **41** #2454.
2. P. R. Chernoff, *Perturbations of dissipative operators with relative bound one*, Proc. Amer. Math. Soc. **33** (1972), 72–74.
3. J. Glimm and A. Jaffe, *Singular perturbations of selfadjoint operators*, Comm. Pure Appl. Math. **22** (1969), 401–414. MR **43** #7955.
4. K. Gustafson, *A perturbation lemma*, Bull. Amer. Math. Soc. **72** (1966), 334–338. MR **32** #4555.
5. T. Kato, *Perturbation theory for linear operators*, Die Grundlehren der math. Wissenschaften, Band 132, Springer-Verlag, Berlin and New York, 1966. MR **34** #3324.
6. ———, *Nonlinear semigroups and evolution equations*, J. Math. Soc. Japan **19** (1967), 508–520. MR **37** #1820.
7. Y. Konishi, *A remark on perturbation of m -accretive operators in Banach space*, Proc. Japan Acad. **47** (1971), 452–455.
8. G. Lumer and R. S. Phillips, *Dissipative operators in a Banach space*, Pacific J. Math. **11** (1961), 679–698. MR **24** #A2248.
9. N. Okazawa, *Two perturbation theorems for contraction semigroups in a Hilbert space*, Proc. Japan Acad. **45** (1969), 850–853. MR **42** #2312.
10. ———, *A perturbation theorem for linear contraction semigroups on reflexive Banach spaces*, Proc. Japan Acad. **47** (1971), 947–949.
11. A. Yoshikawa, *On perturbation of closed operators in a Banach space*, J. Fac. Sci. Hokkaido Univ. **22** (1972), 50–61.
12. K. Yosida, *Functional analysis*, 2nd ed., Die Grundlehren der math. Wissenschaften, Band 123, Springer-Verlag, Berlin and New York, 1968. MR **39** #741.

DEPARTMENT OF APPLIED PHYSICS, WASEDA UNIVERSITY, NISHIOKUBO, SHINJUKU-KU, TOKYO, JAPAN