NILPOTENT ELEMENTS IN BANACH ALGEBRAS

H. BEHNCKE

ABSTRACT. Let $\mathfrak{A}$ be an A^* -algebra such that any maximal abelian $*$ -subalgebra is regular and such that any quasinilpotent element x in $\mathfrak{A}$ satisfies $x^N=0$, with $N<\infty$. Then any irreducible Hilbert space $*$ -representation is at most N -dimensional. If $\mathfrak{A}$ is a C^* -algebra, $\mathfrak{A}$ possesses transcendental quasinilpotent elements if there exists a $\pi \in \hat{\mathfrak{A}}$ with $\dim \pi = \infty$.

This note is a continuation of [1]. We want to establish an intimate connection between the degree of noncommutativity of a Banach algebra $\mathfrak{A}$ and the existence of nilpotent elements in $\mathfrak{A}$. The proofs are straightforward extensions of the matrix situation and of methods used in [1].

Throughout all Banach algebras are complex Banach algebras with norm $|\cdot|$, continuous involution $*$ and an auxiliary norm [3] $|\cdot|_0$ which satisfies $|xx^*|_0 = |x|_0^2$. Such Banach algebras are called A^* -algebras [3]. In A^* -algebras maximal abelian $*$ -subalgebras are necessarily semisimple. We shall further assume that all Banach algebras have a unit, because the adjunction of a unit does not affect the proofs nor the results. A representation π always denotes a $*$ -representation on a Hilbert space H_π . We say that $x \in \mathfrak{A}$ is n -nilpotent, with $n = \infty, 1, 2, \dots$, if x is quasinilpotent and $x^n=0$ but $x^{n-1} \neq 0$.

We begin with a simple lemma, which is probably known.

LEMMA. *Let $\mathfrak{A}$ be an A^* -algebra such that any $x=x^* \in \mathfrak{A}$ has a finite spectrum with at most n points; then $\dim \mathfrak{A} \leq n^2$.*

PROOF. Let $\mathfrak{B}$ be a maximal abelian (semisimple) $*$ -subalgebra of $\mathfrak{A}$. Then the character space $\mathfrak{B}$ of $\mathfrak{B}$ has at most n elements. Hence any $x \in \mathfrak{B}$ has the form $x = \sum_{i \leq n} \lambda_i e_i$, where the e_i are the minimal idempotents of $\mathfrak{B}$. The e_i are selfadjoint and it is easy to see that $\dim e_i \mathfrak{A} e_j \leq 1$.

THEOREM 1. (i) *Let $\mathfrak{A}$ be an A^* -algebra such that any maximal abelian $*$ -subalgebra is regular and such that any quasinilpotent element x in $\mathfrak{A}$ satisfies $x^N=0$, then any irreducible representation π of $\mathfrak{A}$ is at most N dimensional.*

Received by the editors February 10, 1972.

AMS (MOS) subject classifications (1969). Primary 4660, 4665.

Key words and phrases. Nilpotent elements, Banach $*$ -algebras, C^* -algebras.

© American Mathematical Society 1973

(ii) Let $\mathfrak{A}$ be an A^* -algebra such that any maximal abelian $*$ -subalgebra of $\mathfrak{A}$ is regular and assume $\mathfrak{A}$ has an irreducible representation π of dimension $1 < N < \infty$, then $\mathfrak{A}$ has an N -nilpotent element.

(iii) Let $\mathfrak{A}$ be an A^* -algebra such that any irreducible representation of $\mathfrak{A}$ is at most N -dimensional; then any quasinilpotent element x in $\mathfrak{A}$ satisfies $x^N = 0$.

PROOF. (i) Let π be an irreducible representation of $\mathfrak{A}$ and let $\mathfrak{B}$ be a maximal abelian $*$ -subalgebra of $\mathfrak{A}$. By ρ we denote the restriction of π to $\mathfrak{B}$. Then the hull $h(J)$ [3] of $J = \ker \rho$ is a closed set in $\hat{\mathfrak{B}}$. Assume we can find $N+1$ distinct points $t_1, \dots, t_{N+1} \in h(J)$. Then there exist disjoint open neighborhoods U_i of t_i in $\hat{\mathfrak{B}}$. Since $\mathfrak{B}$ is regular we can find elements $a_i \in \mathfrak{B}$ with $\hat{a}_i(t_i) = 1$ and $\hat{a}_i(t) = 0$ for all $t \notin U_i$ with $1 \leq i \leq N+1$. Here $\hat{}$ denotes the Gelfand transform. Then $a_i a_j = a_j a_i = 0$ for $i \neq j$ and $a_i \notin \ker \pi$. Let $c_1, \dots, c_N$ be arbitrary in $\mathfrak{A}$, then the element $a = a_1 c_1 a_2 + \dots + a_N c_N a_{N+1}$ satisfies $a^{N+1} = 0$. By assumption $a^N = a_1 c_1 a_2^2 \cdots a_N^2 c_N a_{N+1} = 0$. This holds for all possible choices of c_i . Since $\pi(a_{N+1}) \neq 0$ there exists a $\xi \in H_\pi$ with $\pi(a_{N+1})\xi = \eta \neq 0$. Then by assumption

$$\pi(a_1 c_1 a_2^2 \cdots a_N^2) \pi(c_N) \eta = 0.$$

Keeping the $c_1, \dots, c_{N-1}$ fixed for a moment, this holds for all c_N . However since π is irreducible the set $\{\pi(c_N)\eta \mid c_N \in \mathfrak{A}\}$ is dense in H_π . Hence $\pi(a_1 c_1 \cdots a_N^2) = 0$ and this holds for any $(N-1)$ -tuple $(c_1, \dots, c_{N-1})$. Continuation of this argument finally leads to $\pi(a_i) = 0$, a contradiction. Hence any selfadjoint element in $\pi(\mathfrak{A})$ has at most N points in its spectrum. By the lemma we get $\dim \pi \leq N$.

(ii) Let π be an irreducible representation of dimension N , and let $e_{i,j}$, with $1 \leq i, j \leq N$, be a system of matrix units in $\pi(\mathfrak{A})$. As in (i) we can find elements $a_i \in \mathfrak{A}$ such that $a_i a_j = a_j a_i = 0$ for $i \neq j$ and $\pi(a_i) = e_{i,i}$. Further let $c_i \in \pi^{-1}(e_{i,i+1})$ then $a = a_1 c_1 a_2 + \dots + a_{N-1} c_{N-1} a_N$ satisfies $a^N = 0$ and

$$\pi(a^{N-1}) = \pi(a_1 c_1 a_2^2 \cdots a_{N-1}^2 c_{N-1} a_N) = e_{1,1} e_{1,2} e_{2,2}^2 \cdots e_{N,N} = e_{1,N} \neq 0.$$

(iii) Assume $\mathfrak{A}$ has a quasinilpotent element x with $x^N \neq 0$.

Then there exists an irreducible representation π with $\pi(x^N) \neq 0$. Since $\dim \pi \leq N$ this is impossible.

For $N=1$ the theorem may be stated in slightly weaker form.

COROLLARY. Let $\mathfrak{A}$ be an A^* -algebra with normal generators $\{x_i\}$ such that the abelian $*$ -subalgebras $\mathfrak{B}_i$ generated by x_i and 1 are regular. Then $\mathfrak{A}$ is commutative iff $\mathfrak{A}$ has no nilpotent elements.

PROOF. Assume $\mathfrak{A}$ has no nilpotent elements and let π be an irreducible representation of $\mathfrak{A}$. Then arguing as above we see that $\text{Sp}(\pi x_i)$ consists of

only one point. Hence $\pi(x_i) = \lambda_i \mathbf{1}$ and $\dim \pi = 1$. Thus $\mathfrak{A}$ is commutative. The converse is trivial, because commutative A^* -algebras are semisimple.

The corollary applies in particular to l^1 -algebras of discrete groups [1].

Part (ii) of Theorem 1 can be extended to $n = \infty$ for C^* -algebras.

THEOREM 2. *Let $\mathfrak{A}$ be a C^* -algebra such that there is a $\pi \in \hat{\mathfrak{A}}$ with $\dim \pi = \infty$. Then $\mathfrak{A}$ has an ∞ -nilpotent element.*

PROOF. Again there is no loss of generality to assume that $\mathfrak{A}$ has a unit. We assume further that $\mathfrak{A}$ has no ∞ -nilpotent elements and derive a contradiction. Let $\pi \in \hat{\mathfrak{A}}$ be an infinite dimensional representation and assume there is a $b = b^* \in \pi(\mathfrak{A})$ such that the spectrum of b contains an interval $[\alpha, \beta]$ with $\beta > \alpha$. Using the functional calculus of C^* -algebras we find an $a = a^* \in \pi(\mathfrak{A})$ with $\text{Sp } a = [0, 1]$. Let $c \in \mathfrak{A}$ be positive hermitean with $\pi(c) = a$ and let f_i be the function.

$$\begin{aligned} f_i(t) &= 0, & t &\leq \frac{1}{i+1}, \\ &= \text{linear}, & \frac{1}{i+1} &\leq t \leq \frac{3i+1}{3i(i+1)}, \\ &= \frac{1}{i^2}, & \frac{3i+1}{3i(i+1)} &\leq t \leq \frac{3i+2}{(i+1)i3}, \\ &= \text{linear}, & \frac{3i+2}{3i(i+1)} &\leq t \leq \frac{1}{i}, \\ &= 0, & t &\geq \frac{1}{i}. \end{aligned}$$

Then the elements $a_i = f_i(c)$ satisfy $a_i a_j = 0$ for $i \neq j$ and $\pi(a_i) = f_i(a) \neq 0$. Now let $q = \sum_{i=1}^{\infty} a_i c_i a_{i+1}$ with $c_i \in \mathfrak{A}$ and $|c_i| \leq 2$. Then

$$q^k = \sum_{i=1}^{\infty} a_i c_i a_{i+1}^2 \cdots c_{i+k-1} a_{i+k}$$

and

$$|q^k| \leq \sum_{i=1}^{\infty} \frac{2^k}{i^2(i+1)^4 \cdots (i+k)^2} \leq \sum_{i=1}^{\infty} \frac{1}{i(i+1) \cdots (i+k)} = \frac{1}{kk!}.$$

Hence q is quasinilpotent for all possible choices of c_i . We shall show now that we can find $c_i \in \mathfrak{A}$ such that $|c_i| \leq 2$ and such that $\pi(q^n) \neq 0$ for all n .

By construction of the a_i there exist unit vectors $\xi_i \in H_\pi$ with $\pi(a_i)\xi_i = (1/i^2)\xi_i$. Since π is irreducible there exist $c_i \in \pi(\mathfrak{A})$ with $|c_i| \leq 2$ and $\pi(c_i)\xi_{i+1} = \xi_i$ [2, 2.8.3]. Let $\{c_i\}$ be a fixed sequence of such operators and

assume that the corresponding q satisfies $q^k=0$. Then

$$\begin{aligned} 0 &= \pi(q^k)\xi_{k+1} = \pi(a_1 \cdots a_k^2)\pi(c_k)\pi(a_{k+1})\xi_{k+1} = \cdots \\ &= \frac{1}{(k+1)^2} \frac{1}{k^4} \cdots \frac{1}{2^4} \xi_1 \neq 0 \quad \text{a contradiction.} \end{aligned}$$

The case when every maximal abelian subalgebra $\mathfrak{B}$ of $\pi(\mathfrak{A})$ has a totally disconnected spectrum can be treated similarly, because $\dim \pi = \infty$.

This theorem is an extension of a result of Topping [4], who showed that any antiliminal C^* -algebra possesses ∞ -nilpotent elements. Our methods however are different and more direct. Theorem 2 shows that any C^* -algebra without ∞ -nilpotent elements is necessarily liminal. This result is not optimal. To see this, let $\mathfrak{A}$ be a weak direct sum of liminal C^* -algebras $\mathfrak{A}_i$, $\mathfrak{A} = \sum' \oplus \mathfrak{A}_i$, such that any $\pi \in \hat{\mathfrak{A}}_i$ satisfies $\dim \pi < \infty$, but such that there exists a $\pi_i \in \hat{\mathfrak{A}}_i$ with $\dim \pi_i = p_i$ and $p_i \rightarrow \infty$. Then by Theorem 1 (ii) there exists a p_i -nilpotent element $a_i \in \mathfrak{A}_i$ with $|a_i| = 1$. Then $a = \sum \oplus a_i \cdot 1/i$ is an ∞ -nilpotent element. This example leads immediately to our next theorem.

THEOREM 3. *Let $\mathfrak{A}$ be a separable C^* -algebra without ∞ -nilpotent elements. Then there exists a positive integer N such that any quasinilpotent element $x \in \mathfrak{A}$ satisfies $x^N = 0$.*

PROOF. (a) By Theorem 2 we may assume $\mathfrak{A}$ to be liminal. Suppose further that $\mathfrak{A}$ has irreducible representations of arbitrary high but finite dimensions. Determine now as in [2, 4.4.4] an open set $\mathfrak{D}'_1 \subset \hat{\mathfrak{A}}$ with the properties a , b and c of [2, 4.4.4]. Then there exist some positive integer n_1 and a nontrivial open subset $\mathfrak{D}_1 \subset \mathfrak{D}'_1 \cap \cap_{n=1}^{n_1} \hat{\mathfrak{A}}$ [2, 3.6.3] or any nontrivial open subset of $\mathfrak{D}'_1$ contains representations of arbitrary high dimensions. Similarly determine an open subset $\mathfrak{D}'_2$ in $\hat{\mathfrak{A}} \setminus \cap_{n=1}^{n_1} \hat{\mathfrak{A}}$. Continuing this process we either find a sequence of nonempty open sets $\mathfrak{D}_i \subset \hat{\mathfrak{A}} - \cap_{n=1}^{n_{i-1}} \hat{\mathfrak{A}}$ with $n_i \rightarrow \infty$ or $\hat{\mathfrak{A}}$ contains a locally compact open set $\mathfrak{D}$ such that any nontrivial open subset of $\mathfrak{D}$ contains representations of arbitrary high dimensions. In the first case let $J \subset \mathfrak{A}$ be the ideal corresponding to $\mathfrak{D} = \bigcup_{i=1}^{\infty} \mathfrak{D}_i$, then J has the form $J = \sum' \oplus J_i$ with $J_i = \mathfrak{D}_i$ [2, 1.9.12]. Thus by our result above J contains an ∞ -nilpotent element. In the second case determine an infinite family $\{\mathfrak{D}_i\}$ of nonempty disjoint open subsets of $\mathfrak{D}$. Let J_i be the ideal corresponding to $\mathfrak{D}_i$. Then the proof of Theorem 2 shows that J_i contains an i -nilpotent element x_i with $|x_i| \leq 2^{-i}$. Clearly $x = \sum \oplus x_i$ is ∞ -nilpotent.

An amusing consequence of Theorem 3 is the following result: Let $\mathfrak{A}$ be a separable C^* -algebra; then $\hat{\mathfrak{A}} = \cap_n \hat{\mathfrak{A}}$ iff for any singly generated subalgebra $\mathfrak{B}$ of $\mathfrak{A}$ we have $\hat{\mathfrak{B}} = \cap_n \hat{\mathfrak{B}}$.

REFERENCES

1. H. Behncke, *Nilpotent elements in group algebras*, Bull. Acad. Polon. Sci. **9** (1971), 197–198.
2. J. Dixmier, *Les C^* -algèbres et leurs représentations*, Cahiers Scientifiques, fasc. 29, Gauthier-Villars, Paris, 1964. MR **30** #1404.
3. C. E. Rickart, *General theory of Banach algebras*, University Series in Higher Math., Van Nostrand, Princeton, N.J., 1960. MR **22** #5903.
4. D. M. Topping, *Transcendental quasi-nilpotents in operator algebras*, J. Functional Analysis **2** (1968), 342–351. MR **38** #1535.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HEIDELBERG, HEIDELBERG, FEDERAL REPUBLIC OF GERMANY

Current address: Fakultät für Mathematik, Universität Bielefeld, 48 Bielefeld, Federal Republic of Germany