ON FINITELY GENERATED SUBGROUPS WHICH ARE OF FINITE INDEX IN GENERALIZED FREE PRODUCTS

A. KARRASS AND D. SOLITAR¹

ABSTRACT. Let $G=(A * B; U)$ be the free product of A and B with the subgroup U amalgamated. Various conditions are given which imply that every finitely generated subgroup H containing a (nontrivial) normal subgroup of G has finite index in G (in such a case we say G has the f.g.c.n. property). In particular, if A is a non-cyclic free group and U is cyclic, then G has the f.g.c.n. property. We use this last result to give a combinatorial proof that Fuchsian groups have the f.g.c.n. property; this was first proved by Greenberg using non-Euclidean geometry.

1. Introduction. We say that a group G has the *finitely generated normal* (f.g.n.) *property* if every nontrivial f.g. normal subgroup of G is of finite index (f.i.) in G .

G has the *finitely generated containing a normal* (f.g.c.n.) *property* if every f.g. subgroup which contains a nontrivial normal subgroup of G is of f.i. in G .

The f.g.n. property does not imply the f.g.c.n. property. For example, consider the group P of all permutations on the integers. The only normal subgroups of P besides the identity and P itself are the subgroup consisting of all finite permutations and the subgroup consisting of all finite even permutations (see [9]); hence the only f.g. normal subgroup of P is the identity subgroup, and so P has the f.g.n. property. On the other hand, the subgroup H generated by the transposition (12) and the infinite cycle $(\dots, -2, -1, 0, 1, 2, \dots)$ contains all transpositions and hence contains the normal subgroup consisting of all finite permutations; since H is countable but P is not, H is not of f.i. in P ; thus P does not have the f.g.c.n. property.

If $G=A * B$ is a proper free product (i.e., $A \neq 1 \neq B$), then G has the f.g.c.n. property (see [1]).

Received by the editors August 23, 1971.

AMS 1970 *subject classifications*. Primary 20E30, 20F05, 20F25.

Key words and phrases. Generalized free products, amalgamated products, small subgroups.

¹ This research was supported by grants from the National Research Council of Canada, Grant #A5602 and Grant #A5614.

© American Mathematical Society 1973

A generalization of this result, proved in [7], is the following: If $G = (A * B; U)$ where U is finite and $A \neq U \neq B$, and if H is a f.g. subgroup containing a normal subgroup of G not contained in U , then H is of f.i. in G ; in particular, if U contains no subgroup normal in both A and B , then G has the f.g.c.n. property.

However, as soon as we try to generalize to the case where U is infinite, even infinite cyclic, we encounter difficulties. For example, if

$$(1) \quad G = \langle a, b; a^2 = b^3 \rangle,$$

then the commutator subgroup of G is f.g. but is of infinite index. Note, however, in (1) the amalgamated subgroup is of finite index in both factors. Indeed, Theorem 1 shows that if $G = (A * B; U)$ where U is of infinite index in A , the subgroups of U are f.g., and A has the f.g.n. property, then G has the f.g.n. property.

To establish the more stringent f.g.c.n. property we require an additional hypothesis, namely, that U be a "small subgroup" of A (a precise definition of small is given in §3). As a special case, suppose $A = (C * D; V) \neq Z_2 * Z_2$ where V is finite, $C \neq V \neq D$, and V contains no nontrivial subgroup normal in both C and D ; then $G = (A * B; U)$ has the f.g.c.n. property provided the subgroups of U are all f.g.

We apply this result to prove that (hyperbolic) Fuchsian groups have the f.g.c.n. property. This was first proved in [3] using geometrical methods and later proved in [4] using topological techniques. Both of these papers actually establish the "f.g. containing a subnormal property". By an observation in [2], the method employed here can be used to establish this more general property.

Theorem 2 may be of independent interest. It implies a generalization of the well-known result that a proper free product cannot be a proper direct product: If G is a proper free product, and $G = HK$ where H, K are f.g. subgroups of G , then H or K is of f.i. in G .

For other conditions which yield the f.g.c.n. property for amalgamated products and also yield the f.g. intersection property see [2].

2. The f.g.n. property.

THEOREM 1. *Let $G = (A * B; U)$ where U is of infinite index in A , A has the f.g.n. property, and the subgroups of U are f.g. Then G has the f.g.n. property.*

PROOF. Let H be a f.g. normal subgroup of G . Since U is of infinite index in A , H is not contained in U . Therefore by Theorem 10 of [7], HU is of f.i. in G . Hence (using the notation of [7]) the collection $\{D_\alpha E_u\}$ of

u -double coset representatives for $G \bmod(H, U)$ is finite. Now for a fixed D_α , the associated collection $\{E_u\}$ is a double coset representative system for $A \bmod(A \cap D_\alpha^{-1} H D_\alpha, U)$; since U is not of finite index in A , $A \cap H = A \cap D_\alpha^{-1} H D_\alpha \neq 1$ and is a normal subgroup of A . Moreover, since every subgroup of U is f.g., it follows as in the proof of Theorem 8 of [7] that $A \cap H$ is f.g., and so $A \cap H$ is of f.i. in A . But HA has f.i. in G ; consequently, H is of f.i. in G . $\square$

To show that the f.g.n. hypothesis on A is necessary (even if $B \neq U$), let $A = \langle x, y; x^2 = y^3 \rangle$, $B = \langle b \rangle$, $U = \text{gp}(y^2) = \text{gp}(b^3)$. Then the commutator subgroup G' of $G = (A * B; U)$ is f.g. ($G'Z$ has f.i. in G , where Z is the center of G) but G' is of infinite index in G .

3. Small subgroups. Let U be a subgroup of a group A . We say that U is a *small subgroup* of A if whenever (H, U) has finite double coset index in A for a f.g. subgroup H , then H itself has f.i. in A .

If A is infinite and U is a small subgroup of A , then U must be of infinite index in A . For otherwise, $(1, U)$ has f.i. in A although 1 has infinite index in A .

Any finite subgroup is small. If U is small in A , then U is small in any finite extension of A .

It is also clear that *any subgroup of a small subgroup of A is small in A .*

Let $A = C * D$, $C \neq 1 \neq D$, be f.g.; then any nontrivial normal subgroup is large (i.e., not small) in A . For, suppose $C \neq Z_2$; it is easy to show that N contains an element n whose reduced form begins in a D -syllable and ends in a C -syllable. Let $H = \text{gp}(n^{-1} C n, D)$. Clearly $[G : (H, N)] = [G : HN] = 1$. On the other hand, if we choose $1 \neq d \in D$ and $1 \neq c \in C$ so that c is different from the last syllable of n , then no positive power of dc is in H ; hence $[G : H]$ is infinite and N is large. If $C = D = Z_2$, then N has finite index in A and N is therefore large.

THEOREM 2. *Let $A = (C * D; V)$ where V is finite and $C \neq V \neq D$. Then any f.g. subgroup of infinite index in A is small.*

PROOF. Let

$$(2) \quad a_1 \cdots a_r$$

be a reduced form of an element of a , i.e., a_i is in one of the factors C, D but a_i, a_{i+1} are not both in the same factor. A *segment* of (2) is a reduced form $a_j a_{j+1} \cdots a_k$, $1 \leq j, k \leq r$. If $g_1 \cdots g_r$ is another reduced form of the element defined by (2) in A , then $g_j \cdots g_k \in V a_j \cdots a_k V$. For, $g_i \cdots g_r a_r^{-1} \cdots a_i^{-1} \in V$ since $g_1 \cdots g_r a_r^{-1} \cdots a_1^{-1} = 1$; hence

$$g_j \cdots g_k = (g_j \cdots g_r a_r^{-1} \cdots a_j^{-1}) a_j \cdots a_k (a_{k+1} \cdots a_r g_r^{-1} \cdots g_{k+1}^{-1}),$$

which is in $V a_j \cdots a_k V$.

We call two elements a, a' of A *equivalent* (denoted $a \sim a'$) if $a' \in VaV$. By the preceding remark, if an element is equivalent to a segment of one reduced form of a , then it is equivalent to the corresponding segment of any other reduced form of a (and in fact to the corresponding segment of an element equivalent to a).

Suppose w is equivalent to a segment of the product xyz but w is not equivalent to a segment of x, y, z, xy or yz . Then $w \sim x_1 y z_1$ where x_1 is a terminal segment of a given reduced form of x and z_1 is an initial segment of a given reduced form of z . We prove this by induction on the sum $|x| + |z|$ of the syllable lengths of the given reduced forms of x and z . Clearly, neither x nor z is in V . Suppose first that $|x| = |z| = 1$; then it is easy to show by examining reduced forms of y and xyz that $w \sim xyz$. Suppose then $|x| + |z| > 2$, and let $x = ax', z = z'b$ where a and b are the first and last syllables respectively of the given reduced forms of x and z . Then w is equivalent to a segment of $a(x'yz')b$. If w is not equivalent to a segment of $x'yz', (ax')yz',$ or $x'y(z'b)$, then by the preceding case, $w \sim xyz$; on the other hand, if w is equivalent to a segment of $x'yz', (ax')yz',$ or $x'y(z'b)$, then by inductive hypotheses, $w \sim x_1 y z_1$, where x_1 and z_1 are as asserted.

Let H be any f.g. subgroup of infinite index in A . We show that there exists an element w of A such that w is not equivalent to any segment of any element of H . Our proof is based on the proof for the corresponding result in free groups given by G. Higman in [5]. Suppose every element w of A is equivalent to some segment of some element of H . Let $h_1, \dots, h_s$ be given reduced forms for a finite set of generators of H . Since V is finite, there are only finitely many elements equivalent to a segment of $h_i^{\varepsilon_i}$, $\varepsilon_i = \pm 1$. Suppose now w is equivalent to a segment of $h_{i_1}^{v_1} \cdots h_{i_n}^{v_n}$, $v_i = \pm 1$, where n is minimal. If $n > 1$, then w is equivalent to a segment of xyz where $x = h_{i_1}^{v_1}, y = h_{i_2}^{v_2} \cdots h_{i_{n-1}}^{v_{n-1}}, z = h_{i_n}^{v_n}$; hence by the above, w is equivalent to $x_1 y z_1$ where x_1 and z_1 are segments of $h_i^{\varepsilon_i}$. Since $y \in H$, it follows that A is the union of finitely many sets of the form $a_i H b_i = (a_i H a_i^{-1}) a_i b_i$, i.e., A is the union of finitely many cosets of finitely many conjugates of H . But Lemma 4.1 in [8] then implies that H must be of f.i. in A , contrary to hypothesis.

To prove Theorem 2, let U be a f.g. subgroup of infinite index in A . We show (H, U) has infinite double coset index in A . For, suppose A is the union $H g_i U, i = 1, \dots, n$. Let w_1, w_2 be reduced forms not equivalent to any segments of any elements of H, U respectively. Let p be a reduced form such that $w_1 p w_2$ is reduced and $|p| > \max |g_i|$. For some $j, g_j = h^{-1} w_1 p w_2 u^{-1}$, with $h \in H, u \in U$. Since w_1 is not equivalent to an initial segment of h , the last syllable of $h^{-1} w_1$ comes from the same factor as the last syllable of w_1 ; similarly, the first syllable of $w_2 u^{-1}$ comes from the same

factor as the first syllable of w_2 . Hence $|g_i| \geq |p|$, contrary to the choice of p .

Thus U is a small subgroup of A if U is f.g. and of infinite index in A . $\square$

COROLLARY. *Let $A = C * D$, $C \neq 1 \neq D$. If H is a f.g. subgroup of A and contains a nontrivial normal subgroup N of A then H is of f.i. in A .*

PROOF. We first show that C and D are f.g. For suppose say C is infinitely generated. Since H is f.g., there exist f.g. subgroups C' , D' of C , D respectively, such that $H < C' * D'$. Hence, $N < C' * D'$. If $c \in C - C'$, then $cnc^{-1} \notin C' * D'$ if $n \neq 1$. Thus C , D are f.g.

But if H has infinite index in A , H is small and therefore N is small; but N is large by the remark immediately preceding Theorem 2. $\square$

This corollary was proved in [1] by a different method.

COROLLARY. *Suppose $A = (C * D; V)$, V finite, $C \neq V \neq D$. If $A = HK$ where H , K are f.g. subgroups of A , then H or K is of f.i. in A . In particular, if A is the direct product $H \times K$, then C , D , V can be written in the form $C = C' \times Y$, $D = D' \times Y$, $V = V' \times Y$ where Y is either H or K .*

PROOF. Since the double coset index of (H, K) is 1, the first assertion holds. If $A = H \times K$, then H or K must be finite, say K . Since K is finite and normal, K must be in C or D , and hence within V . Take $C' = C \cap H$, $D' = D \cap H$, $V' = V \cap H$, $Y = K$. $\square$

4. The f.g.c.n. property.

THEOREM 3. *Let $G = (A * B; U)$ where U is a small subgroup of A , $A \neq U \neq B$ and the subgroups of U are all f.g. If H is a f.g. subgroup of G and contains a normal subgroup N , $N \triangleleft U$, then H is of f.i. in G . In particular, if U contains no nontrivial subgroup normal in both A and B , then G has the f.g.c.n. property.*

PROOF. By Theorem 10 of [7], (H, U) has finite double coset index in G . Hence as in the proof of Theorem 1, for a fixed α -double coset representative D_α , the double coset index of $(A \cap D_\alpha^{-1} H D_\alpha, U)$ is finite in A and $A \cap D_\alpha^{-1} H D_\alpha$ is f.g. Since U is small in A , $A \cap D_\alpha^{-1} H D_\alpha$ is of f.i. in A . Moreover, since (H, A) is of f.i. in G , it follows that H is of f.i. in G . $\square$

As R. Burns [2] has observed, Theorem 10 of [7] holds if N is assumed merely to be subnormal in G ; consequently, Theorem 3 holds with "normal" replaced by "subnormal".

COROLLARY. *Suppose $A = (C * D; V) \neq Z_2 * Z_2$, V finite, $C \neq V \neq D$, and V contains no nontrivial subgroup normal in both C and D . Then*

$G = (A * B; U)$ has the f.g.c.n. property provided the subgroups of U are f.g.

PROOF. Clearly V is not of index two in C and D . Hence A contains a free subgroup of rank two. Since the subgroups of U are all f.g., U cannot be of finite index in A . Therefore U is small in A and cannot contain a nontrivial normal subgroup of A . $\square$

5. The f.g.c.n. property for Fuchsian groups. A (hyperbolic) Fuchsian group (or equivalently, a discrete group of orientation-preserving motions of the hyperbolic plane) G is either a free product of cyclic groups or has a presentation

$$(3) \quad G = \langle a_1, b_1, \dots, a_n, b_n, c_1, \dots, c_t; \\ c_1^{\gamma_1}, \dots, c_t^{\gamma_t}, c_1^{-1} \cdots c_t^{-1} [a_1, b_1] \cdots [a_n, b_n] \rangle$$

with $n, t \geq 0$, $\gamma_i > 1$, and $\mu(G) > 0$ where

$$\mu(G) = 2n - 2 + (1 - \gamma_1^{-1}) + (1 - \gamma_2^{-1}) + \cdots + (1 - \gamma_t^{-1}).$$

Greenberg [3] has shown geometrically that these discrete hyperbolic groups have the f.g.c.n. property. We sketch here a combinatorial proof of this result.

If G is a free product of cyclic groups, then G has the f.g.c.n. property. Suppose G is as in (3). Then G is known to be a finite extension of a group of the form (3) with $t=0$ and $n \geq 2$; also G cannot contain a finite nontrivial normal subgroup (for combinatorial arguments see the last part of [6]). Now such a subgroup has the f.g.c.n. property, since it can be written in the form $(A * B; U)$ with $A = \langle a_1, b_1 \rangle$, $B = \langle a_2, b_2, \dots, a_n, b_n \rangle$, $U = \text{gp}([b_1, a_1]) = \text{gp}([a_2, b_2] \cdots [a_n, b_n])$, which satisfies the conditions of the corollary to Theorem 3. Consequently G itself has the f.g.c.n. property.

Similarly, one can show that G has the "f.g. containing a subnormal" property.

REFERENCES

1. B. Baumslag, *Intersections of finitely generated subgroups in free products*, J. London Math. Soc. **41** (1966), 673-679. MR **33** #7396.
2. R. Burns, *On the finitely generated subgroups of an amalgamated product of two groups*, Trans. Amer. Math. Soc. **169** (1972), 293-306.
3. L. Greenberg, *Discrete groups of motions*, Canad. J. Math. **12** (1960), 414-426. MR **22** #5932.
4. H. B. Griffiths, *A covering-space approach to theorems of Greenberg in Fuchsian, Kleinian, and other groups*, Comm. Pure Appl. Math. **20** (1967), 365-399; Correction, ibid. **21** (1968), 521-522. MR **35** #2282; MR **38** #1671.
5. G. Higman, *Some problems and results in the theory of groups*. II, Notes of a Mini-Conference, Oxford, 12th and 13th August 1966, pp. 15-16.

6. A. H. M. Hoare, A. Karrass and D. Solitar, *Subgroups of infinite index in Fuchsian groups*, Math. Z. **125** (1972), 59–69.
7. A. Karrass and D. Solitar, *The subgroups of a free product of two groups with an amalgamated subgroup*, Trans. Amer. Math. Soc. **150** (1970), 227–255. MR **41** #5499.
8. B. H. Neumann, *Groups covered by permutable subsets*, J. London Math. Soc. **29** (1954), 236–248. MR **15**, 931.
9. J. Schreier and S. Ulam, *Über die Permutationsgruppe der natürlichen Zahlenfolge*, Studia Math. **4** (1933), 134–141.

DEPARTMENT OF MATHEMATICS, YORK UNIVERSITY, DOWNSVIEW, ONTARIO, CANADA