ON THE COEFFICIENTS OF AN ASYMPTOTIC EXPANSION OF SPHERICAL FUNCTIONS ON SYMMETRIC SPACES

JOHN J. H. MILLER AND D. J. SIMMS

ABSTRACT. The asymptotic expansion of a spherical function on a symmetric space of noncompact type, obtained by Harish-Chandra, is a finite linear combination of expansions of the form $\Phi_\theta = \sum_\mu \Gamma_\mu(\theta) e^{\theta - \mu}$. In this paper it is proved that $\lim_{t \rightarrow \infty} \Gamma_\mu(te - \bar{\rho})$ is finite and rational for any e , where $\bar{\rho}$ is the restriction of half the sum of the positive roots.

1. Introduction. Let G be a connected real semisimple Lie group with finite centre and K a maximal compact subgroup. Let $G = KAN$ be an Iwasawa decomposition of G , relative to a system $\{\alpha\}$ of simple roots of G , and among their restrictions $\{\bar{\alpha}\}$ to the Lie algebra $\mathfrak{A}$ of A choose a system $\bar{\alpha}_1, \dots, \bar{\alpha}_l$ of simple restricted roots of the symmetric space G/K . Denote by L the lattice of all integral combinations $\mu = \sum_{i=1}^l m_i \bar{\alpha}_i$ and let L^+ be the set of μ with each m_i nonnegative. Write $\nu \leq \mu$ if $\mu - \nu \in L^+$. Regard each μ in the complex dual Λ^C of $\mathfrak{A}$ as a linear form on Λ^C via the Killing form $\langle \cdot, \cdot \rangle$. Let Δ_+ be the set of positive roots of G , $\bar{\rho}$ the restriction to $\mathfrak{A}$ of half the sum of the positive roots; and $Q = \prod_{\beta \in P_+} (e^\beta - e^{-\beta})$ where $P_+ = \{\beta \in \Delta_+ | \bar{\beta} \neq 0\}$.

For each $\mu \in L$ let Γ_μ be the rational function on Λ^C defined by the recursion relation

$$\begin{aligned} \Gamma_\mu &= 0 \quad \text{if } \mu \in L - L^+, \\ \Gamma_0 &= 1, \\ c_\mu \Gamma_\mu &= \sum_{\beta \in \Delta_+} \sum_{k=1}^{\infty} \gamma(\beta)_{\mu - 2k\bar{\beta}} \Gamma_{\mu - 2k\bar{\beta}} \quad \text{if } \mu \in L^+ - \{0\}, \end{aligned} \tag{1.1}$$

where the coefficients are the functions on Λ^C given by

$$c_\mu = \langle \mu, \mu - 2\bar{\rho} \rangle - 2\mu, \quad \gamma(\bar{\beta})_\mu = 2\langle \mu, \bar{\beta} \rangle - 2\bar{\beta}. \tag{1.2}$$

Received by the editors March 9, 1972.

AMS (MOS) subject classifications (1970). Primary 43A90; Secondary 17B20, 22E30, 22E60.

Key words and phrases. Symmetric space of noncompact type, spherical function, coefficient of asymptotic expansion, radial limit.

According to Harish-Chandra [3], each spherical function on G/K , when transferred to $\mathfrak{A}$ by the exponential, can be written as a finite linear combination of asymptotic expansions of the form

$$\Phi_\theta = \sum_{\mu \in L} \Gamma_\mu(\theta) e^{\theta - \mu}.$$

In [7] we showed that for $e \in \Lambda$ the radial limit $\lim_{t \rightarrow \infty} \Gamma_\mu(ite - \bar{\rho})$ is finite and independent of e except possibly for those exceptional directions of e for which $\langle e, \nu \rangle = 0$ for some $0 \leq \nu \leq \mu$. In this paper we show, by a different method, that the radial limit $\lim_{t \rightarrow \infty} \Gamma_\mu(te - \bar{\rho})$ is finite for all $e \in \Lambda^C$ without exception. We also calculate $\Gamma_\mu(\theta)$ explicitly when μ is a multiple of a simple restricted root which is not itself a multiple of any other positive restricted root. This shows in particular that the radial limit in an exceptional direction may differ from the limit in a nonexceptional direction.

2. The radial limits.

THEOREM. Let $e, a \in \Lambda^C$, $\mu \in L^+$ and suppose that $\langle e, \nu \rangle \neq 0$ for all $0 \leq \nu \leq \mu$. Then $\lim_{t \rightarrow \infty; t \in C} \Gamma_\mu(te + a)$ exists and is equal to the positive rational number A_μ defined by

$$e^{\bar{\rho}} Q^{-1/2} = \prod_{\beta \in P^+} (1 - e^{-2\beta})^{-1/2} = \sum_{\mu \in L} A_\mu e^{-\mu}.$$

For all $e \in \Lambda^C$, without exception, $\lim_{t \rightarrow \infty; t \in C} \Gamma_\mu(te - \bar{\rho})$ exists and is a rational number.

PROOF. Let the rational functions C_μ and the rational numbers D_μ be defined by

$$Q^{1/2} \Phi_\theta = e^{\theta + \bar{\rho}} \sum_{\mu \in L} C_\mu(\theta) e^{-\mu}, \quad Q^{-1/2} \nabla^2 (Q^{1/2}) = \sum_{\mu \in L} D_\mu e^{-\mu},$$

where ∇^2 is the Laplacian on $\mathfrak{A}$. We note that A_μ , C_μ and D_μ are all zero for $\mu \in L - L^+$, $A_0 = 1$, $C_0 \equiv 1$, $D_0 = \langle \bar{\rho}, \bar{\rho} \rangle$, and $A_\mu \geq 0$ for all $\mu \in L$. By a result of Gangolli [2] in a form given by Helgason [5, Chapter II, §2 p. 38] we have:

$$\{\langle \mu, \mu \rangle - 2\langle \mu, \theta + \bar{\rho} \rangle\} C_\mu(\theta) = \sum_{\nu \in L; \nu \neq \mu} D_{\mu - \nu} C_\nu(\theta).$$

Now fix $e, a \in \Lambda^C$ and put $\theta = te + a$, $t \in C$, to get

$$(2.1) \quad \{\langle \mu, \mu - 2a - 2\bar{\rho} \rangle - 2t\langle \mu, e \rangle\} C_\mu(te + a) = \sum_{\nu \in L; \nu \neq \mu} D_{\mu - \nu} C_\nu(te + a).$$

In particular, for $a = -\bar{\rho}$, we have

$$(2.2) \quad \{\langle \mu, \mu \rangle - 2t\langle \mu, e \rangle\} C_\mu(te - \bar{\rho}) = \sum_{\nu \in L; \nu \neq \mu} D_{\mu - \nu} C_\nu(te - \bar{\rho}).$$

We now distinguish two cases, depending on a and on the direction of e .

Case (i). For all $a, e \in \Lambda^C$ such that $\langle e, v \rangle \neq 0$ for all $0 \leq v \leq \mu$ ('non-exceptional' directions of e): In this case we let $t \rightarrow \infty$ in (2.1) and use induction on μ to conclude that

$$(2.3) \quad \lim_{t \rightarrow \infty} C_\mu(te + a) = 0$$

for all $\mu \in L^+, \mu \neq 0$.

Case (ii). For $a = -\bar{\rho}$ and for all $e \in \Lambda^C$: In this case we let $t \rightarrow \infty$ in (2.2) and use induction on μ to conclude that

$$(2.4) \quad \lim_{t \rightarrow \infty} C_\mu(te - \bar{\rho})$$

is finite and rational for all $\mu \in L$. Now

$$\sum_{\mu \in L} \Gamma_\mu(\theta) e^{\theta - \mu} = \Phi_\theta = \left(\sum_{\mu \in L} A_\mu e^{-\mu} \right) \cdot \left(\sum_{\mu \in L} C_\mu(\theta) e^{\theta - \mu} \right);$$

therefore

$$\Gamma_\mu(\theta) = \sum_{v \in L} A_{\mu-v} C_v(\theta).$$

For Case (i), $\langle e, v \rangle \neq 0$ for all $0 \leq v \leq \mu$, so that

$$\lim_{t \rightarrow \infty} \Gamma_\mu(te + a) = \sum_{v \in L} A_{\mu-v} \lim_{t \rightarrow \infty} C_v(te + a) = A_\mu$$

by (2.3). Also, Case (ii), we have for all $e \in \Lambda^C$

$$\lim_{t \rightarrow \infty} \Gamma_\mu(te - \bar{\rho}) = \sum_{v \in L} A_{\mu-v} \lim_{t \rightarrow \infty} C_v(te - \bar{\rho})$$

which is finite and rational by (2.4). This completes the proof.

REMARK. The Theorem shows that the rational numbers A_μ are the same as those introduced in [7] by the recursion relation

$$(2.5) \quad \mu A_\mu = \sum_{\beta \in I^+} \sum_{k=1}^{\infty} \tilde{\beta} A_{\mu-2k\beta}.$$

This recursion relation can easily be obtained by applying the gradient operator ∇ to $e^{\bar{\rho}} Q^{-1/2}$ to get

$$\begin{aligned} \sum_{\mu \in L} \mu A_\mu e^{-\mu} &= \sum_{\beta \in P_+} \tilde{\beta} e^{-2\tilde{\beta}} (1 - e^{-2\tilde{\beta}})^{-1} \prod_{\beta \in P_+} (1 - e^{-2\tilde{\beta}})^{-1/2} \\ &= \sum_{\beta \in I^+} \tilde{\beta} \sum_{k=1}^{\infty} e^{-2k\tilde{\beta}} \sum_{\mu \in L} A_\mu e^{-\mu}. \end{aligned}$$

Equating the coefficients of $e^{-\mu}$ gives (2.5).

3. **Some calculations and estimates for Γ_μ and A_μ .** The following proposition deals with the special case $q=0$ of [4, §5, p. 304]. It gives an improvement of Helgason's estimate for this case.

PROPOSITION. *Let α be a root of G such that $\bar{\alpha}$ is one of the restricted simple roots $\bar{\alpha}_1, \dots, \bar{\alpha}_l$. Let p be the number of positive roots of G whose restriction to $\mathfrak{A}$ is equal to $\bar{\alpha}$, and suppose that $\bar{\alpha}$ is not a scalar multiple of any other restricted root. Let $\eta = \langle \bar{\alpha}, \bar{\alpha} \rangle^{-1} \bar{\alpha}$. Then for all $m \geq 1$, $\theta \in \Lambda^C$ we have $\eta(\bar{\rho}) = \frac{1}{2}p$ and*

$$(3.1) \quad \Gamma_{2m\bar{\alpha}}(\theta) = \prod_{r=1}^m \left(1 + \frac{p-2}{2r} \right) \left(1 + \frac{p-2}{2r-p-2\eta(\theta)} \right).$$

PROOF. Put μ equal to $2m\bar{\alpha}$ and $2(m-1)\bar{\alpha}$ in turn in (1.1) to get

$$\begin{aligned} c_{2m\bar{\alpha}} \Gamma_{2m\bar{\alpha}} &= \sum_{k=1}^{\infty} p\gamma(\alpha)_{2(m-k)\bar{\alpha}} \Gamma_{2(m-k)\bar{\alpha}}, \\ c_{2(m-1)\bar{\alpha}} \Gamma_{2(m-1)\bar{\alpha}} &= \sum_{k=1}^{\infty} p\gamma(\alpha)_{2(m-k-1)\bar{\alpha}} \Gamma_{2(m-k-1)\bar{\alpha}}. \end{aligned}$$

Subtracting we get

$$(3.2) \quad c_{2m\bar{\alpha}} \Gamma_{2m\bar{\alpha}} - c_{2(m-1)\bar{\alpha}} \Gamma_{2(m-1)\bar{\alpha}} = p\gamma(\alpha)_{2(m-1)\bar{\alpha}} \Gamma_{2(m-1)\bar{\alpha}}.$$

It is an easy and well-known consequence of [1, VI, §1.6, Corollary 1] that $\langle \bar{\alpha}, \bar{\rho} \rangle = \frac{1}{2}p \langle \bar{\alpha}, \bar{\alpha} \rangle$. Thus $\eta(\bar{\rho}) = \frac{1}{2}p$. From (1.2) we have

$$\begin{aligned} c_{2m\bar{\alpha}}(\theta) &= \langle 2m\bar{\alpha}, 2m\bar{\alpha} - 2\bar{\rho} - 2\theta \rangle = 4m[m - \eta(\bar{\rho} + \theta)] \langle \bar{\alpha}, \bar{\alpha} \rangle, \\ \gamma_{2m\bar{\alpha}}(\theta) &= \langle 4m\bar{\alpha}, \bar{\alpha} \rangle - \langle 2\bar{\alpha}, \theta \rangle = [4m - 2\eta(\theta)] \langle \bar{\alpha}, \bar{\alpha} \rangle. \end{aligned}$$

Substitution in (3.2) gives, with the use of $\eta(\bar{\rho}) = \frac{1}{2}p$,

$$\begin{aligned} \Gamma_{2m\bar{\alpha}}(\theta) &= \frac{[4(m-1) + 2p][m-1-\eta(\theta)]}{4m[m-\frac{1}{2}p-\eta(\theta)]} \Gamma_{2(m-1)\bar{\alpha}}(\theta) \\ &= (1 + (p-2)/2m)(1 + (p-2)/(2m-p-2\eta(\theta))) \Gamma_{2(m-1)\bar{\alpha}}(\theta). \end{aligned}$$

This gives (3.1).

COROLLARIES. *Let $m \geq 1$.*

(i) *Since $\eta(\bar{\rho}) = \frac{1}{2}p$ we have, for all $\lambda \in \Lambda^C$,*

$$\Gamma_{2m\bar{\alpha}}(\lambda - \bar{\rho}) = \prod_{r=1}^m \left(1 + \frac{p-2}{2r} \right) \left(1 + \frac{p-2}{2r-2\eta(\lambda)} \right).$$

(ii) *In a 'nonexceptional' direction of $e \in \Lambda^C$ given by $\langle \bar{\alpha}, e \rangle \neq 0$ we have*

$\eta(e) \neq 0$ and therefore

$$A_{2m\bar{\alpha}} = \lim_{t \rightarrow \infty} {}'\Gamma_{2m\bar{\alpha}}(te - \bar{\rho}) \\ = \lim_{t \rightarrow \infty} \prod_{r=1}^m \left(1 + \frac{p-2}{2r}\right) \left(1 + \frac{p-2}{2r-2t\eta(e)}\right) = \prod_{r=1}^m \left(1 + \frac{p-2}{2r}\right).$$

(iii) In an 'exceptional direction' given by $\langle \bar{\alpha}, \lambda \rangle = 0$ we have $\eta(\lambda) = 0$ and therefore by (i) and (ii)

$${}'\Gamma_{2m\bar{\alpha}}(\lambda - \bar{\rho}) = \prod_{r=1}^m \left(1 + \frac{p-2}{2r}\right)^2 = A_{2m\bar{\alpha}}^2.$$

(iv) If $p=1$ then for $\lambda \in \Lambda$ we have

$$|{}'\Gamma_{2m\bar{\alpha}}(i\lambda - \bar{\rho})| = \left| \prod_{r=1}^m \left(1 - \frac{1}{2r}\right) \left(1 - \frac{1}{2r-2i\eta(\lambda)}\right) \right| < \prod_{r=1}^m \left(1 - \frac{1}{2r}\right)$$

since $i\eta(\lambda)$ is pure imaginary.

(v) If $p=2$ then ${}'\Gamma_{2m\bar{\alpha}}(\theta) = 1$ for all $\theta \in \Lambda^c$.

(vi) If $p > 2$ and $\lambda \in \Lambda$, then the absolute value of $1 + (p-2)/(2r-2i\eta(\lambda))$ is greater than 1 and less than or equal to $1 + (p-2)/2r$. Therefore, using (ii) we have $A_{2m\bar{\alpha}} < |{}'\Gamma_{2m\bar{\alpha}}(i\lambda - \bar{\rho})| \leq A_{2m\bar{\alpha}}^2$.

REMARK. The conditions of the Proposition are satisfied provided that α is represented by a node of the form $\circ$, and not of the form $\odot$ in Tables 4 and 8 on pp. 119 and 146 of [6, Chapter VII].

ACKNOWLEDGEMENT. We should like to thank Sigurdur Helgason for stimulating discussions and for making available to us the manuscript of [5] before publication.

REFERENCES

1. N. Bourbaki, *Éléments de mathématiques*. XXVI. *Groupes et algèbres de Lie*, Actualités Sci. Indust., no. 1285, Hermann, Paris, 1960. MR 24 #A2641.
2. R. Gangolli, *On the Plancherel formula and the Paley-Wiener theorem for spherical functions on semisimple Lie groups*, Ann. of Math. (2) 93 (1971), 150-165.
3. Harish-Chandra, *Spherical functions on a semisimple Lie group*. I, Amer. J. Math. 80 (1958), 241-310. MR 20 #925.
4. S. Helgason, *An analogue of the Paley-Wiener theorem for the Fourier transform on certain symmetric spaces*, Math. Ann. 165 (1966), 297-308. MR 36 #6545.
5. ———, *A duality for symmetric spaces with applications to group representations*, Advances in Math. 5 (1970), 1-154. MR 41 #8587.
6. O. Loos, *Symmetric spaces*. II: *Compact spaces and classification*, Benjamin, New York, 1969. MR 39 #365b.
7. John J. H. Miller and D. J. Simms, *Radial limits of the rational functions ${}'\Gamma_\mu$ of Harish-Chandra*, Proc. Roy. Irish Acad. Sect. A 68 (1969), 41-47. MR 40 #7391.

SCHOOL OF MATHEMATICS, TRINITY COLLEGE, DUBLIN, IRELAND

Current address (D. J. Simms): Mathematisches Institut der Universität Bonn, 53 Bonn, Wegelerstrasse 10, Germany