A NOTE ON TORSION-FREE ABELIAN GROUPS OF FINITE RANK

W. WICKLESS AND C. VINSONHALER

ABSTRACT. Let G be a torsion-free abelian group of rank n and $X = \{x_1, \dots, x_n\}$ a maximal set of rationally independent elements in G . It is well known that any $g \in G$ can be uniquely written $g = \alpha_1 x_1 + \dots + \alpha_n x_n$, for some $\alpha_1, \dots, \alpha_n \in \mathbb{Q}$, the rational numbers. This enables us to define, for any such (G, X) , a collection of subgroups of $\mathbb{Q}$ and "natural" isomorphisms, denoted by $S(G, X)$. It is known that if G is of rank two, then G may be recovered from $S(G, X)$ in a natural way. The following result is obtained for groups of rank greater than two:

THEOREM. Let G, G' be torsion free abelian groups of finite rank with $S(G, X) = S(G', X')$ for suitable X, X' . Let F, F' be the free subgroups of G, G' generated by X, X' . Then $G/F \cong G'/F'$.

An additional condition is given for pairs $(G, X), (G', X')$ such that $S(G, X) = S(G', X')$ implies $G \cong G'$.

1. Schemes of groups. Let G be a group¹ of rank n and $X = \{x_1, \dots, x_n\}$ a maximal set of rationally independent elements in G . It is well known that any $g \in G$ can be uniquely written $g = \alpha_1 x_1 + \dots + \alpha_n x_n$, for some $\alpha_1, \dots, \alpha_n \in \mathbb{Q}$, the rational numbers. This enables us to define, for any such (G, X) , a collection of subgroups of $\mathbb{Q}$ as follows:

DEFINITION 1. Let $1 \leq i \leq n$ and $i \notin J \subset \{1, 2, \dots, n\}$. Then

$$\begin{aligned} A_J^i &= A_J^i(G, X) \\ &= \{\alpha \in \mathbb{Q} \mid \alpha_1 x_1 + \dots + \alpha_{i-1} x_{i-1} + \alpha x_i + \alpha_{i+1} x_{i+1} + \dots + \alpha_n x_n \in G \\ &\quad \text{for some set of } \alpha_j \in \mathbb{Q} \text{ with } \alpha_j = 0 \quad \forall j \in J\}. \end{aligned}$$

The A_J^i are clearly subgroups of the additive rationals and contain the integers, $\mathbb{Z}$. Furthermore, $A_J^i \supseteq A_{J'}^i$, if $J \subseteq J'$. For convenience, we write $A^i = A_\emptyset^i$. The A_J^i can be used to define a collection of "natural" isomorphisms.

LEMMA 1. Given $1 \leq i \neq j \leq n$, with $i, j \notin J \subset \{1, 2, \dots, n\}$, there exists an isomorphism

$$A_J^i / A_{\{j\} \cup J}^i \cong A_J^j / A_{\{i\} \cup J}^j.$$

Presented to the Society, September 1, 1972; received by the editors March 20, 1972.

AMS (MOS) subject classifications (1970). Primary 20K15.

¹ We write "group" for torsion-free abelian group throughout.

PROOF. Let $\alpha_i \in A_J^i$ and write $\sum_{k=1}^n \alpha_k x_k \in G$ for some choice of $\alpha_k \in Q$, $k \neq i$, with $\alpha_k = 0$ for $k \in J$. Then $(\alpha_i + A_{\{j\} \cup J}^i)^\theta = \alpha_j + A_{\{i\} \cup J}^j$ is the required isomorphism.

We call the collection of groups A_J^i and isomorphisms θ the scheme of (G, X) , and denote it by $S(G, X)$.

LEMMA 2. *Given $1 \leq i, j, k \leq n$ with $i, j, k \notin J \subset \{1, \dots, n\}$ and i, j, k distinct, then*

$$\begin{array}{ccc} A_J^i / A_{\{j\} \cup J}^i & \xrightarrow{\theta} & A_J^j / A_{\{i\} \cup J}^j \\ \uparrow NAT & & \uparrow NAT \\ A_{\{k\} \cup J}^i / A_{\{k, j\} \cup J}^i & \xrightarrow{\theta} & A_{\{k\} \cup J}^j / A_{\{k, i\} \cup J}^j \end{array}$$

is a commutative diagram where the θ 's are the isomorphisms of Lemma 1 and NAT is the composition of the obvious inclusion and factor maps.

PROOF. The proof follows immediately from the definition of the maps θ .

If G has rank 2, Beaumont and Wisner [2] showed that G may be recovered from the groups A_J^i and isomorphism θ in the following way: Let x_1, x_2 be independent elements in G and $\{A^1, A_2^1, A^2, A_1^2, \theta\}$ the scheme of $(G, \{x_1, x_2\})$. Then $G' = \{(\alpha, \beta) \mid \alpha \in A_1, (\alpha + A_2^1)^\theta = \beta + A_1^2\}$ is isomorphic to G . Moreover, $S(G', \{(1, 0), (0, 1)\}) = S(G, \{x_1, x_2\})$.

The question then arises: To what extent does $S(G, X)$ determine G in general? The results which follow give a partial answer.

Since any group G of rank n with maximal independent set $X = \{x_1, \dots, x_n\}$ can be identified with a subgroup of Q^n under the map $\alpha_1 x_1 + \dots + \alpha_n x_n \rightarrow (\alpha_1, \dots, \alpha_n)$, it suffices to consider pairs (G, X) with $Z^n \subseteq G \subseteq Q^n$ and $X = \{(1, 0, 0, \dots, 0), (0, 1, 0, \dots, 0), \dots, (0, 0, \dots, 0, 1)\}$. Let (G, X) and (G', X) be two such pairs.

LEMMA 3. *If $S(G, X) = S(G', X)$ then given any element $(\alpha_1, \alpha_2, \dots, \alpha_n) \in G$ and fixed integers $i \neq j$, with $1 \leq i, j \leq n$, there exists an element $(\alpha'_1, \alpha'_2, \dots, \alpha'_n) \in G'$ with $\alpha_i = \alpha'_i, \alpha_j = \alpha'_j$. Moreover, it is possible to choose $\alpha'_k = 0$ whenever $\alpha_k = 0$.*

PROOF. By symmetry, we may take $i = 1, j = 2$. Let $K = \{k \mid \alpha_k = 0\}$. Since $\alpha_1 \in A_K^1$ and $A_K^1(G, X) = A_K^1(G', X)$, there exists $(\alpha_1, \beta_2, \beta_3, \dots, \beta_n) \in G'$ with $\beta_k = 0$ for $k \in K$. Thus, $\beta_2 + A_{\{1\} \cup K}^2 = [\alpha_1 + A_{\{2\} \cup K}^1]^\theta = \alpha_2 + A_{\{1\} \cup K}^2$. Hence, $\alpha_2 - \beta_2 \in A_{\{1\} \cup K}^2$, and there exists $(0, \alpha_2 - \beta_2, \gamma_3, \gamma_4, \dots, \gamma_n) \in G'$ with $\gamma_k = 0$ for $k \in K$. Then $(\alpha_1, \alpha_2, \alpha'_3, \dots, \alpha'_n) = (0, \alpha_2 - \beta_2, \gamma_3, \dots, \gamma_n) + (\alpha_1, \beta_2, \dots, \beta_n) \in G'$ with $\alpha'_k = 0$ for $k \in K$.

LEMMA 4. Let G, G', X be as in Lemma 3, and p a prime integer. Let $g=(\alpha_1, \dots, \alpha_n) \in G$ and $g'_1, \dots, g'_t \in G'$ be such that $\bar{g}=g+Z^n$, $\bar{g}'_1=g'_1+Z^n, \dots, \bar{g}'_t=g'_t+Z^n$ are independent elements in the p -primary component of Q^n/Z^n . Then there exists $g'=(\alpha'_1, \dots, \alpha'_n) \in G'$, with $\text{order}(\bar{g}') \geq \text{order}(\bar{g})=p^k$, such that $\bar{g}'_1, \dots, \bar{g}'_t, \bar{g}'$ are independent. If $\alpha_r=a_r/p^k$, $\alpha_s=a_s/p^{k'}$, for integers a_r, a_s with $(a_r, p)=(a_s, p)=1$, $k' \leq k$, then g' can be chosen so that $c\alpha_r=\alpha'_r$, $c\alpha_s=\alpha'_s$, where c is an integer with $(c, p)=1$. Furthermore, we can choose $\alpha'_i=0$ whenever $\alpha_i=0$.

PROOF. We proceed by induction on the number, m , of nonzero coordinates of g . If $m=1, 2$, then $g \in G'$ by Lemma 3 and the result follows. Suppose now that $g=(\alpha_1, \dots, \alpha_m, 0, 0, \dots, 0)$. We assume, without loss of generality, that all integral components of g are zero, and, by renumbering if necessary, that the nonzero coordinates appear first. Write $\alpha_i=a_i/p^{k_i}$, $1 \leq k_i \leq k$, where $(a_i, p)=1$. Again by renumbering, assume $r=1, s=2$. Thus, $k_1=k$. By Lemma 3, there exists

$$h_1 = (a_1/p^{k_1}, a_2/p^{k_2}, \beta_3, \dots, \beta_m, 0, \dots, 0) \in G'.$$

Choose an integer c with $(c, p)=1$ such that

$$h_2 \equiv ch_1 = (ca_1/p^{k_1}, ca_2/p^{k_2}, b_3/p^{l_3}, \dots, b_m/p^{l_m}, 0, \dots, 0).$$

Note that $\{\bar{g}'_1, \dots, \bar{g}'_t, c\bar{g}\}$ is still independent. (The integer c times a power of p is divisible by the denominators of $\beta_3 \dots \beta_m$.)

Case I. Suppose $l_i=k_i$ and $p|ca_i-b_i$ for $3 \leq i \leq m$. Then $\text{order}(h_2)=p^k$. We show $\{\bar{g}'_1, \dots, \bar{g}'_t, \bar{h}_2\}$ is independent. It suffices to show $\{p^{j_1-1}\bar{g}'_1, \dots, p^{j_{t-1}-1}\bar{g}'_{t-1}, p^{k-1}\bar{h}_2\}$ is independent in $\bar{G}'[p]$, where $\text{order } \bar{g}'_i=p^{j_i}$. Suppose $u_1p^{j_1-1}\bar{g}'_1 + \dots + u_{t-1}p^{j_{t-1}-1}\bar{g}'_{t-1} + up^{k-1}\bar{h}_2=0$ with $(p, u)=1$. Clearly, $u_1p^{j_1-1}\bar{g}'_1 + \dots + u_{t-1}p^{j_{t-1}-1}\bar{g}'_{t-1} + up^{k-1}c\bar{g} \neq 0$. Subtraction yields $up^{k-1}(c\bar{g}-\bar{h}_2) \neq 0$. But, since $p|ca_i-b_i$, $i=3, 4, \dots, m$, $up^{k-1}(c\bar{g}-\bar{h}_2)=0$, a contradiction. Thus, in Case I, $g'=h_2$ is the desired element of G' .

Case II. Assume without loss of generality that $l_3 \neq k_3$, or that $p \nmid ca_3-b_3$. It is then easy to show that $1/p^{k_3} \in A_{\{1, m+1, m+2, \dots, n\}}^3$. Thus there exist $\Gamma_1=(0, x, ca_3/p^{k_3}, x_4, \dots, x_m, 0, 0, \dots, 0) \in G$ and (by Lemma 3) $\Gamma'_1=(0, x, ca_3/p^{k_3}, x'_4, \dots, x'_m, 0, \dots, 0) \in G'$. Applying Lemma 3 again to $\Gamma_2=cg-\Gamma_1$, we have

$$\Gamma'_2 = (ca_1/p^{k_1}, ca_2/p^{k_2} - x, 0, y'_4, \dots, y'_m, 0, 0, \dots, 0) \in G'.$$

Consider $h_2-\Gamma_1-\Gamma'_2=(0, 0, b_3/p^{l_3}-ca_3/p^{k_3}, z'_4, \dots, z'_m, 0, 0, \dots, 0)$. Since $l_3 \neq k_3$ or $p \nmid b_3-ca_3$, it follows that $1/p^{k_3} \in A_{\{1, 2, m+1, m+2, \dots, n\}}^3$. Let

$$\Gamma_3 = (0, 0, ca_3/p^{k_3}, v_4, \dots, v_m, 0, 0, \dots, 0) \in G$$

and

$$\Gamma'_3 = (0, 0, ca_3/p^{k_3}, v'_4, \dots, v'_m, 0, 0, \dots, 0) \in G'.$$

Finally, let

$$\Gamma'_4 = \Gamma'_1 + \Gamma'_2 - \Gamma'_3 = (ca_1/p^{k_1}, ca_2/p^{k_2}, 0, w'_4, \dots, w'_m, 0, 0, \dots, 0).$$

If $\{\bar{g}'_1, \dots, \bar{g}'_t, \Gamma'_4\}$ is independent, it is easy to construct the desired element $g' \in G'$. If $\{\bar{g}'_1, \dots, \bar{g}'_t, c\bar{g} - \Gamma_3\}$ is independent, the result follows from the induction hypothesis, after multiplication by a suitable integer prime to p .

Otherwise, Γ'_4 is dependent on, and Γ_3 is independent of the set $\{\bar{g}'_1, \dots, \bar{g}'_t\}$. Multiplying Γ_3 by a suitable integer d , prime to p , and applying the induction hypothesis, we obtain

$$\Gamma''_3 = (0, 0, cda_3/p^{k_3}, v''_4, \dots, v''_m, 0, 0, \dots, 0) \in G'$$

independent of $\bar{g}'_1, \dots, \bar{g}'_t$. Now $\Gamma''_3 + d\Gamma'_4$ is still independent, and we can choose e prime to p such that $g' = e(\Gamma''_3 + d\Gamma'_4)$ is the desired element.

LEMMA 5. *Let G, G', X be as in Lemma 3. Let $\bar{g}_1, \dots, \bar{g}_t$ be independent elements in the p -primary component of $\bar{G}$. Then there exist $\bar{g}'_1, \dots, \bar{g}'_t$ independent in the p -primary component of $\bar{G}'$ with $\text{order}(\bar{g}'_i) \geq \text{order}(\bar{g}_i)$.*

PROOF. The result follows by using induction on t and applying Lemma 4. We now prove the main theorem.

THEOREM A. *Let G, G' be groups of rank n with maximal independent sets Y, Y' respectively, such that $S(G, Y) = S(G', Y')$. Let F, F' be the free subgroups of G, G' generated by Y, Y' respectively. Then $G/F \cong G'/F'$.*

PROOF. As before, we identify G and G' with subgroups of Q^n containing Z^n , such that Y and Y' are identified with the set X of standard basis vectors. Write

$$G/Z^n = \sum \oplus A_p, \quad G'/Z^n = \sum \oplus A'_p$$

where A_p and A'_p are the respective p -primary components. Note that A_p and A'_p are subgroups of $(Q^n/Z^n)_p$, and therefore

$$A_p = \sum_{i=1}^r \oplus Z(p^{k_i}),$$

$$A'_p = \sum_{i=1}^{r'} \oplus Z(p^{k'_i}) \quad \text{where } 1 \leq k_i \leq \infty, \quad 1 \leq k'_i \leq \infty.$$

Applying Lemma 5, we have that $A_p \cong A'_p$ for all p .

Note that the converse to Theorem A is false. It is easy to construct nonisomorphic subgroups G and G' of $Q \oplus Q$ which contain $Z \oplus Z$ with $G/Z \oplus Z \cong G'/Z \oplus Z$. Since groups of rank two are determined up to isomorphism by their schemes, $S(G, X) \neq S(G', X')$ for any X, X' .

2. The intersection property. In this section we give a condition under which a group may be recovered from its scheme, $S(G, X)$.

DEFINITION 2. The groups A_J^i associated with (G, X) are said to satisfy the intersection property iff $A_J^i \cap A_{J'}^i = A_{J \cup J'}^i$ for all $1 \leq i \leq n$ and $i \notin J \cup J'$, where J and J' are subsets of $\{1, \dots, n\}$. In this case we will also say (G, X) satisfies the intersection property.

REMARKS. It is immediate that if G has rank ≤ 2 , (G, X) satisfies the intersection property for any maximal independent set $X \subseteq G$. If (G_i, X_i) has the intersection property for $i=1, 2, \dots, n$, then $(\sum_{i=1}^n \oplus G_i, \bigcup_{i=1}^n X_i)$ has the intersection property, where X_i' denotes the natural embedding of X_i into the direct sum. Let $v = \sum_{k=0}^\infty v_k p^k$, $w = \sum_{k=0}^\infty w_k p^k$ be p -adic units such that $1, v, w$ are rationally independent. Let $V_0 = W_0 = 0$, $V_i = \sum_{k=0}^{i-1} v_k p^k$, $W_i = \sum_{k=0}^{i-1} w_k p^k$, for $i > 1$. It can be shown using the methods of [1] that

$$G = \{(r/p^k, r/p^k V_k + t, r/p^k W_k + l) \mid (r, p) = 1, t, l \in \mathbb{Z}\}$$

is a strongly indecomposable rank 3 group; and it is easy to verify directly that (G, X) has the intersection property with $X = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$.

THEOREM B. Let (G, X) satisfy the intersection property for G a group of rank n and $X = \{x_1, \dots, x_n\}$ a maximal independent subset. Let $S(G, X) = \{A_J^i, \theta\}$. Then

$$\bar{G} = \{(\alpha_1, \dots, \alpha_n) \in \mathbb{Q}^n \mid \alpha_i \in A^i, (\alpha_i + A_j^i)^\theta = \alpha_j + A_j^i, i, j = 1, \dots, n\}$$

is isomorphic to G under the map $(\alpha_1, \dots, \alpha_n) \rightarrow \sum_{i=1}^n \alpha_i x_i$.

PROOF. We show that $(\alpha_1, \dots, \alpha_n) \in \bar{G}$ iff $\sum_{i=1}^n \alpha_i x_i \in G$. Clearly, by definition of the maps θ , $\sum_{i=1}^n \alpha_i x_i \in G$ implies $(\alpha_1, \dots, \alpha_n) \in \bar{G}$. Let $(\alpha_1, \dots, \alpha_n) \in \bar{G}$. Then $\alpha_1 \in A^1$, so $g_1 = \alpha_1 x_1 + \alpha_2' x_2 + \dots + \alpha_n' x_n \in G$ for some $\alpha_2', \dots, \alpha_n'$. But $\alpha_2' + A_1^2 = (\alpha_1 + A_2^1)^\theta = \alpha_2 + A_1^2$, so $\alpha_2 - \alpha_2' \in A_1^2$. Thus, there exists $g_2 = (\alpha_2 - \alpha_2') x_2 + \alpha_3'' x_3 + \dots + \alpha_n'' x_n \in G$. Now $g_1 + g_2 = \alpha_1 x_1 + \alpha_2 x_2 + (\alpha_3' + \alpha_3'') x_3 + \dots + (\alpha_n' + \alpha_n'') x_n \in G$. This yields $\alpha_3 - (\alpha_3' + \alpha_3'') \in A_1^3 \cap A_2^3$. Since (G, X) satisfies the intersection property, we have $\alpha_3 - (\alpha_3' + \alpha_3'') \in A_{12}^3$, and there exists $g_3 = [\alpha_3 - (\alpha_3' + \alpha_3'')] x_3 + \alpha_4''' x_4 + \dots + \alpha_n''' x_n \in G$. We continue in this way, obtaining $g_1, \dots, g_n$, such that $\sum_{i=1}^n \alpha_i x_i = g_1 + g_2 + \dots + g_n \in G$.

COROLLARY. Let (G, X) satisfy the intersection property, and let $S(G, X) = S(G', X')$. Then $G \cong G'$.

REMARK (ADDED IN PROOF). Theorem B and the above Corollary do not hold in general. There exist two nonisomorphic groups of rank 3 which have the same schemes with respect to the standard basis.

REFERENCES

1. R. A. Beaumont and R. S. Pierce, *Torsion-free rings*, Illinois J. Math. **5** (1961), 61–98. MR **26** #6212.
2. R. A. Beaumont and R. J. Wisner, *Rings with additive group which is a torsion-free group of rank two*, Acta Sci. Math. (Szeged) **20** (1959), 105–116. MR **21** #5651.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CONNECTICUT
06268