ADDITIVITY AND INDEFINITE INTEGRATION FOR MCSHANE'S P -INTEGRAL¹

C. H. SCANLON

ABSTRACT. Suppose $[a, b]$ is a closed real number interval, $A = \{(p, q]; (p, q] \subset (a, b)\}$, and U is a real valued function on $[a, b] \times A$. For $c \in (a, b)$, necessary and sufficient conditions are given for P -integrability of U on $(a, c]$ and $(c, b]$ to imply P -integrability on $(a, b]$. Suppose U is P -integrable on $(a, b]$ and $g(x) = P \int_a^x U$ for each $x \in [a, b]$. Necessary and sufficient conditions are given for g to be respectively continuous, bounded, and of bounded variation.

1. Preliminaries. Suppose $a = t_0 < t_1 < \cdots < t_n = b$ and $x_1, x_2, \dots, x_n \in [a, b]$. Then $\Pi = \{(x_1, (t_0, t_1]), \dots, (x_n, (t_{n-1}, t_n])\}$ is called a P -partition of $[a, b]$. If δ is a positive real valued function on $[a, b]$, then the statement that Π is δ -fine means that $(t_{i-1}, t_i) \subset (x_i - \delta(x_i), x_i + \delta(x_i))$ for $i = 1, \dots, n$.

The statement that U is P -integrable on $(a, b]$ means that there is a real number denoted by $P \int_a^b U$ such that if $\varepsilon > 0$ there is a positive real valued function δ on $[a, b]$ such that if $\Pi = \{(x_i, (t_{i-1}, t_i])\}_{i=1}^n$ is any δ -fine P -partition of $(a, b]$ then

$$\left| \sum_{i=1}^n U(x_i, (t_{i-1}, t_i]) - P \int_a^b U \right| < \varepsilon.$$

Variations in the following lemma can be found in ([1], [2], [3]).

LEMMA K. *If U is P -integrable on $[a, b]$ and $\varepsilon > 0$, there is a positive real valued function δ on $[a, b]$ such that if $\Pi = \{(x_i, (t_{i-1}, t_i])\}_{i=1}^n$ is any δ -fine P -partition of $[a, b]$ then*

$$\sum_{i=1}^n \left| U(x_i, (t_{i-1}, t_i]) - P \int_{t_{i-1}}^{t_i} U \right| < \varepsilon.$$

2. Additivity. E. J. McShane [1] proved that if U is P -integrable on $(a, b]$ and $c \in (a, b)$ then U is P -integrable on $(a, c]$ and $(c, b]$ and that

Received by the editors June 15, 1972.

AMS (MOS) subject classifications (1970). Primary 26A42; Secondary 28A25.

Key words and phrases. Integration, indefinite integration, additivity.

¹ This work was partially supported by Arkansas State University research funds.

© American Mathematical Society 1973

$P \int_a^b U = P \int_a^c U + P \int_c^b U$. The following theorem gives necessary and sufficient conditions for a form of the converse to hold.

THEOREM 2.1. *Suppose $c \in (a, b)$ and U is P -integrable on $(a, c]$ and $(c, b]$. Then each of the following statements is equivalent to each of the others:*

- (i) U is integrable on $(a, b]$,
- (ii) if $\varepsilon > 0$, there is a positive number γ such that if $\sup\{a, c - \gamma\} < x < c < y < \inf\{b, c + \gamma\}$, then $|U(c, (x, c]) + U(c, (c, y]) - U(c, (x, y])| < \varepsilon$, and
- (iii) if $\varepsilon > 0$ there is a positive number λ such that if

$$\sup\{a, c - \lambda\} < x < c < y < \inf\{b, c + \lambda\},$$

then $|P \int_x^c U + P \int_c^y U - U(c, (x, y])| < \varepsilon$.

PROOF. Suppose (i) is true and $\varepsilon > 0$. By Lemma K there is a positive function δ on $[a, b]$ satisfying Lemma K for $\varepsilon/3$. Then if $\sup\{a, c - \delta(c)\} < x < c < y < \inf\{c + \delta(c), b\}$,

$$(1) \quad \left| P \int_x^c U - U(c, (x, c]) \right| < \varepsilon/3,$$

$$(2) \quad \left| P \int_c^y U - U(c, (c, y]) \right| < \varepsilon/3, \quad \text{and}$$

$$(3) \quad \left| P \int_x^y U - U(c, (x, y]) \right| < \varepsilon/3.$$

Combining these inequalities yields

$$|U(c, (x, c]) + U(c, (c, y]) - U(c, (x, y])| < \varepsilon.$$

Thus (i) $\Rightarrow$ (ii). Also, (3) yields

$$\left| P \int_x^c U + P \int_c^y U - U(c, (x, y]) \right| < \varepsilon/3 < \varepsilon.$$

Thus (i) $\Rightarrow$ (iii).

Assume (ii) is true, let $\varepsilon > 0$, and let γ be the positive number from (ii) for $\varepsilon/3$. There exist positive functions δ_1 and δ_2 corresponding to the intervals $(a, c]$ and $(c, b]$, which satisfy Lemma K for $\varepsilon/3$. Let

$$\begin{aligned} \delta(x) &= \inf\{\delta_1(x), c - x\} \quad \text{if } x \in [a, c), \\ &= \inf\{\delta_1(c), \delta_2(c), \gamma\} \quad \text{if } x = c, \quad \text{and} \\ &= \inf\{\delta_2(x), x - c\} \quad \text{if } x \in (c, b]. \end{aligned}$$

Suppose $\Pi = \{(t_i, (x_{i-1}, x_i))\}_{i=1}^n$ is a δ -fine P -partition of $(a, b]$. If $c = x_j$ for some j ,

$$\begin{aligned} & \left| \sum_{i=1}^n U(t_i, (x_{i-1}, x_i)) - P \int_a^c U - P \int_c^b U \right| \\ & \leq \left| \sum_{i=1}^j U(t_i, (x_{i-1}, x_i)) - P \int_a^c U \right| + \left| \sum_{i=1}^n U(t_i, (x_{i-1}, x_i)) - P \int_c^b U \right| \\ & < \varepsilon/3 + \varepsilon/3 < \varepsilon. \end{aligned}$$

If $c \in (x_{j-1}, x_j)$ for some j ,

$$\begin{aligned} & \left| \sum_{i=1}^n U(t_i, (x_{i-1}, x_i)) - P \int_a^c U - P \int_c^b U \right| \\ & \leq \sum_{i=1}^{j-1} \left| U(t_i, (x_{i-1}, x_i)) - P \int_{x_{i-1}}^{x_i} U \right| + \left| U(c, (x_{j-1}, c]) - P \int_{x_{j-1}}^c U \right| \\ & \quad + \left| U(c, (c, x_j]) - P \int_c^{x_j} U \right| + \sum_{i=j+1}^n \left| U(t_i, (x_{i-1}, x_i)) - P \int_{x_{i-1}}^{x_i} U \right| \\ & \quad + |U(c, (x_{j-1}, c]) + U(c, (c, x_j]) - U(c, (x_{j-1}, x_j])| \\ & < \varepsilon/3 + \varepsilon/3 + \varepsilon/3 = \varepsilon. \end{aligned}$$

Thus (ii) $\Rightarrow$ (i).

Suppose (iii) is true and $\varepsilon > 0$. Let λ be the positive number from (iii) for $\varepsilon/3$. Let δ_1 and δ_2 be positive functions on $(a, c]$ and $(c, b]$ respectively which satisfy Lemma K for $\varepsilon/3$. Choose $\gamma = \inf\{\delta_1(c), \delta_2(c), \lambda\}$. Then if $\sup\{a, c - \gamma\} < x < c < y < \inf\{b, c + \gamma\}$,

$$\begin{aligned} & |U(c, (x, c]) + U(c, (c, y]) - U(c, (x, y])| \\ & \leq \left| U(c, (x, c]) - P \int_x^c U \right| + \left| U(c, (c, y]) - P \int_c^y U \right| \\ & \quad + \left| P \int_x^c U + P \int_c^y U - U(c, (x, y]) \right| \\ & < \varepsilon/3 + \varepsilon/3 + \varepsilon/3 = \varepsilon. \end{aligned}$$

Thus (iii) $\Rightarrow$ (ii).

3. Indefinite integration. Suppose U is P -integrable on $(a, b]$ and $g(x) = P \int_a^x U$ for each $x \in [a, b]$. g is called the indefinite integral of U on $[a, b]$.

THEOREM 3.1. g is bounded on $[a, b]$ if and only if there is a positive function δ on $[a, b]$ and a number M such that if $(c, d] \subset (a, b)$, $t \in [a, b]$, and $(c, d] \subset (t - \delta(t), t + \delta(t))$, then $|U(t, (c, d])| < M$.

PROOF OF NECESSITY. Suppose g is bounded and let L denote a bound of $|g|$. There is a positive function δ satisfying Lemma K for $\varepsilon=1$. Suppose $(c, d] \subset (a, b]$, $t \in [a, b]$ and $(c, d] \subset (t - \delta(t), t + \delta(t))$. Then there is a δ -fine P -partition of $(a, b]$ with $(t, (c, d])$ as a member. Thus $|U(t, (c, d]) - P \int_c^d U| < 1$ and $|U(t, (c, d])| < 1 + |P \int_c^d U| \leq 1 + |g(d) - g(c)| \leq 1 + 2L$.

PROOF OF SUFFICIENCY. Suppose U , δ , and M satisfy the hypothesis and assume g is not bounded. Let $\{t_i\}$ denote an increasing sequence of numbers in $[a, b]$ such that $|g(t_i)| > i$ for each i and let $t \in (a, b]$ denote the limit of $\{t_i\}$. If no such sequence exists, there would be a similar decreasing sequence and a similar argument would follow. By Lemma K, there exists a positive function δ' on $[a, t]$ such that if

$$\Pi = \{(t_i, (x_{i-1}, x_i))\}_{i=1}^n$$

is a δ' -fine P -partition of $(a, t]$, then

$$\sum_{i=1}^n \left| U(t_i, (x_{i-1}, x_i]) - P \int_{x_{i-1}}^{x_i} U \right| < 1.$$

Let $\gamma = \inf\{\delta(t), \delta'(t)\}$ and j a positive integer such that $0 < |t_j - t| < \gamma$ and $|g(t_i) - g(t)| > M + 1$. Since there is a δ' -fine P -partition of $(a, t]$ with $(t, (t_j, t])$ an element,

$$\left| U(t, (t_j, t]) - \int_{t_j}^t U \right| < 1,$$

thus $|g(t) - g(t_j)| - 1 < |U(t, (t_j, t])|$ and $M < |U(t, (t_j, t])|$ which contradicts the hypothesis.

THEOREM 3.2. g is continuous on $[a, b]$ if and only if for each $\varepsilon > 0$ there is a positive function δ on $[a, b]$ such that if $(c, d] \subset (a, b]$, $t \in [a, b]$ and $(c, d] \subset (t - \delta(t), t + \delta(t))$, then $|U(t, (c, d])| < \varepsilon$.

PROOF OF NECESSITY. Suppose g is on $[a, b]$ and $\varepsilon > 0$. Let δ' be positive real from the definition of uniform continuity for $\varepsilon/2$ and δ'' be the positive function from Lemma K for $\varepsilon/2$. Let $\delta(t) = \inf\{\delta', \delta''(t)\}$. Then if $t \in [a, b]$ and $(p, q] \subset (a, b]$ such that $(p, q] \subset (t - \delta(t), t + \delta(t))$, by Lemma K

$$\left| U(t, (p, q]) - P \int_p^q U \right| < \varepsilon/2,$$

$$|U(t, (p, q])| < \varepsilon/2 + |g(q) - g(p)|,$$

and since $|p - q| < \delta$, $|U(t, (p, q])| < \varepsilon$.

PROOF OF SUFFICIENCY. Suppose U satisfies the hypothesis and $\varepsilon > 0$.

Let δ be the positive function from the hypothesis for $\varepsilon/2$ and δ'' be the positive function from Lemma K for $\varepsilon/2$. Let $\delta = \inf\{\delta', \delta''\}$. Then if $x, y \in [a, b]$ and $(x, y) \subset (x - \delta(x), x + \delta(x))$,

$$\begin{aligned} |g(x) - g(y)| &= \left| P \int_x^y U \right| \\ &\leq \left| P \int_x^y U - U(x, (x, y)) \right| + |U(x, (x, y))| \\ &< \varepsilon/2 + \varepsilon/2 = \varepsilon. \end{aligned}$$

Thus g is continuous.

THEOREM 3.3. g is of bounded variation on $[a, b]$ if and only if $|U|$ is P -integrable on $(a, b]$. Furthermore, if g is of bounded variation on $[a, b]$, then the total variation of g on $[a, b]$ is $\int_a^b |U|$.

PROOF OF NECESSITY. Suppose g is of bounded variation and let Vg denote the total variation of g on $[a, b]$ and let $\varepsilon > 0$. There is a partition $a = x_0 < x_1 < \cdots < x_n = b$ such that $0 \leq Vg - \sum_{i=1}^n |g(x_i) - g(x_{i-1})| < \varepsilon/2$.

Let

$$\begin{aligned} \delta'(t) &= \inf\{x_i - t, t - x_{i-1}\} \quad \text{if } t \in (x_{i-1}, x_i), \\ &= \frac{1}{2} \inf\{x_i - x_{i-1}, x_{i+1} - x_i\} \quad \text{if } t = x_i \text{ for } i = 1, \dots, n-1, \\ &= \frac{1}{2}(x_n - x_{n-1}) \quad \text{if } t = x_n, \\ &= \frac{1}{2}(x_1 - x_0) \quad \text{if } t = x_0. \end{aligned}$$

Let δ'' be the positive function from Lemma K for $\varepsilon/2$ and $\delta = \inf\{\delta', \delta''\}$.

Suppose $\Pi = \{(t_i, (y_{i-1}, y_i))\}_{i=1}^m$ is a δ -fine P -partition of $(a, b]$. Then $a = y_2 < y_1 < \cdots < y_m = b$ is a refinement of $a = x_0 < x_1 < \cdots < x_m = b$ and thus $0 \leq Vg - \sum_{i=1}^m |g(y_i) - g(y_{i-1})| < \varepsilon/2$ and $0 \leq Vg - \sum_{i=1}^m |P \int_{y_{i-1}}^{y_i} U| < \varepsilon/2$. It follows from Lemma K that

$$\left| \sum_{i=1}^m |U(t_i, (y_{i-1}, y_i))| - \sum_{i=1}^m \left| P \int_{y_{i-1}}^{y_i} U \right| \right| < \frac{\varepsilon}{2}.$$

Combining inequalities yields

$$\left| \sum_{i=1}^m |U(t_i, (y_{i-1}, y_i))| - Vg \right| < \varepsilon.$$

Therefore $|U|$ is P -integrable and $P \int_a^b |U| = Vg$.

PROOF OF SUFFICIENCY. Suppose $|U|$ is P -integrable and $\varepsilon > 0$. Let δ'' be a positive function which satisfies Lemma K for both U and $|U|$ for $\varepsilon/2$.

Let $a = x_0 < x_1 < \cdots < x_n = b$ be any partition of $[a, b]$. Define δ as in the necessary proof and let $\Pi = \{(t_i, (y_{i-1}, y_i))\}_{i=1}^m$ be any δ -fine P -partition of $(a, b]$.

Lemma K yields

$$\sum_{i=1}^m \left| U(t_i, (y_{i-1}, y_i]) - P \int_{y_{i-1}}^{y_i} U \right| < \frac{\varepsilon}{2}$$

and thus

$$\sum_{i=1}^m ||U(t_i, (y_{i-1}, y_i])| - |g(y_i) - g(y_{i-1})|| < \frac{\varepsilon}{2}.$$

Also, Lemma K gives

$$\sum_{i=1}^m \left| |U(t_i, (y_{i-1}, y_i])| - P \int_{y_{i-1}}^{y_i} |U| \right| < \frac{\varepsilon}{2}.$$

Combining inequalities yields

$$\left| \sum_{i=1}^m |g(y_i) - g(y_{i-1})| - P \int_a^b |U| \right| < \varepsilon.$$

Therefore it follows from the definition of bounded variation that g is of bounded variation and that Vg is $P \int_a^b |U|$.

REFERENCES

1. E. J. McShane, *A Riemann-type integral that includes Lebesgue-Stieltjes, Bochner and stochastic integrals*, Mem. Amer. Math. Soc. No. 88 (1969). MR 42 #436.
2. A. Kolmogoroff, *Untersuchungen über den Integralbegriff*, Math. Ann. 103 (1930), 654-696.
3. Ralph Henstock, *Theory of integration*, Butterworths, London, 1963. MR 28 #1274.

DEPARTMENT OF MATHEMATICS, ARKANSAS STATE UNIVERSITY, STATE UNIVERSITY,
ARKANSAS 72467