K-THEORY OF COMMUTATIVE REGULAR RINGS

ANDY R. MAGID

ABSTRACT. Pierce's representation of a commutative regular ring as a sheaf of fields is used to compute the K -theory of the ring: K_1 is units (Robert's Theorem) and K_2 is generated by symbols.

Pierce [4] shows how a commutative regular (in the sense of von Neumann) ring R can be represented as the ring of all global sections of a sheaf $\mathcal{R}$ of fields over a zero-dimensional compact space X . Because of the zero-dimensionality of X , data over the stalks of $\mathcal{R}$ can be extended to similar global data, and thus much of the theory of R follows immediately from the corresponding theory for fields. This brief note uses this technique to compute the K -theory of R : $K_0(R)$ is the ring $C(X, \mathbb{Z})$ of continuous functions from X to the integers $\mathbb{Z}$ (this is due to Pierce [4, 16.4, p. 67]); $K_1(R)$ is the group of units of R (this is due to Roberts [5, p. 425]); and $K_2(R)$ is generated by the universal Steinberg symbols. The proofs use only the fact that the K_i are functors of finite type and hence apply to other such functors. This gives new proofs of earlier results on the Brauer group [2, 1.10, p. 117].

We begin by recalling Pierce's general construction: to each commutative ring R is associated a compact, zero-dimensional Hausdorff space $X(R)$ and a sheaf $\mathcal{R}$ of connected rings on $X(R)$ such that $R = \Gamma(X(R), \mathcal{R})$ [4, 4.4, p. 17].

When R is regular, this is just the usual sheaf of local rings on $\text{Spec}(R)$ [6, 2.4].

Now suppose F is an abelian group valued functor on the category of commutative rings. Then $F \circ \mathcal{R}$ is a presheaf of abelian groups on $X(R)$. We denote its associated sheaf by $\#F$. Further study of $\#F$ requires the following hypothesis.

DEFINITION (SEE [1, 1.5, p. 24]). F is of *finite type* if F commutes with finite products and arbitrary direct limits.

PROPOSITION 1. *In the above notation, suppose F is of finite type. Then the group of global sections of $\#F$ is isomorphic to $F(R)$ and for each x in $X(R)$, the stalk $(\#F)_x$ is isomorphic to $F(R_x)$.*

Received by the editors October 6, 1972.

AMS (MOS) subject classifications (1970). Primary 13D15; Secondary 13A30.

Key words and phrases, K -theory, von Neumann regular ring, functor of finite type.

© American Mathematical Society 1973

PROOF. For the first part, see [2, 1.9, p. 117]. For the second part, $(\#F)_x = F \circ \mathcal{R}_x = \text{dir lim } F(\Gamma(U, \mathcal{R})) = F(\text{dir lim } \Gamma(U, \mathcal{R}))$ (since F is of finite type) and $\text{dir lim } \Gamma(U, \mathcal{R}) = R_x$, the direct limit being over open neighborhoods of x .

We note the proposition applies to the following functors of finite type: K_i , $i=0, 1, 2$, K_2 being in the sense of Milnor [3, p. 40], the Picard group, the multiplicative group G_m and the Brauer group. (Regarding the latter, see also [2, 1.10, p. 117].) We also will consider the universal Steinberg symbol functor [3, 11.1, p. 93] defined as follows: for any commutative ring T , $Us(T)$ is the multiplicative abelian group with one generator (a, b) for each pair of units a, b of T and relations forcing (a, b) to be bimultiplicative and $(a, b) = 1$ if $a + b = 1$. Clearly Us is a functor of finite type.

PROPOSITION 2. *Let R be a commutative regular ring and $F \rightarrow G$ a natural transformation of functors of finite type from commutative rings to abelian groups which is an isomorphism when the rings are fields. Then $F(R) \rightarrow G(R)$ is also an isomorphism.*

PROOF. The natural transformation $F \rightarrow G$ induces a sheaf morphism $f: \#F \rightarrow \#G$. For each x in $X(R)$, R_x is a field [4, 10.3, p. 41] and we have a commutative diagram

$$\begin{array}{ccc} (\#F)_x & \longrightarrow & (\#G)_x \\ \downarrow & & \downarrow \\ F(R_x) & \longrightarrow & G(R_x) \end{array}$$

The vertical maps are isomorphisms by Proposition 1 and the lower horizontal map is an isomorphism by hypothesis. Thus the upper map is an isomorphism. Thus f , being an isomorphism at each stalk, is an isomorphism and hence induces an isomorphism of global sections. But, by Proposition 1 again, these global sections are $F(R)$ and $G(R)$ respectively.

COROLLARY 3. *Let R be a commutative regular ring. Then:*

- (a) $K_0 R = C(X(R), Z)$,
- (b) $K_1 R = G_m(R)$,
- (c) $K_2 R = Us(R)$.

PROOF. The transformation $K_0 \rightarrow C(X(\), Z)$ is given by the rank homomorphism, the transformation $K_1 \rightarrow G_m$ is given by the determinant and the transformation $Us \rightarrow K_2$ is given by symbols [3, p. 74]. These are all isomorphisms for fields, the first two being classical and the third a theorem of Matsumoto [3, 11.1, p. 93]. Thus Proposition 2 applies and the corollary follows.

Roberts also computes relative K_1 for commutative regular rings [5, p. 425]. We outline another approach to his result.

LEMMA 4. *Let R be a commutative regular ring, I an ideal of R . Then $G_m(R) \rightarrow G_m(R/I)$ is onto.*

PROOF. Let r in R go to a unit of R/I , and let $Rr = Re$ where e is idempotent. I is the intersection of the maximal ideals M containing it, and if I is contained in M , r , being a unit modulo I , is not, so e is not in M and hence $1 - e$ is. This holds for all M containing I , so $1 - e$ is in I . Let $s = 1 - e + r$. If s is in the maximal ideal M , then if $e \in M$, $r \in M$ so also $1 - e \in M$, which is impossible, so e is not in M and $1 - e$ is in M , hence $r \in M$, hence $e \in M$, again an impossibility. Thus s is in no maximal ideal, hence is a unit congruent to r modulo I .

COROLLARY 5. *Let R be a commutative regular ring, I an ideal of R . Then $K_1(R, I) = \text{Kernel}(G_m(R) \rightarrow G_m(R/I))$.*

PROOF. We have an exact sequence [3, 6.2, p. 54]

$$K_2R \rightarrow K_2R/I \rightarrow K_1(R, I) \rightarrow K_1R \rightarrow K_1R/I$$

which by Corollary 5 becomes

$$Us(R) \rightarrow Us(R/I) \rightarrow K_1(R, I) \rightarrow G_m(R) \rightarrow G_m(R/I).$$

By Lemma 4 the first map is onto, and the result follows by exactness.

Proposition 1 gives information about rings of continuous functions $C(X, T)$ where X is a compact zero-dimensional topological space and T a commutative ring with the discrete topology. However, a more elementary argument suffices in this special case, which we now record.

PROPOSITION 6. *Let $R = C(X, T)$ be as above and let F be a functor of finite type. Then $F(R)$ is isomorphic to $C(X, F(T))$ (here $F(T)$ carries the discrete topology).*

PROOF. A finite partition of X is a cover of X by finitely many disjoint open subsets. The partitions of X are partially ordered by refinement and $X = \text{proj lim } P$ where P ranges over the partitions of X . Thus $C(X, T) = \text{dir lim } C(P, T)$, and $C(P, T)$ is a finite product of copies of T . Then since F is of finite type, $F(C(X, T)) = \text{dir lim } F(C(P, T)) = \text{dir lim } C(P, F(T)) = C(X, F(T))$.

By the Stone representation theorem, Boolean rings are of the type treated in Proposition 6 (where T is the field of two elements). Thus the K -theory of Boolean rings may be computed:

COROLLARY 7. *Let R be a Boolean ring. Then: $K_0R = C(X(R), \mathbb{Z})$ and K_1 and K_2 of R are trivial.*

PROOF. The final assertion follows from Steinberg's result [3, 9.9, p. 75] that K_2 of a finite field is trivial.

One may also use Proposition 6 to compute K -theory of rings of continuous integer valued functions, and also Brauer groups of such rings and Boolean rings as in [2, 1.12, p. 118].

REFERENCES

1. M. Artin, *Algebraic approximation of structures over complete local rings*, Inst. Hautes Études Sci. Publ. Math. **36** (1969), 23–58. MR **42** #3087.
2. A. Magid, *Pierce's representation and separable algebras*, Illinois J. Math. **15** (1971), 114–121. MR **42** #7713.
3. J. Milnor, *Introduction to algebraic K-theory*, Ann. of Math. Studies, no. 72, Princeton Univ. Press, Princeton, N.J., 1971.
4. R. Pierce, *Modules over commutative regular rings*, Mem. Amer. Math. Soc. No. 70 (1967). MR **36** #151.
5. L. Roberts, K_1 of a commutative von Neumann regular ring, Proc. Amer. Math. Soc. **32** (1972), 425–426.
6. O. Villamayor and D. Zelinsky, *Galois theory with infinitely many idempotents*, Nagoya Math. J. **35** (1969), 83–98. MR **39** #5555.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OKLAHOMA 73069