ERRATUM TO VOLUME 27

R. B. Hughes, *Zero sets of functions from non-quasi-analytic classes*, Proc. Amer. Math. Soc. **27** (1971), 539–542.

Page 540, line 22, reads “ $\cdots$ and $h_2(t, k)$ in $C^\infty(R)$ such that $\cdots$ ”.

It should read

“ $\cdots$ and $h_2(t, k)$, h_1 in $C^\infty([0, \infty))$ and h_2 in $C^\infty((-\infty, 4\lambda])$ such that $\cdots$ ”.

DEPARTMENT OF MATHEMATICS, BOISE STATE COLLEGE, BOISE, IDAHO 83707

ERRATUM TO VOLUME 31

R. J. Plemmons and R. E. Cline, *The generalized inverse of a nonnegative matrix*, Proc. Amer. Math. Soc. **31** (1972), 46–50.

The first statement of Theorem 2 in the above article is incorrect. In particular one can construct square matrices $A \geq 0$ having no zero rows or columns with the Moore-Penrose generalized inverse $A^+ \geq 0$, which is not diagonally equivalent to a doubly stochastic matrix. As an example take

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

Then $A^+ = \frac{1}{2}A^T$, while A is clearly not diagonally equivalent to a doubly stochastic matrix. This item is not directly related to the main topic of the paper and the mistake affects no other results in the paper.

The authors wish to thank Professor D. J. Hartfiel for informing them of the error.

DEPARTMENTS OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF TENNESSEE,
KNOXVILLE, TENNESSEE 37916

© American Mathematical Society 1973