AN EXTREMAL PROPERTY OF SOME CAPACITARY MEASURES IN E_n^1

BURGESS DAVIS AND JOHN L. LEWIS

ABSTRACT. The capacitary measure on an arc of the circle is known (via conformal mapping) to be that measure of a class of measures which has the largest potential at certain points of the plane. Here it is shown that the analogous result is true in E_n .

1. Introduction. Let E_n be Euclidean n space, $n \geq 3$; denote vectors $(x_1, \dots, x_n) \in E_n$ by x ; and let $|x|$ be the norm of x .

Put $S = \{x \in E_n : |x| = 1\}$ and for a given positive c , $1 \leq c < \infty$, let $H(c)$ denote the class of those positive measures μ on S of total mass 1 which satisfy

$$(1.1) \quad \phi(x) \leq c \quad \text{for each } x \in E_n.$$

Here $\phi(x) = \int_S |x - y|^{2-n} d\mu(y)$ is the potential of μ .

Let $x \in E_n$, $|x| \neq 0$, $|x| \neq 1$. In this paper we characterize the measure in $H(c)$ which has the largest potential at x . The corresponding problem, where now ϕ is a logarithmic potential, occurs in E_2 in the study of the class S^* of starlike univalent functions f in $\{|x| < 1\}$, normalized by $f(0) = 0$, $f'(0) = 1$, and satisfying $|f| \leq e^{2c}$. Indeed, if ϕ is given, then the function f defined by $\log(|f(z)|/|z|) = 2\phi(z)$, $|z| < 1$, $f'(0) = 1$, is in S^* . Conversely, given $f \in S^*$, $\frac{1}{2} \log(|f(z)|/|z|)$ is a potential whose associated measure is in $H(c)$. Hence the problem of characterizing the measure in $H(c)$ which has largest potential at x is equivalent to finding the function f in S^* whose modulus is largest at x . Due to this equivalence, in E_2 the measure can be characterized by using conformal mapping. The measure has the same form in E_n . That is

THEOREM 1. *Let $x \in E_n$, $|x| \neq 0$, $|x| \neq 1$, $n \geq 3$. Let λ be such that the cap $\{y : y \in S, |x - y| \leq \lambda\}$ has capacity $1/c$. Then if e is the capacitary measure on this cap, the measure ce has a greater potential at x than any other measure in $H(c)$.*

Received by the editors September 11, 1972.

AMS (MOS) subject classifications (1970). Primary 31B15.

Key words and phrases. Capacity measure, potential.

¹ Research partially supported by NSF Grant GP 19222.

With minor changes our method provides a new proof of the corresponding result in E_2 .

We remark that T. J. Suffridge in E_2 [3] and J. L. Lewis in E_3 [2] have considered a related problem for potentials bounded below in $\{x: |x| \leq 1\}$ by a constant.

2. Several lemmas. Given α , $0 < \alpha \leq \pi$, let $C_\alpha = \{x \in S: x_1 > \cos \alpha\}$. Let ∂C_α denote $\{x \in S: x_1 = \cos \alpha\}$ and $\bar{C}_\alpha = \partial C_\alpha \cup C_\alpha$. Let μ_α be the capacity measure on $\bar{C}_\alpha$ and let P_α be its potential. Then

$$(2.1) \quad P_\alpha(x) = \int_{\bar{C}_\alpha} |x - y|^{2-n} d\mu_\alpha(y).$$

P_α is continuous on $E_n \cup \{\infty\}$, harmonic in $E_n - \bar{C}_\alpha$, 1 on $\bar{C}_\alpha$ and 0 at ∞ , and is superharmonic in E_n . For fixed c , $1 \leq c < \infty$, let $\alpha_0 = \alpha_0(c)$ be such that $\Phi = cP_{\alpha_0}$ satisfies $\Phi(0) = 1$, that is the potential of $c\mu_{\alpha_0}$ at 0 is 1. The existence of α_0 is easily shown. We first prove

LEMMA 1. *If ϕ is the potential of a measure in $H(c)$ then for $0 < \alpha \leq \pi$*

$$\int_{\bar{C}_\alpha} \phi d\mu_\alpha \leq \int_{\bar{C}_\alpha} \Phi d\mu_\alpha.$$

PROOF. The lemma is trivial if $\alpha \leq \alpha_0$ since $\Phi = c$ on $\bar{C}_{\alpha_0}$. If $\alpha > \alpha_0$ we interchange the order of integration in $\int_{\bar{C}_\alpha} \phi d\mu_\alpha$ and obtain for ϕ with μ in $H(c)$ that

$$\int_{\bar{C}_\alpha} \phi d\mu_\alpha = \int_S P_\alpha(y) d\mu(y) \leq \int_S d\mu(y) = 1.$$

Similarly we have

$$\int_{\bar{C}_\alpha} \Phi d\mu_\alpha = \int_S cP_\alpha(y) d\mu_{\alpha_0} = \int_{\bar{C}_{\alpha_0}} c d\mu_{\alpha_0} = 1.$$

Next we note that

- (i) P_α has continuous one sided normal derivatives on C_α .
- (ii) $\mu_\alpha(C_\theta)$ is a continuous function of θ , $0 < \theta \leq \pi$.
- (i) is easily verified using the Kelvin transformation to map $\{x: |x| < 1\}$ onto a halfspace (see Helms [1, pp. 36, 37]), and the reflection principle.
- (ii) follows from the fact that μ_α is capacity measure on $\bar{C}_\alpha$. Let σ denote Lebesgue measure on S and σ_n the Lebesgue measure of S . Differentiating (2.1) under the integral sign we find for rx , $0 < r < 1$, $|x| = 1$, that

$$(2.2) \quad r^{2-n/2} \frac{\partial}{\partial r} [r^{n/2-1} P_\alpha(rx)] = \left(\frac{n}{2} - 1\right) \int_{\bar{C}_\alpha} \frac{1 - r^2}{|rx - y|^n} d\mu_\alpha(y).$$

The last integral is a constant multiple, $(n/2-1)\sigma_n$, of the Poisson integral with boundary measure μ_α evaluated at rx . It follows from this fact that

$$(2.3) \quad (n/2-1)^{-1}\sigma_n^{-1} \lim_{r \rightarrow 1} r^{2-n/2}(\partial/\partial r)[r^{n/2-1}P_\alpha(rx)] = d\mu_\alpha(x)/d\sigma$$

almost everywhere with respect to σ . Since (i) is true we deduce that μ_α is absolutely continuous with respect to σ on C_α and thereupon that equality holds in (2.3) whenever $x \in \partial C_\alpha$. Since we also have (ii) it follows that μ_α is absolutely continuous on $\bar{C}_\alpha$.

We observe from the boundary values of P_α and (2.3) that the values of $d\mu_\alpha(x)/d\sigma$ depend only on the x_1 coordinate of x . Hence if we define θ by $x_1 = \cos \theta$, $0 \leq \theta < \alpha$, and put $h_\alpha(\theta) = d\mu_\alpha(x)/d\sigma$, $0 \leq \theta < \alpha$, then h is well defined. With this notation we prove

LEMMA 2. *If $0 < \alpha \leq \pi$, then h_α is a nonnegative nondecreasing function of θ on $[0, \alpha]$ which is not everywhere 0.*

PROOF. Let x_0 be that point on S with $x_1 = 1$, $x_i = 0$ if $i \neq 1$. We first show for $0 < r < 1$ that

$$(2.4) \quad P_\alpha(ry) \leq P_\alpha(rx)$$

when $x, y \in C_\alpha$ and $|x - x_0| < |y - x_0|$. To prove (2.4) let K be the hyperplane through the origin for which $\tilde{x} = y$, where $\tilde{x}$ denotes the reflection of x with respect to K . Let G denote the halfspace bounded by K which contains x . Define f in $K \cup G$ by $f(z) = P_\alpha(\tilde{z})$, $z \in K \cup G$. Then $f - P_\alpha \leq 0$ on $K \cup (C_\alpha \cap G)$. Since $f - P_\alpha$ is bounded and harmonic in $G - \bar{C}_\alpha$ it follows from the maximum principle that $f \leq P_\alpha$ in G and consequently (2.4) is true.

Using (2.4) we get

$$\int_0^s \frac{\partial}{\partial r} [r^{n/2-1}P_\alpha(ry)] dr \leq \int_0^s \frac{\partial}{\partial r} [r^{n/2-1}P_\alpha(rx)] dr$$

for $0 < s \leq 1$. Since equality holds for $s = 1$ it follows that

$$\lim_{r \rightarrow 1} (\partial/\partial r)[r^{n/2-1}P_\alpha(rx)] \leq \lim_{r \rightarrow 1} (\partial/\partial r)[r^{n/2-1}P_\alpha(ry)].$$

Using (2.3) we conclude that h_α is nondecreasing on $[0, \alpha]$. Finally, h_α is somewhere positive since $\mu_\alpha(C_\alpha) > 0$.

3. Proof of Theorem 1. It clearly suffices to prove Theorem 1 for points x of the form $x = tx_0$, $0 < t < \infty$, $t \neq 1$. Let Q be the Poisson kernel. Let $\phi \in H(c)$, $\phi \neq \Phi$, and suppose by way of contradiction that $\Phi(tx_0) \leq \phi(tx_0)$. Put

$$(3.1) \quad F(\theta) = \int_{C_\theta} Q(tx_0, y)[\Phi(y) - \phi(y)] d\sigma(y), \quad 0 < \theta \leq \pi.$$

Then $\Phi(tx_0) - \phi(tx_0) = F(\pi)$, and so the following lemma immediately gives Theorem 1.

LEMMA 3. *If $\phi \neq \Phi$, then $F(\alpha) > 0$, $\alpha_0 \leq \alpha \leq \pi$.*

PROOF. We observe $F(\alpha) \geq 0$, $0 < \alpha < \alpha_0$, and $F(\alpha_0) > 0$, since $\Phi = c$ on $\bar{C}_{\alpha_0}$. Hence if Lemma 3 is false there exists δ , $\alpha_0 < \delta \leq \pi$, such that $F(\alpha) > 0$, $\alpha_0 \leq \alpha < \delta$, and $F(\delta) = 0$.

Define $\hat{Q}$ on $[0, \pi]$ by $\hat{Q}(\theta) = Q(tx_0, y)$ where y satisfies $y_1 = \cos \theta$, and define $g_\delta(\theta) = h_\delta(\theta)/\hat{Q}(\theta)$. Since $\hat{Q}$ is strictly decreasing on $[0, \pi]$, Lemma 2 implies g_δ is nondecreasing and nonconstant on $[\alpha_0, \delta)$.

If $\alpha_0 < \beta < \delta$ then

$$(3.2) \quad \int_{C_\beta} (\Phi - \phi) d\mu_\delta = \int_0^\beta g_\delta(\theta) dF(\theta) = F(\beta)g_\delta(\beta) - \int_0^\beta F(\theta) dg_\delta(\theta).$$

The left-hand integral approaches

$$\int_{C_\delta} (\Phi - \phi) d\mu_\delta \quad \text{as } \beta \rightarrow \delta,$$

since μ_δ is absolutely continuous with respect to σ . Also

$$\int_\beta^\delta g_\delta(\theta) |dF(\theta)| \geq F(\beta)g_\delta(\beta)$$

and so $\lim_{\beta \rightarrow \delta} F(\beta)g_\delta(\beta) = 0$. Taking the limit as $\beta \rightarrow \delta$ in (3.2), we get

$$\int_{C_\delta} (\Phi - \phi) d\mu_\delta = - \int_0^\delta F(\theta) dg_\delta(\theta).$$

The right-hand side is negative since g_δ is nondecreasing and nonconstant on $[0, \delta]$ and F is positive on $[\alpha_0, \delta)$. However by Lemma 1 the left-hand integral is nonnegative. We have reached a contradiction. Hence Lemma 3 is true.

4. **Remark.** We note that Φ also solves the corresponding minimum problem. That is

$$(4.1) \quad \Phi(-tx_0) < \min_{|x|=t} \phi(x),$$

when ϕ is a potential of a measure on $H(c)$, ϕ is not a rotation of Φ , $0 < t < \infty$, and $t \neq 1$. The proof is exactly the same except we substitute $-Q(-tx_0, y)$ for $Q(tx_0, y)$ in (3.1). It appears likely that (4.1) is also true for $t=1$, although we have not been able to prove this.

REFERENCES

1. L. L. Helms, *Introduction to potential theory*, Pure and Appl. Math., vol. 22, Interscience, New York, 1969. MR 41 #5638.
2. J. L. Lewis, *A potential theory problem in three space* (to appear).
3. T. J. Suffridge, *A coefficient problem for a class of univalent functions*, Michigan Math. J. 16 (1969), 33–42. MR 39 #1646.

STATISTICS CENTER, RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY 08903

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, LEXINGTON, KENTUCKY 40506